

## Optical Photometry of the Eclipsing LMC Supersoft Source CAL 87

K.A. Southwell, P.A. Charles

*Department of Astrophysics, Keble Road, Oxford, OX1 3RH, UK*

D. O'Donoghue

*Department of Astronomy, University of Cape Town, South Africa*

C. Alcock, R.A. Allsman, D. Alves, T.S. Axelrod, D.P. Bennett, K.H. Cook, K.C. Freeman, K. Griest, J. Guern, M.J. Lehner, S.L. Marshall, B.A. Peterson, M.R. Pratt, P.J. Quinn, A.W. Rodgers, C.W. Stubbs, W. Sutherland, D.L. Welch

*The MACHO Collaboration*

**Abstract.** We present optical photometry of the eclipsing supersoft source, CAL 87. We find the eclipse structure to be stable over  $\sim 4$  y, derive an improved ephemeris of  $T_o = \text{HJD } 2450111.5144(3) + 0.442674(7)E$ , and see new structure in the light curve morphology.

### 1. MACHO Project Photometry

We show in Fig. 1 optical photometry of the eclipsing supersoft source, CAL 87, obtained via the MACHO project from 1992 Aug – 1996 May (see Alcock et al. 1996). The broad primary eclipse is  $\sim 0.2$  mag deeper in the blue, and




Figure 1. The blue and red MACHO light curves of CAL 87. The data are folded and averaged into 50 phase bins.

a shallower secondary dip is seen in both passbands at  $\phi \sim 0.5$ . The latter probably results from obscuration of the irradiated face of the donor star (van den Heuvel et al. 1992). The  $B - R$  colour (not plotted) shows pronounced




Figure 2. SAAO high speed white light curve of CAL 87. Note the obvious structure at  $\phi \sim 0.82 - 0.87$  and  $\phi > 1.1$ .

reddening through the primary eclipse, and also marginal evidence for a slight reddening centred on  $\phi \sim 0.55$ . Examination of the yearly data-sets from 1992-1996 failed to reveal any significant changes in the orbital period or overall morphology, although the step in the egress at  $\phi \sim 1.1$  (see also Fig. 2) sometimes seemed more pronounced. We derive  $P = 0.442676 \pm 0.000007$  d from the complete dataset in each filter, consistent with previous measurements (e. g. Schmidtke et al. 1993).

## 2. High speed photometry

Fig. 2 displays a white light curve of a single eclipse, consisting of continuous 10 – 20 s integrations, obtained with the UCT CCD and SAAO 1.9 m telescope. We derive an ephemeris of  $T_o = \text{HJD } 2450111.5144(3)$  for the time of minimum light. The extended ingress and kink at  $\phi \sim 0.82 - 0.87$  may be explained by variable obscuration by the thickened accretion disc rim (Callanan et al. 1989; Armitage & Livio 1996; Alcock et al. 1996). A turn-over is observed in the egress for  $\phi > 1.1$ . This is similar to the variability seen in Z Cha (Cook & Warner 1984), caused by the presence of a hot spot on the disc. In CAL 87, an analogous component may be the strongly irradiated inner-surface of the accretion disc bulge.

## References

- Alcock, C. et al. 1996, MNRAS, submitted  
 Armitage, P. J. & Livio, M. 1996, ApJ, in press  
 Callanan, P. J., Machin, G., Naylor, T. & Charles, P. A. 1989, MNRAS, 241, 37P  
 Cook, M. C. & Warner, B. 1984, MNRAS, 207, 705  
 Schmidtke, P. C., McGrath, T. K., Cowley, A. P. & Frattare, L. M. 1993, PASP, 105, 863  
 van den Heuvel, E. P. J., Bhattacharya, D., Nomoto, K. & Rappaport, S. A. 1992, A&A, 262, 97