

EPP0677

Comorbidities of anxiety disorders in bipolar patients: therapeutic complexityS. Bahetta^{1*} and N. El moussaoui²Psychiatric hospital arrazi, Sale and ²Psychiatric hospital arrazi, Morocco, Morocco

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.973

Introduction: Numerous clinical and epidemiological studies show that the rate of comorbidity of anxiety disorders is high in bipolar patients compared to the general population. This is associated with a poorer prognosis, poorer functioning and higher suicidal risk. Anxiety comorbidity should therefore be carefully investigated.

Objectives: Our main objectives are to explore the therapeutic complexity of anxiety disorders in patients with bipolar disorder. To investigate the existence of psycho-pathological links and vulnerabilities between bipolar disorder and anxiety disorders.

Methods: through a clinical vignette and a review of the existing literature on the comorbidity of anxiety disorders and bipolar disorders, and the resulting therapeutic issues

Results: Anxiety comorbidity is quite common in the bipolar population. In the American National Comorbidity Survey (NCS), lifetime comorbidity is close to 90%. Two recent French clinical studies show the existence of at least one anxiety disorder in approximately 25% of bipolar subjects (24% and 27.2%), which will have an impact on the course of the bipolar disorder, with a particular increase in the risk of suicide, hence the importance of adequate treatment. This treatment faces two obstacles: the risk of manic episodes under antidepressants and the risk of dependence on benzodiazepines. Emphasis is also placed on non-drug approaches, including cognitive-behavioural and psycho-educational therapies.

Conclusions: Anxiety comorbidity is not without consequence on the evolution of bipolar disorder. Its particularly high prevalence means that it cannot be neglected or ignored in current practice.

Disclosure of Interest: None Declared

EPP0678

Work Stress, Overcommitment and Somatization among Early Childhood Professionals during COVID-19: A Cross-sectional Study

S. Gritzka*, P. Angerer and M. Diebig

Institut für Arbeits-, Sozial- und Umweltmedizin, Heinrich-Heine-Universität Düsseldorf, Düsseldorf, Germany

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.974

Introduction: In addition to posing major public health challenges, the COVID-19 pandemic has affected psychological and physical stressors in workplaces as well as strain by workers worldwide. Even prior COVID-19, the working conditions of early childhood professionals (ECPs) were described as critical leading to (psycho) somatic complaints. Despite the high societal relevance of ECPs and the potential increase in stressors caused by COVID-19, there is

a lack of empirical evidence to which extent work-related demands had an adverse effect on ECPs health during COVID-19.

Objectives: We aimed to obtain updated prevalence rates of somatic symptoms of ECPs as well as unfavourable working conditions inducing stress (in terms of an effort-reward imbalance; ERI) in childcare. The overall objective was to establish associations between work stress and health during the early phase of the pandemic. Given the tendency of ECPs to overcommit themselves, we further explored the potential moderating role of overcommitment.

Methods: Between June 2020 and May 2021, questionnaire data was collected using validated instruments (i.e., ERI, PHQ-15) from ECPs in childcare centers as well as from family providers ($N = 1.009$). ECPs were informed about the study and contacted through the youth welfare office. Multiple logistic regression tested the influence of ERI and overcommitment on the severity of somatic symptoms. It was controlled for age, gender as well as leadership position. SPSS 28.0 was used to analyze the data.

Results: The ERI ratio of the sample averaged 1.28 ($SD = 0.45$), with 72.7% of subjects exceeding the critical cut-off value > 1 , indicating a gratification crisis. The averaged sum score of the PHQ-15 was 8.99 ($SD = 5.43$). Based on a PHQ-15 cut-off ≥ 10 , the overall prevalence of somatization at a moderate to high level is estimated to be 44.6%. The mean overcommitment score was 15.14 ($SD = 3.55$) and 23.4% of ECPs were in the highest overcommitment tertile. ERI ($OR = 4.12$, 95% $CI: 2.73 - 6.22$) and overcommitment ($OR = 5.20$, 95% $CI: 2.17 - 12.47$) were associated with an increased likelihood of greater severity of somatic symptoms. Yet, the interaction effect between both predictors remained non-significant. Being female and having no leadership position were predictors for a moderate to high level of somatization.

Conclusions: The results demonstrate the high relevance of work stress for somatic health among ECPs in the midst of a pandemic. Given the high overall prevalence of somatic symptoms and the female-dominated work sector, ECPs may be at high risk for somatoform disorders. There remains a strong need for action to reduce work-related stress in order to decrease the somatic symptom burden. Large-scale and longitudinal studies are needed to further investigate coping with and persistence of somatic symptoms among this occupational group.

Disclosure of Interest: None Declared

EPP0679

The effect psychobiotics on anxiety symptomsY. Denysov^{1*}, G. Putyatin¹, S. Moroz² and V. Semenikhina²

¹Psychiatry, psychotherapy, addictology and medical psychology, Donetsk national medical university, Kropyvnytskyi and ²Psychiatry, Municipal Enterprise "Dnipro Multiprofile Clinical Hospital for the Provision of Psychiatric Care of Dnipropetrovsk Regional Council, Dnipro, Ukraine

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.975

Introduction: Psychobiotics are a group of probiotics that affect the central nervous system related functions and behaviors mediated by the gut-brain-axis via immune, humoral, neural, and metabolic pathways to improve not only the gastrointestinal function but also the antidepressant and anxiolytic capacity.