

An outbreak of multidrug-resistant *Salmonella enterica* subsp. *enterica* serotype Typhimurium, DT104L linked to dried anchovy in Singapore

M. L. LING¹*, K. T. GOH², G. C. Y. WANG¹, K. S. NEO² AND T. CHUA²

¹Department of Pathology, Singapore General Hospital, Singapore

²Quarantine and Epidemiology Department, Ministry of the Environment, Singapore

(Accepted 2 October 2001)

SUMMARY

Multidrug-resistant *Salmonella enterica* subsp. *enterica* serotype Typhimurium DT104L was first reported in Singapore from mid-July to mid-October 2000. *Salmonella* strains isolated from clinical laboratories were submitted to a reference laboratory for serotyping, phage-typing and pulsed-field gel electrophoresis (PFGE) using *Xba*I restriction endonuclease. An epidemiological investigation was conducted to determine the source of infection and mode of transmission using a structured questionnaire. A total of 33 cases involving mainly infants and toddlers were detected in the 3-month long outbreak. The outbreak strain was of the R-type ACGSTS_u, i.e. resistant to ampicillin, chloramphenicol, gentamicin, streptomycin, tetracycline and sulphonamide. PFGE showed all isolates had an indistinguishable pattern, indicating a common source of infection. Consumption of imported dried anchovy was found to be the vehicle of transmission after adjusting for all confounding variables in the case-control study using stepwise logistic regression (OR 25.6; 95% CI 3.9–167.9; *P* = 0.001). Imported dried seafood should be properly processed, packed, labelled, and thoroughly cooked to prevent transmission of multidrug-resistant *S.* Typhimurium.

INTRODUCTION

The emergence of salmonella strains which are resistant to commonly used antimicrobials is of importance to clinicians, microbiologists and those responsible for the control of foodborne infectious diseases. *Salmonella enterica* subsp. *enterica* serotype Typhimurium definitive type 104 (DT104) with resistance to ampicillin, chloramphenicol, streptomycin, sulphonamide and tetracycline (ACSSuT) was first reported in the United Kingdom in 1984 [1, 2]. Since then, it has been isolated with increasing frequency in other European countries, Japan and the United States [3–6]. In the United Kingdom, it is the second most prevalent type of salmonella isolated from humans [7]. Outbreaks related to this strain have also been reported in Ireland, Northern California and

Denmark [8–10] and the vehicles linked with foodborne transmission include chicken, pork sausage and meat paste, contaminated beef, turkey and raw-milk cheese [11–14].

In July 2000, an unusual increase in the isolation of *S.* Typhimurium DT104L with resistance to ampicillin, chloramphenicol, gentamicin, streptomycin, tetracycline and sulphonamide or R-type ACGSTS_u, was noted. The isolates came mainly from the stool cultures of children under 18 months of age. We report here the findings of our microbiological and epidemiological investigations.

METHODS

Microbiological investigations

S. Typhimurium was identified by serotyping using slide and/or tube agglutination following the

* Author for correspondence: Department of Pathology, Singapore General Hospital, Outram Road, Singapore 169608.

Kauffman–White scheme and phage typed by the method of Ward et al. [15]. The isolates were tested for susceptibility to 14 antimicrobial agents by agar disk diffusion tests on Mueller–Hinton agar (BBL Microbiology Systems, Cockeysville, MD) according to the National Committee for Clinical Laboratory Standards [16]. The agents were ampicillin (10 µg), ceftriaxone (30 µg), chloramphenicol (30 µg), ciprofloxacin (5 µg), co-trimoxazole (1.25/23.75 µg), gentamicin (10 µg), amikacin (30 µg), nalidixic acid (30 µg), kanamycin (30 µg), streptomycin (10 µg), tetracycline (30 µg), neomycin (30 µg), sulphonamide (0.25 mg) and trimethoprim (5 µg).

Isolates were subcultured on Tryptic Soy Agar (TSA) containing 5% sheep blood overnight at 37 °C and colonies were inoculated into 10 ml of Tryptic Soy Broth (TSB) medium. After overnight incubation at 37 °C, the cells were pelleted by centrifugation and resuspended in sterile saline equal to the weight of the cell pellet. This suspension (20 µl) was added to 300 µl of TEN buffer (0.1 M Tris–HCL, 0.15 M NaCl, 0.1 M EDTA [pH 7.5]). 340 µl of 2% low-melting-point agarose (FMC BioProducts, Rockland, ME) was added to the TEN buffer mixture and pipetted into plug moulds. The agarose plugs were incubated for 5 h at 37 °C in 2 ml EC buffer (6 mM Tris, 1 M NaCl, 0.01 M EDTA, 0.5% Brij 58, 0.2% sodium deoxycholate, 0.5% Sarkosyl [pH 7.5] with 200 µl lysozyme (20 mg/ml)), washed twice with 5 ml 1 × CHEF TE buffer (10 mM Tris–HCL, 1 mM EDTA), deproteinated in 2 ml ES buffer (400 mM EDTA, 1% Sarkosyl) with 100 µl proteinase K (20 mg/ml) and incubated overnight at 50 °C. The plugs were washed four times with 5 ml 1 × CHEF TE buffer and genomic DNA was digested with restriction endonuclease *Xba*I (New England Biolabs). Electrophoresis was carried out with a GenePath apparatus (Bio-Rad Laboratories, Richmond, CA) in gels of 1% agarose in 0.5 × Tris-borate-EDTA buffer for 25 h at 200 V at a temperature of 14 °C with ramped times of 2–40 s. The gels were stained with ethidium bromide (0.2 µg/ml) and photographed under UV transillumination. The DNA size standards used were a lambda ladder PFG marker of size range 50–1000 kb (New England, Biolabs).

Epidemiological investigations

A case was defined as an infant or toddler aged between 1 month and 36 months whose stool cultures were positive for *S. Typhimurium* DT104L during the

period July–August 2000. Each case was traced and investigated to determine the source of infection and the mode of transmission. The home of each child was visited to look for other unreported cases among contacts. The parent, guardian, or domestic maid was interviewed by two trained environmental health officers using a structured questionnaire. As well as the basic epidemiological data (age, gender, ethnic group, residential address and onset of clinical symptoms), the questionnaire included a list of food items consumed and other risk factors likely to be linked to transmission, such as food handling and personal hygiene practices in the kitchen and contact with household pets with diarrhoea. For each notified case, 2–3 controls comprising children of the same age group (within 6 months of the age of the case), gender and ethnic group living in the same neighbourhood, and with no recent travel history and gastro-intestinal symptoms during the previous 2 weeks, were selected and interviewed. The questionnaire covered 3 days preceding the onset of symptoms of the cases, and for the controls, 3 days preceding the interview. All interviews were carried out within 3 weeks of onset of illness of the reported cases.

Environmental investigation

Samples of milk powder, cereal and other food items were obtained from the homes of the cases and submitted to the Food and Water Laboratory, Singapore General Hospital, for isolation of *Salmonella* species.

Statistical analyses

Differences in proportions between cases and controls were first compared using χ^2 or Fisher's exact test and odds ratio with 95% confidence interval derived. Multiple logistic regression was used to adjust simultaneously for potentially confounding factors [17]. All calculations were performed using SPSS and Excel microcomputer based software. A *P*-value of < 0.05 was considered to be statistically significant in a two-tailed test.

RESULTS

During the period 13 July–17 October 2000, 33 isolates of *S. Typhimurium* DT104L of R-type ACGSTS_u were identified. Their PFGE patterns were

Table 1. Age-gender distribution of 33 reported cases of *S. enterica subsp. enterica serotype Typhimurium DT104L*, Singapore

Age group	Male	Female	Total
< 6 months	4	2	6
6–11 months	6	4	10
1 year	3	4	7
2 years	3	0	3
3 years	1	0	1
4 years	0	0	0
5–14 years	0	2	2
15–24 years	0	0	0
25–34 years	0	1	1
35–44 years	0	1	1
55+ years	2	0	2
Total	19	14	33

indistinguishable from each other. Seventeen cases were detected in July, 10 in August, 4 in September and 2 in October. Except for 4 adults aged between 31 and 66 years, all others were children and infants as young as 2 days of age. There were 19 males and 14 females (Table 1). None of them had a recent travel history outside Singapore. Cases were not clustered in any particular locality. No secondary cases were identified in the same household.

The main presenting clinical symptoms of 19 cases investigated were watery diarrhoea (100%), fever (78.9%), vomiting (52.6%) and abdominal cramps (26.3%). About 58% had bloody stools. Seventeen of them were hospitalized and 2 were treated as outpatients. No death was reported.

Results of the case-control study based on 19 cases and 55 controls using univariate analyses, implicated porridge with 'ikan bilis' (dried anchovy) as the vehicle of transmission (OR 8.2; 95% CI 2.5–26.9; $P < 0.0001$). No other food items such as egg, dairy products, chicken, pork, beef, fruit or raw vegetables were incriminated (Table 2). Among the risk factors analysed, there was a strong inverse relationship between handwashing before preparing food and the development of illness (OR 0.17; 95% CI 0.04–0.69; $P = 0.02$). However, there was no significant association between illness and contact with animals or household members with diarrhoea. After adjusting for confounding variables using forward stepwise logistic regression, porridge with 'ikan bilis' remained significantly associated with the illness (OR 25.6; 95% CI 3.9–167.9; $P = 0.001$).

None of the food samples taken from the homes of six cases, including one sample of ground dried anchovies, was positive for *Salmonella* species. However, *Salmonella* group E was isolated from 1 of the 16 samples of dried anchovies subsequently purchased from the retail outlets.

DISCUSSION

Outbreaks of salmonellosis caused by multidrug-resistant strains have occurred previously in Singapore: *S. Typhimurium* in 1971–97, *S. Oranienburg* in 1978–9, and *S. Blockley* in 1983. However, the vehicles of transmission remained undetermined. The predominant serotypes in 1999 were *S. Enteritidis* (20.3%), *S. Stanley* (12.8%), *S. Weltevreden* (10.8%), followed by *S. Typhimurium* (7.3%) [18].

Unlike in Western Europe and North America, where consumption of unpasteurized dairy products, poultry, pork and beef products has been associated with foodborne transmission of multidrug resistant *S. Typhimurium*, no such relationship could be established in this outbreak [11–14]. The only food vehicle linked to transmission was the consumption of ground 'ikan bilis' (dried anchovies). Risk factors identified in the United States and the United Kingdom, such as contact with farm animals and pets, were also not significantly associated with this outbreak [19]. The inverse relationship between handwashing before preparing food and the development of illness was not found to be statistically significant after adjusting for confounding variables.

Salmonella is one of the most prevalent enteric pathogens encountered in seafood in Thailand [20]. It has been isolated from ready-to-eat imported seafood such as cooked shrimp, shellfish, fish paste, smoked fish, salted dried fish and caviar in the United States [21]. Outbreaks of salmonellosis caused by ingestion of contaminated cuttlefish chips as snacks have also been reported in Japan [22].

The 'ikan bilis' are imported from Thailand, Vietnam, Malaysia and Indonesia and sold at retail outlets throughout the country. These small anchovies caught in fishing villages in these countries are lightly salted and dried in the open air. This method of processing 'ikan bilis' is subject to gross environmental contamination, including excreta of birds, rats and flies. After processing, the dried seafood is packed in gunny sacks and transported by container trucks,

Table 2. Results of matched case-control analysis in an outbreak of *S. enterica subsp. enterica serotype Typhimurium DT104L*, Jul to Oct 2000

Food items/risk factors	Cases (n = 19)	Controls (n = 55)	OR	95% CI	P-value
Egg					
Uncooked	0/19	1/55	—	—	0.74
Partially cooked	4/19	12/55	0.96	0.27–3.42	0.94
Cooked	3/19	23/55	0.26	0.07–1.00	0.04
Diary products					
Powdered milk formula	18/19	48/55	2.63	0.30–22.86	0.37
Pasteurized milk	1/19	6/55	0.45	0.05–4.03	0.47
Reconstituted/UHT milk	0/19	1/55	—	—	0.74
Cereals with milk	8/19	21/55	1.18	0.41–3.40	0.76
Baby food	1/19	1/55	3.00	0.18–50.47	0.43
Cheese	9/19	13/55	2.91	0.97–8.69	0.05
Yoghurt	0/19	4/55	—	—	0.30
‘Yakult’	2/19	12/55	0.42	0.09–2.09	0.28
‘Vitagen’	2/19	9/55	0.60	0.12–3.07	0.54
Porridge					
Alone	6/19	10/55	2.08	0.64–6.80	0.22
With chicken	4/19	26/55	0.30	0.09–1.01	0.05
With pork	7/19	29/55	0.52	0.18–1.53	0.23
With beef	0/19	4/55	—	—	0.30
With liver	1/19	6/55	0.45	0.05–4.03	0.47
With ‘ikan bilis’ (dried anchovy)	14/19	14/55	8.2	2.5–26.9	0.0001
Breast fed					
With supplements	0/19	1/55	—	—	0.74
Without supplements	0/19	4/55	—	—	0.30
Fruits	11/19	21/55	2.23	0.77–6.43	0.14
Fruit juices	4/19	11/55	1.07	0.30–3.86	0.92
Raw vegetables	0/19	2/55	—	—	0.98
Animal contact	0/19	3/55	—	—	0.57
Household contact with diarrhoea	0/19	5/55	—	—	0.24
Handwashing before preparing food	13/19	51/55	0.17	0.04–0.69	0.02
Storage of ‘ikan bilis’ in chiller	4/19	10/55	1.2	0.33–4.40	0.78

fishing boats or lorries to Singapore. Further contamination could have occurred during transport, storage and display at retail outlets where it was sold without proper packing and labelling. The shelf life of the product is 1–2 weeks. It has a high turnover rate at the retail outlet and is sold out within a few days. The dried anchovy is a rich source of calcium and protein and is a popular ingredient for the preparation of a variety of foods, including soup stock. It is commonly deep fried and served with chilli. If used as an infant feed, the common practice is to grind it and then cook it with porridge. In the households where cases had occurred, the ‘ikan bilis’ was first ground and stored in a container at room temperature for subsequent use. It was either sprinkled on or added to

freshly cooked porridge, instead of boiling with the porridge, and served.

The finding of a group E *Salmonella* in the dried anchovies showed that this product can be a potential food vehicle for transmission of salmonellosis caused by other serotypes. The means of prevention of salmonellosis through imported dried seafood is to implement hazard analyses and critical control points to reduce contamination. Products should be properly labelled and packed before transportation in a closed container to retail outlets. The practice of sprinkling or adding ground ‘ikan bilis’ on cooked porridge should be discouraged as the temperature attained is inadequate to destroy the salmonella present in the food.

ACKNOWLEDGEMENTS

We thank Dr Mavis Yeo for critically reviewing the manuscript and offering useful suggestions. We are grateful to Dr Ho Ling and Dr Raymond Lin for their early notification of the outbreak cases.

REFERENCES

- Threlfall EJ, Ward LR, Frost JA, Willshaw GA. Spread of resistance from food animals to man – the UK experience. *Acta Vet Scand Suppl* 2000; **93**: 63–8.
- Hosek G, Leschincky D, Irons S, Safranek TJ. Multidrug-resistant *Salmonella* serotype *typhimurium* – United States, 1996. *MMWR* 1997; **46**: 308–10.
- Natassi A, Mammina C. Surveillance of multidrug-resistant strains of *Salmonella enterica* serotype *typhimurium* in southern Italy in the years 1992–1997. *Eur J Epidemiol* 2000; **16**: 135–9.
- Sameshima T, Akinab M, Izumiya H, et al. *Salmonella typhimurium* DT104 from livestock in Japan. *Jpn J Infect Dis* 2000; **53**: 15–6.
- Matsumoto M, Suzuki M, Hiramatsu R, et al. An increase in multi-drug resistant isolates of *Salmonella typhimurium* from healthy carriers in Aichi, Japan. *Jpn J Infect Dis* 2000; **53**: 164–5.
- Besser TE, Goldoft M, Pritchett LC, et al. Multidrug-resistant *Salmonella typhimurium* DT104 infections of humans and domestic animals in the Pacific Northwest of the United States. *Epidemiol Infect* 2000; **124**: 193–200.
- Threlfall EJ, Frost JA, Ward LR, Rowe B. Increasing spectrum of resistance in multidrug-resistant *Salmonella typhimurium*. *Lancet* 1996; **347**: 1053–4.
- Skov MN, Angen O, Chriel M, Olsen JE, Bisgaard M. Risk factors associated with *Salmonella enterica* serovar *typhimurium* infection in Danish broiler flocks. *Poult Sci* 1999; **78**: 848–54.
- Berends BR, Van Knapen F, Mossel DA, Burt SA, Snijders JM. *Salmonella* sp. on pork at cutting plants and at the retail level and the influence of particular risk factors. *Int J Food Microbiol* 1998; **44**: 207–17.
- Anonymous. Outbreak of *Salmonella* serotype *typhimurium* infection associated with eating raw ground beef – Wisconsin, 1994. *JAMA* 1996; **275**: 353–4.
- Grein T, O’Flanagan D, McCarthy T, Bauer D. An outbreak of multidrug-resistant *Salmonella typhimurium* food poisoning at a wedding reception. *Ir Med J* 1999; **92**: 238–41.
- Villar RG, Macek MD, Simons S, et al. Investigation of multidrug-resistant *Salmonella* serotype *typhimurium* DT104 infections linked to raw-milk cheese in Washington. *JAMA* 1999; **281**: 1811–6.
- Cody SH, Abbott SL, Marfin AA, et al. Two outbreaks of multidrug-resistant *Salmonella* serotype *typhimurium* DT104 infections linked to raw-milk cheese in Northern California. *JAMA* 1999; **281**: 1805–10.
- Molbak K, Baggesen DL, Aarestrup FM, et al. An outbreak of multidrug-resistant, quinolone-resistant *Salmonella enterica* serotype *typhimurium* DT104. *N Engl J Med* 1999; **341**: 1420–5.
- Ward LR, De Sa JDH, Rowe B. A phage-typing scheme for *Salmonella enteritidis*. *Epidemiol Infect* 1987; **99**: 291–4.
- National Committee for Clinical Laboratory Standards. Performance Standards for Antimicrobial Disk Susceptibility Testing, 7th edn. Approved standards M2-A7. Wayne, PA: National Committee for Clinical Laboratory Standards, 2000.
- Glantz SA, Slinker BK. Primer of applied regression and analysis of variance. New York: McGraw Hill, 1990: 512–68.
- Annual Report Department of Pathology, Singapore General Hospital, 1999.
- Wall PG, Morgan D, Lamden K, et al. A case control study of infection with an epidemic strain of multi-resistant *Salmonella* PT104 in England and Wales. *CDR* 1994; **4**: R126–31.
- Pilantanapak A, Jayanetra P, Panbangred W, Klungthong C, Bangtrakulnonth A. Construction of specific DNA probe for the detection of *Salmonella* in food. *SE Asia J Trop Med Publ Hlth* 1997; **28**: 73–81.
- Heinitz ML, Ruble RD, Wagner DE, Tatini SR. Incidence of *Salmonella* in fish and seafood. *J Food Prot* 2000; **63**: 579–92.
- Hamada K, Tsuji H, Masuda K, Uemura K. Outbreak of salmonellosis caused by ingestion of cuttlefish chips: epidemiological analysis by pulse-field gel electrophoresis. *Jpn J Infect Dis* 1999; **52**: 53–4.