

# Molecular and spatial epidemiology of human campylobacteriosis: source association and genotype-related risk factors

P. MULLNER<sup>1,2\*</sup>, T. SHADBOLT<sup>3</sup>, J. M. COLLINS-EMERSON<sup>1</sup>,  
A. C. MIDWINTER<sup>1</sup>, S. E. F. SPENCER<sup>1</sup>, J. MARSHALL<sup>1</sup>,  
P. E. CARTER<sup>4</sup>, D. M. CAMPBELL<sup>2</sup>, D. J. WILSON<sup>5</sup>, S. HATHAWAY<sup>2</sup>,  
R. PIRIE<sup>4</sup> AND N. P. FRENCH<sup>1</sup>

<sup>1</sup> Molecular Epidemiology and Veterinary Public Health Laboratory, Hopkirk Research Institute, Massey University, Palmerston North, New Zealand

<sup>2</sup> New Zealand Food Safety Authority, Science Group, Wellington, New Zealand

<sup>3</sup> MidCentral Public Health, Rata House, Palmerston North Hospital, Palmerston North, New Zealand

<sup>4</sup> Institute for Environmental Science and Research, Kenepuru Science Centre, Porirua, New Zealand

<sup>5</sup> Department of Human Genetics, University of Chicago, Chicago, IL, USA

(Accepted 3 December 2009; first published online 9 February 2010)

## SUMMARY

The epidemiology of human campylobacteriosis is complex but in recent years understanding of this disease has advanced considerably. Despite being a major public health concern in many countries, the presence of multiple hosts, genotypes and transmission pathways has made it difficult to identify and quantify the determinants of human infection and disease. This has delayed the development of successful intervention programmes for this disease in many countries including New Zealand, a country with a comparatively high, yet until recently poorly understood, rate of notified disease. This study investigated the epidemiology of *Campylobacter jejuni* at the genotype-level over a 3-year period between 2005 and 2008 using multilocus sequence typing. By combining epidemiological surveillance and population genetics, a dominant, internationally rare strain of *C. jejuni* (ST474) was identified, and most human cases (65·7%) were found to be caused by only seven different genotypes. Source association of genotypes was used to identify risk factors at the genotype-level through multivariable logistic regression and a spatial model. Poultry-associated cases were more likely to be found in urban areas compared to rural areas. In particular young children in rural areas had a higher risk of infection with ruminant strains than their urban counterparts. These findings provide important information for the implementation of pathway-specific control strategies.

**Key words:** Bacterial typing, *Campylobacter*, molecular epidemiology, spatial modelling, surveillance.

## INTRODUCTION

Cases of campylobacteriosis in New Zealand have been increasing steadily in the last decade, peaking at

379 cases/100 000 people per annum in 2006 [1]. This situation has been accompanied by a considerable amount of media attention with public calls for urgent action to reduce the unacceptable burden of human

\* Author for correspondence: Dr P. Mullner, Molecular Epidemiology and Veterinary Public Health Laboratory, Hopkirk Research Institute, Massey University, Private Bag 11 222, Palmerston North, NZ. (Email: petra@epi-interactive.com)

disease [2]. Internationally rare clones of *Campylobacter* could be isolated during an epidemic [3], but their role in the long-term occurrence of disease remained unclear. The complex epidemiology of campylobacteriosis has hampered the development of successful control programmes for this pathogen in New Zealand and elsewhere. *Campylobacter* spp. have been isolated from a wide range of food sources including poultry [4, 5], red meat [6] and milk [7], as well as environmental sources such as wild birds [8] and contaminated water [9]. In addition a variety of risk factors for acquiring campylobacteriosis, such as consumption of offal and dog-ownership have previously been identified [10, 11].

The ecological situation in New Zealand is unique: there is extensive agricultural usage of land and the islands are geographically remote, resulting in human, animal and pathogen populations that are relatively isolated. The country has among the highest enteric infectious disease rates in industrialized countries [12], and the high ratio of food-producing animals to humans, and frequent use of rural water supplies in New Zealand have been proposed as underlying causes [13]. The poultry supply is also different from many other developed countries, in that suppliers are almost entirely focused on the domestic market and for biosecurity reasons no raw poultry products are imported into the country. Despite little movement of animals and animal products into the country as a result of tight biosecurity measures, New Zealand is, however, exposed to a relatively large number of international travellers [14].

Previous studies have indicated that in New Zealand [15], as in other countries [16], the species *Campylobacter jejuni* is responsible for the majority (82.6%) of human campylobacteriosis cases. *C. jejuni* differs in its epidemiology from other *Campylobacter* spp., as illustrated in a case-case comparison for *C. jejuni* and *C. coli* [17]. There is evidence that these differences also apply to different strains [18], which may affect inferences made at the species level such as the estimation of risk factors and population attributable risk. In addition, commonly used typing schemes for *Campylobacter* spp., such as serotyping, have struggled to provide sufficient resolution for detailed epidemiological studies based on the differences between strains. The increasing availability of genotyping methods allows for the study of transmission patterns and risk factors at the strain level, and this is changing the way epidemiologists study infectious diseases [19]. Multilocus sequence typing (MLST) has

emerged as a genotyping technique for *Campylobacter* [20]. It has revealed differences in the epidemiology of individual genotypes [21] and allows for the identification of source-associated strains [22, 23]. Based on the sequence of seven, relatively conserved, house-keeping genes, this approach has several properties, such as efficiency, accuracy, reproducibility and the opportunity to share data via an openly accessible database. These properties make it suitable as a typing tool for epidemiological investigations of genetically diverse pathogens [24, 25]. In addition MLST has informed recently developed epidemiological and population-genetic source attribution models such as the Hald and island models [26]. These models aim to apportion human cases to disease sources, and hence estimate the relative importance of each source. They are therefore a valuable tool for decision-making in food safety. Furthermore, the results from the genetic analysis performed by the asymmetric island model [27] can be used to assign pathogen strains to likely sources for further molecular epidemiological analysis.

In response to the high burden of human disease in New Zealand a long-term sentinel surveillance site was established in the Manawatu region to better understand key determinants of human campylobacteriosis by combining longitudinal epidemiological surveillance and genotyping data. In this paper we investigate the epidemiology of *C. jejuni* infection at the genotype level using data from this site. Patterns of human cases are described and spatial and multi-variable analyses are used to identify risk factors for source-associated strains. We compare the molecular epidemiology of *C. jejuni* in New Zealand with findings from overseas, and discuss the implications of our results for the prevention and control of this important zoonosis.

## MATERIAL AND METHODS

### Study population

The study was conducted within the Manawatu Health District of New Zealand's North Island. The population at the 2006 census consisted of 155 072 people [28] living in a mixture of rural and urban dwellings, including both coastal and inland areas. The main urban centre is the city of Palmerston North, which has a population of about 75 000 people.

### Case definition

A case was defined as a laboratory-confirmed sporadic *C. jejuni* case in the Manawatu Health District between 1 March 2005 and 29 February 2008. Outbreak-related cases, except the index case, were excluded. Campylobacteriosis is a notifiable disease under the Health Act 1956 in New Zealand. Data from New Zealand's national notifiable disease database EpiSurv [29] were obtained for cross-validation with notified case numbers in the study region. EpiSurv is operated by the Institute of Environmental Science and Research Ltd (ESR) on behalf of the Ministry of Health. General practitioners and public health units are required to report notified cases of campylobacteriosis to ESR for nationwide disease surveillance.

### Case ascertainment

Cases were prospectively recruited via the regional diagnostic laboratory, which tests all human pathological samples from the region. Samples were tested with a commercial enzyme immunoassay that detects a *Campylobacter*-specific antigen, shared by *C. jejuni* and *C. coli* (ProSpecT; Remel, USA). Positive faecal swabs (Amies Charcoal transport swabs; Copan, Italy) were sent to the Hopkirk Research Institute for isolation of presumptive *Campylobacter*. In brief, swabs were cultured on modified cefoperazone charcoal desoxycholate agar plates (Fort Richard, NZ) and in Bolton Broth (Lab M, UK) and incubated at 42 °C in a microaerobic atmosphere for 2 days. A single colony resembling *Campylobacter* spp. was subcultured to Columbia horse blood agar (Fort Richard) and incubated microaerobically at 42 °C for 2 days before DNA preparations were made. Isolates were confirmed as *C. jejuni* by polymerase chain reaction. If multiple samples were collected from a single patient, only one isolate was selected.

### Genotyping

After speciation, MLST of *C. jejuni* isolates was performed to assign isolates to a sequence type (ST) using seven housekeeping genes: *aspA* (aspartase A), *glnA* (glutamine synthase), *gltA* (citrate synthase), *glyA* (serine hydroxymethyltransferase), *pgm* (phosphoglucomutase), *tkt* (transketolase) and *uncA* (ATP synthase  $\alpha$ -subunit) based on the method outlined by Dingle *et al.* [20].

### Epidemiological surveillance

Three years of human data were acquired; these were initially collected using routine public health service methods which were then superseded by a sentinel surveillance approach with data collection targets. Case information between February 2005 and June 2006 was acquired using both mailed questionnaires and telephone interviews. To enhance the quality of the surveillance data gathered, from 1 July 2006 notified cases of campylobacteriosis were contacted by telephone with a target of 95% of cases interviewed and 95% of datasets completed. The interviews were conducted using the standard structured EpiSurv case report form used for enteric diseases in New Zealand [30]. The questionnaire gathered demographic variables, risk factor information for variables within the incubation period, as well as case characteristics. Investigators were trained to ensure consistency and standardization of interviews.

### Data handling and statistical analysis

The epidemiological dataset was merged with the genotyping information in an Access database, which was linked into a Bionumerics database for analysis. Data were quality controlled and validated by techniques such as consistency check, range check and batch totals. R, version 2.7.0, was used for statistical analysis [31]. Distributions were compared using  $\chi^2$  tests and Fisher's exact test for count data.

All STs isolated from human cases were assigned a probability that they were acquired from individual animal reservoirs using the asymmetric island model [27]. Data for this assignment were collected from parallel structured studies at the sentinel surveillance site over the same time period [22, 26, 32]. To attribute STs to their most likely source, we first computed the attribution probabilities of each ST to each source using the asymmetric island model. The distribution of the attribution probabilities of the most likely source of each ST (i.e. the maximum attribution probability among the sources for a given ST) was assessed. Based on this distribution, a cut-off point was determined to provide a reasonable balance between the number of STs assigned to one of the three sources, and the confidence as to their correct assignment. STs with an estimated probability of at least 60% of having originated from a certain source were assigned to that source, thereby creating four different groups of human cases: poultry-associated ST474,

other poultry-associated STs, ruminant-associated STs and STs not associated with any of these sources.


The spatial location of each notified human case was provided at the meshblock level. These are the smallest regions defined for the New Zealand census, each containing between zero and 498 individuals, with a median size of 78. The spatial resolution was therefore much higher in urban areas. Urban and rural areas were defined following the Urban/Rural Profile Classification provided by Statistics New Zealand [33]. The risk of being a case of campylobacteriosis in each meshblock was described using a Bayesian hierarchical model with a Markov random field prior [34]. Relative risk surfaces were prepared for human cases from the first three of the four groupings of STs produced by the asymmetric island model. STs not associated with a particular source were not analysed as there were insufficient data.

More precisely, the model is defined as follows.  $Y_{i,t}$  represents the number of notified cases from a particular group of STs of campylobacteriosis in meshblock  $i$  and week  $t$ . We assumed that  $Y_{i,t} \sim \text{Poisson}(n_i \lambda_i)$ , where  $n_i$  is the usually resident population of meshblock  $i$  (obtained from the most recent census) and  $\lambda_i$  represents the expected risk of notification for this meshblock. We assumed that the logs of the  $\lambda_i$  follow a Gaussian Markov Random Field prior (also called a Gaussian intrinsic autoregression) in which the risk in each meshblock is assumed to be similar to the mean risk of the neighbouring meshblocks. More formally, we assumed the following full conditional distributions:

$$\log(\lambda_i) \sim N\left(\sum_{j \in n(i)} \frac{\log(\lambda_j)}{|n(i)|}, \frac{1}{\kappa |n(i)|}\right), \quad (1)$$

where  $n(i)$  is the set of meshblocks that are neighbours to meshblock  $i$ . For the hyperparameter  $\kappa$  we assumed the mildly informative and conjugate gamma-distributed prior:  $\kappa \sim \text{Gamma}(1, 0.5)$ . Several prior distributions were examined and this one produced the best balance between aiding convergence of the Markov chain and allowing sufficient flexibility in the model to ensure that the data could be well represented in the posterior.

In order to investigate the associations between covariate information and infection with poultry- and ruminant-associated strains, a multivariable logistic regression model was constructed. The binary response variable was whether or not the infection was with a poultry-associated strain vs. a ruminant strain.



**Fig. 1.** The number of ELISA positive (—), culture positive (---) and genotyped (.....) human samples from the Manawatu surveillance site, New Zealand, in each month between March 2005 and February 2008.

Multiple covariates, both numerical (such as age) and categorical (such as urban/rural living classification) were analysed using the defined source associations of strains and odds ratios (OR) were calculated where appropriate. Variables for inclusion in the final models were chosen by exploratory univariable analysis with a  $P$  value cut-off of 0.2. Final variables were chosen from all variables that met this criterion via a backwards elimination. The final model was of the form:

$$\text{logit}(y_i) = \beta_0 + \beta_1 \text{age} + \beta_2 \text{rural}, \quad (2)$$

where age was treated as a continuous variable, and rural living was binary.

## RESULTS

### Description of cases

Examination of laboratory and surveillance data revealed that 774 (75%) of the 1036 swabs submitted between 1 March 2005 and 29 February 2008 could be considered 'primary samples' that originated in the Manawatu area. The remainder were duplicate samples, or samples from outside the study area. Of these 662 (86%) were culture positive and 581 (88%) were confirmed as *C. jejuni*, of which 502 (86%) provided complete MLST data. The remainder provided partial profiles or could not be resuscitated (Fig. 1). The comparison of demographic data in *C. jejuni* cases for which an isolate was genotyped and *C. jejuni* cases for

Table 1. *Distribution of selected demographic characteristics in surveillance-linked C. jejuni cases for which a bacterial isolate was genotyped (typed) and cases for which no isolate was genotyped (non-typed), Manawatu Health District, New Zealand 2005–2008*

	Typed ( <i>n</i> = 459)†		Non-typed ( <i>n</i> = 64)‡		<i>P</i> value*
	No. of cases	%	No. of cases	%	
Sex					
Male	248	54.3	40	63.5	0.21
Female	209	45.7	23	36.5	
Age group (years)					
< 20	141	30.7	19	29.7	0.23
20–50	184	40.0	32	50.0	
> 50	134	29.2	13	20.0	

\* *P* value from a  $\chi^2$  test.

† Sex was unknown for two cases.

‡ Sex was unknown for one case.

which no isolate was genotyped revealed no significant difference between typed and non-typed cases (Table 1). The response rate to the questionnaire, applied to all notified cases in the Manawatu Health District, increased after the introduction of data collection targets in July 2006 from 58% to 97% with a 96–100% completion of case-report data fields. A significant difference could be observed for the distribution of gender ( $\chi^2=4.12$ , D.F. = 1, *P* = 0.04) before (*n* = 207) and after (*n* = 252) the changes to the interview process, with an increase in male respondents as a result of the enhanced surveillance. The age distribution of respondents was not significantly different between the two periods ( $\chi^2=4.15$ , D.F. = 2, *P* = 0.13).

### Incidence

A total of 971 cases from the Manawatu region were notified to the national notifiable disease database EpiSurv during the study period compared to 774 cases included in the laboratory recruited study. The number of cases reflected the number of notifications in each week with only minor aberrations (data not shown). Case numbers increased during each southern hemisphere summer, with an additional peak of cases in winter in 2006. Annual notification rates (per 100 000 population) in the region for the 3-year period were 206 for the first year, 232 for the second and 188 in the final year of the study.

### Distribution and source association of genotypes

A total of 51 different MLST genotypes were detected in our study. Seven STs each accounted for more than

20 cases and amounted to a total of 66% of all cases (Table 2). The most dominant genotype was ST474 accounting for the largest proportion of human cases in most quarters, contributing a total of 154 cases (30.7%) over the study period. Other major genotypes were ST48 and ST45, which could be isolated from 8.4% and 8.2%, respectively, of human cases. Common STs, occurring in over 20 cases, were STs 53, 50, 190 and 354. A total of 23 STs were poultry-associated including STs 474, 45, 48, 190 and 354. Sixteen STs were associated with ruminant sources including STs 61, 38, 42 and 2026 (Table 2).


### Risk factor analysis

Of the 502 cases typed using MLST 459 (90.4%) could be linked to a notified case of campylobacteriosis and thus provided the dataset for the risk factor analysis. A total of ten cases associated with overseas travel were excluded, as well as one case with missing information on key risk factors, bringing the total number of cases in the dataset to 448. This included 141 ST474 cases, 209 cases with other poultry-associated STs, 70 cases with ruminant-associated STs and 28 cases which could not be assigned to a probable source. The genotype distribution between linked and unlinked cases was compared using a  $\chi^2$  test and no significant differences between the two groups could be observed ( $\chi^2=58.98$ , D.F. = 51, *P* = 0.21). The age distributions for ruminant- and poultry-associated strains are shown in Figure 2. Ruminant-associated strains were relatively more common in cases in children aged < 10 years compared to adults.

Table 2. Frequency and source association of human *C. jejuni* MLST genotypes, Manawatu Health District, New Zealand, 2005–2008

Genotype	Absolute frequency	Relative frequency (%)	Source association*	Source probability (%)
ST21	7	1.4	Ruminant	61
ST25	1	0.2	Poultry	97
ST38	13	2.6	Ruminant	88
ST42	19	3.8	Ruminant	67
ST45	41	8.2	Poultry	95
ST48	42	8.4	Poultry	98
ST50	23	4.6	Poultry	73
ST52	17	3.4	Poultry	98
ST53	27	5.4	Poultry	63
ST61	14	2.8	Ruminant	87
ST81	1	0.2	Ruminant	92
ST137	1	0.2	Poultry	64
ST190	21	4.2	None	—
ST219	1	0.2	Ruminant	62
ST257	12	2.4	Poultry	98
ST354	23	4.6	Poultry	98
ST403	1	0.2	Ruminant	99
ST422	3	0.6	Ruminant	85
ST436	4	0.8	Ruminant	80
ST451	8	1.6	Poultry	98
ST459	1	0.2	Ruminant	86
ST474	154	30.7	Poultry	89
ST520	8	1.6	Poultry	66
ST578	1	0.2	Ruminant	94
ST583	9	1.8	Poultry	95
ST658	2	0.4	None	—
ST677	5	1.0	Poultry	70
ST829	1	0.2	Poultry	82
ST1457	1	0.2	None	—
ST1517	4	0.8	Poultry	68
ST1581	1	0.2	Poultry	98
ST1707	1	0.2	None	—
ST2026	11	2.2	Ruminant	88
ST2219	1	0.2	None	—
ST2343	1	0.2	None	—
ST2345	3	0.6	Poultry	98
ST2350	2	0.4	Ruminant	82
ST2391	1	0.2	Poultry	98
ST3072	1	0.2	Ruminant	88
ST3222	1	0.2	Ruminant	85
ST3538	1	0.2	None	—
ST3676	2	0.4	Ruminant	71
ST3711	2	0.4	None	—
ST3712	2	0.4	None	—
ST3715	1	0.2	None	—
ST3717	1	0.2	Poultry	98
ST3718	1	0.2	Poultry	88
ST3720	1	0.2	Ruminant	60
ST3727	1	0.2	None	—
ST3784	1	0.2	Poultry	91
ST3792	1	0.2	Poultry	92
Total	502			

\* Source association was determined by applying the island model.



**Fig. 2.** (a) Age distribution of cases associated with poultry (■,  $n=351$ ) and ruminant sources (□,  $n=70$ ) in the Manawatu between March 2005 and February 2008. (b) Age distribution of urban cases associated with poultry (■,  $n=295$ ) and ruminant sources (□,  $n=46$ ). (c) Age distribution of rural cases associated with poultry (■,  $n=55$ ) and ruminant sources (□,  $n=24$ ). The error bars represent 95% confidence intervals. The comparison is within group (i.e. poultry- or ruminant-associated cases) across age categories; e.g. 29% of ruminant-associated cases were aged <10 years compared to 7% in adults aged  $\geq 70$  years.

In contrast the age distribution of poultry-associated strains was more even across age categories, with no obvious association with children. This difference in age profiles for ruminant- compared to poultry-associated cases was significant ( $\chi^2=6.48$ , D.F. = 1,  $P=0.011$ ). Figure 2 also shows the age distributions for both poultry- and ruminant-associated cases divided into rural and urban residency. In rural areas ruminant-associated cases were relatively more common in younger children, while in urban areas ruminant cases were much more evenly distributed across all age groups.

Table 3 shows results from the multivariable case-case analysis comparing the risk factors for infection with poultry strains compared to ruminant strains. Age and location were identified as predictors. Older people were more likely to be infected with poultry strains compared to younger people; e.g. the estimated OR of being a poultry-associated rather than a ruminant-associated case for a 65-year-old adult compared to a 5-year-old child was  $OR=2.1$  ( $e^{0.012*60}$ ). Conversely children were therefore more likely to be infected with ruminant strains compared to adults. People in rural communities were less likely to be infected with poultry strains ( $OR=0.39$  for rural compared to urban dwellers) and therefore more likely to be infected with ruminant strains, while the opposite was evident for people living in urban communities.

### Spatial analysis

Figure 3 shows the relative-risk surfaces for the different ST groups. Poultry-associated cases were more prevalent in urban areas than rural areas, whereas for ruminant-associated cases the reverse was evident. There was a higher risk of notification with ST474 in and around Palmerston North, and very little increased risk in the smaller urban centre of Levin. For the other poultry-associated STs, there was a smaller increase in risk near Palmerston North, but a more significant increase in risk around the urban areas of Levin and Foxton. The relative risk for ST474 shows the most spatial variation, followed by other poultry and then ruminant STs, which exhibited the least variation of the three groups.

### DISCUSSION

We used data collected from a 3-year longitudinal study to explore the spatial and molecular epidemiology of *C. jejuni* at the genotype level, and findings from this study have influenced the development of disease control strategies for this pathogen in New Zealand [26]. Through the integration of molecular and population genetic tools into epidemiological approaches new insight has been gained into the determinants of New Zealand's most important zoonosis. Our sentinel sampling strategy, gathering longitudinal data concurrent in both space and time, differs from other studies with similar objectives that have used data from different geographical regions and time periods [27, 35]. A concentration of

Table 3. Results of a multivariate case-case risk-factor analysis for infection with poultry strains vs. ruminant strains, Manawatu region of New Zealand 2005–2008

Exposure	Estimate	95% CI	OR (95% CI)	P value
Intercept	1.462			
Age in years (continuous)	0.012	0 to 0.023	—	0.052
Living in a rural area	−0.949	−1.528 to −0.369	0.39 (0.22 to 0.69)	0.001

OR, Odds ratio; CI, confidence interval.

resources in the defined site provided a rich source of information and included advantages such as (i) standardized microbiological and epidemiological methods, (ii) high-quality data on individual cases as well as (iii) timeliness and cost-effectiveness.

An estimate of the population at risk was available in this study, allowing incidence rates of notified cases to be calculated. Manawatu Health District was an area of moderate disease rate in the study period (188–232 cases/100 000) compared to the rest of New Zealand, where in some regions and time periods more than 500 cases/100 000 population were notified [1]. From 2005 to 2007 for the whole of New Zealand notification rates ranged from 302 to 379 cases/100 000 population. Rates in our study population were above those of comparable countries such as USA (12.71/100 000), EU-25 (46.1/100 000) or Australia (111.8/100 000) [36–38]. Previous studies of the national notification system have confirmed that rates of campylobacteriosis in New Zealand are comparatively high by international comparison [39]. The observed 2006 nationwide increase in cases during the southern hemisphere winter [3] is also evident in our study as well as the expected seasonal increase in campylobacteriosis during the warmer months [40].


A good correlation was observed between the number of cases statutorily notified in the Manawatu region and the number of faecal samples from primary cases submitted to the laboratory. The figures were not identical as the notified cases contained additional non-laboratory-confirmed cases associated with outbreaks; these were excluded in our case definition. This was highlighted in the data from November 2006 where the number of notifications deviated from the number of cases included in the study as a result of a major outbreak which included 26 outbreak-associated cases [3]. In addition differences between notifications and study cases in each month may have been caused by a delay between laboratory confirmation and notifications of cases, as well as technical problems such as lost or unidentifiable samples, or

sample submission to laboratories outside the study region. Not all cases included in the study were genotyped and the potential misclassification and differences between characteristics of patients with and without genotyped isolates present potential sources of bias. We consider, supported by a temporal analysis of the dataset, that the typed samples are representative of human cases in the region over the study period. In addition our analysis revealed no significant differences between the two groups, and we therefore consider the typed cases to be broadly representative of all notified cases. No evidence was found that cases without a linked surveillance report were different from cases with a report, and we therefore conclude that cases included in the analysis are representative of the study population.

Although we found over 50 different STs in human samples in our study, most human cases (66%) were associated with only seven different STs. These major human genotypes were generally associated with a consistent number of human cases over the time period and appear to be endemic to New Zealand. ST474 was the dominant genotype found in our study and was also the most commonly isolated genotype in the winter epidemic in 2006 in New Zealand [3]. Surprisingly this ST has only rarely been reported overseas [3] and besides human cases has been detected predominantly in samples from poultry in New Zealand [32]. The link between poultry and human cases of campylobacteriosis has long been identified [41] and poultry sources have recently been estimated to cause the majority of campylobacteriosis cases in New Zealand [22], Scotland [23] and England [27]. Although this study found a dominant clone and several other STs, such as ST2345, that are currently only found in samples collected in New Zealand, many genotypes isolated in this study can be commonly found in other countries [42, 43]. This includes major human pathogens such as STs 45, 48 and 53.

The spatial analysis (Fig. 3) showed that cases in urban areas were more likely to be infected with





**Fig. 3.** (a) Relative risk surfaces of human campylobacteriosis cases of ST474, (b) other poultry-associated strains and (c) ruminant-associated strains in the Manawatu between March 2005 and February 2008. The box shows an enlargement of the city of Palmerston North.

poultry-associated STs than rural areas. Conversely, cases in rural areas were more likely to be infected with ruminant-associated STs than urban areas. This supports the findings from a similar study in Scotland [35] where urban and rural areas were also shown to have different epidemiological patterns. Figure 3*b* shows an increased risk of poultry-associated strains around the small towns of Levin and Foxton; however, the same increase in risk was not present for ST474 cases (Fig. 3*a*). This may be because ST474 is associated with one particular poultry company [26] which has a relatively high market share in Palmerston North. A second company (not associated with ST474) has a processing plant near Foxton, and this may explain why there is an increased risk of non-ST474 poultry-associated cases near Levin and Foxton, either through occupational or environmental exposure or increased local market share of poultry products. The increased risk for infection with ruminant strains in rural areas (Fig. 3*c*) may be associated with increased exposure to faecal matter or poorer quality drinking water in these areas [9, 44, 45]. Similarly, the observed differences in age distributions for ruminant and poultry genotypes may represent different transmission pathways, such as an increased exposure to ruminant faecal matter in young children, as well as differences in immunity between age groups [46, 47]. Different exposure pathways are therefore likely to exist in rural and urban communities, and this should influence the development of control strategies.

This research suggests that the high rates of campylobacteriosis in New Zealand were associated with a dominant, internationally rare clone, namely ST474, as well the presence of internationally common human genotypes such as ST45. ST474 was also the most commonly isolated genotype in the winter epidemic in 2006 in New Zealand [3]. Currently, this ST has only one submission to the *Campylobacter* PubMLST database [48] from a chicken sample in the Czech Republic, and has only been reported sporadically outside New Zealand [27, 49, 50]. If ST474 were to display phenotypic traits associated with, for example, higher virulence, this might help to explain the relatively high incidence of campylobacteriosis in New Zealand. However, there is currently only limited evidence for this: studies by Taboada and Pope indicated that some *Campylobacter* strains, including ST474 of New Zealand origin, may have higher virulence as measured by an invasion assay [51, 52]. Further research into pathogenicity and

virulence genes of this dominant clone could reveal determinants for human infection and commensal colonization and may explain the dominance of this ST in New Zealand. The identification of ST-specific risk factors will facilitate the identification of tailor-made control programmes and by prioritizing dominant STs, more efficient control programmes can be designed.

Instead of using a network of surveillance sites this study used only one region and the importance of transmission routes is likely to vary regionally [53]. The generalization of our results from the sentinel surveillance site to the general population is supported by validation studies performed in two other regions where a similar distribution of MLST types was observed [32]. New Zealand provides a unique ecology and this is likely to have influenced the national distribution of MLST genotypes and the epidemiology of human cases [26]. However, some of the results from this study are in accord with those made in a similar study in Scotland [35]. These include the differences in risks of infection in rural and urban communities.

Source attribution studies from the Manawatu sentinel surveillance site [22, 26] provided strong evidence that poultry was the leading cause of human campylobacteriosis in New Zealand, causing an estimated 54–80% of cases. In contrast, environmental sources played a relatively minor role. This is supported by the dramatic decline in human notified cases in 2008 to a 16-year low, which is a probably a consequence of poultry industry-specific interventions that were implemented based on these findings. The analysis presented here complements those results by improving our knowledge of risk factors and transmission pathways to implement more refined control strategies; for example to protect young children from disease through exposure to farm animal faecal matter in rural settings in addition to reducing transmission through poultry products in urban and rural areas.

In conclusion, the combination of molecular and epidemiological tools in a sentinel surveillance site has improved our understanding of the epidemiology of campylobacteriosis in New Zealand. The analysis of risk factors associated with MLST groupings has revealed disease characteristics that will influence the design and implementation of control measures. The results from this study underline the importance of studying *Campylobacter* spp. beyond the strain level. Extending this work in space and time by, for

example, including data from other sites where risk factors may differ and integrating temporal components into the analysis could further improve our understanding of the commonly observed regionality and seasonality of *Campylobacter* infections [53].

## ACKNOWLEDGEMENTS

The authors are grateful to all members of the Hopkirk Molecular Epidemiology Team, Environmental Science and Research (ESR), MidCentral Health, Public Health Services, NZFSA (Science Group), and MedLab Central (in particular Lynn Rogers) for their contributions. This work was supported by the New Zealand Food Safety Authority through their Postgraduate PhD Programme: ‘Estimating the contribution of different food sources to the burden of human campylobacteriosis in the Manawatu’. In addition financial support was provided through the New Zealand Food Safety Authority project ‘Enhancing Surveillance of Potentially Foodborne Enteric Diseases in New Zealand: Human Campylobacteriosis in the Manawatu’ (FDI/236/2005).

## DECLARATION OF INTEREST

None.

## REFERENCES

1. **Institute of Environmental Science and Research Limited (ESR)**. Public Health Surveillance – EpiSurv. (<http://www.survesrcrinz/epiSurv/index/php>). Accessed 16 January 2009.
2. **Baker M, et al.** Regulation of chicken contamination urgently needed to control New Zealand’s serious campylobacteriosis epidemic. *New Zealand Medical Journal* 2006; **119**, no. 1243.
3. **McTavish SM, et al.** Wide geographical distribution of internationally rare *Campylobacter* clones within New Zealand. *Epidemiology and Infection* 2008; **136**: 1244–1252.
4. **Chrystal ND, et al.** Counts of *Campylobacter* spp. and prevalence of *Salmonella* associated with New Zealand broiler carcasses. *Journal of Food Protection* 2008; **71**: 2526–2532.
5. **Snow LC, et al.** Survey of the prevalence of *Salmonella* on commercial broiler farms in the United Kingdom, 2005/06. *Veterinary Record* 2008; **163**: 649–654.
6. **Wong TL, et al.** Prevalence, numbers, and subtypes of *Campylobacter jejuni* and *Campylobacter coli* in uncooked retail meat samples. *Journal of Food Protection* 2007; **70**: 566–573.

7. Gillespie IA. Milkborne general outbreaks of infectious intestinal disease, England and Wales, 1992–2000. *Epidemiology and Infection* 2003; **130**: 461–468.
8. French N, *et al.* Molecular epidemiology of *Campylobacter jejuni* isolated from wild bird faecal material in children's playgrounds. *Applied and Environmental Microbiology* 2008; **75**: 779–783.
9. Savill MG, *et al.* Enumeration of *Campylobacter* in New Zealand recreational and drinking waters. *Journal of Applied Microbiology* 2001; **91**: 38–46.
10. Gillespie IA, *et al.* Demographic determinants for *Campylobacter* infection in England and Wales: implications for future epidemiological studies. *Epidemiology and Infection* 2008; **136**: 1717–1725.
11. Stafford RJ, *et al.* Population-attributable risk estimates for risk factors associated with *Campylobacter* infection, Australia. *Emerging Infectious Diseases* 2008; **14**: 895–901.
12. Lake RJ, *et al.* Estimated number of cases of foodborne infectious disease in New Zealand. *New Zealand Medical Journal* 2000; **113**: 278–281.
13. Crump JA, Murdoch DR, Baker MG. Emerging infectious diseases in an island ecosystem: The New Zealand perspective. *Emerging Infectious Diseases* 2001; **7**: 767–772.
14. Statistics New Zealand. International visitors (<http://searchstats.govt.nz/search?w=international%20visitors>). Accessed 16 September 2009.
15. Devane ML, *et al.* The occurrence of *Campylobacter* subtypes in environmental reservoirs and potential transmission routes. *Journal of Applied Microbiology* 2005; **98**: 980–990.
16. Oporto B, *et al.* Prevalence and strain diversity of thermophilic *campylobacters* in cattle sheep and swine farms. *Journal of Applied Microbiology* 2007; **103**: 977–984.
17. Gillespie IA, *et al.* A case-case comparison of *Campylobacter coli* and *Campylobacter jejuni* infection: a tool for generating hypotheses. *Epidemiology and Infection* 2002; **128**: 111–118.
18. McCarthy ND, *et al.* A three-year population based investigation of human *Campylobacter jejuni* epidemiology using sequence typing and patient survey data. *Zoonoses and Public Health* 2007; **54**: 37.
19. Murray M. Determinants of cluster distribution in the molecular epidemiology of tuberculosis. *Proceedings of the National Academy of Sciences* 2002; **99**: 1538–1543.
20. Dingle KE, *et al.* Multilocus sequence typing system for *Campylobacter jejuni*. *Journal of Clinical Microbiology* 2001; **39**: 14–23.
21. Sopwith W, *et al.* Identification of potential environmentally adapted *Campylobacter jejuni* Strain, United Kingdom. *Emerging Infectious Diseases* 2008; **14**: 1769–1773.
22. Mullner P, *et al.* Source attribution of food borne zoonoses in New Zealand: a modified Hald model. *Risk Analysis* 2009; **29**: 970–984.
23. Sheppard SK, *et al.* *Campylobacter* genotyping to determine the source of human infection. *Clinical Infectious Diseases* 2009; **48**: 1072–1078.
24. Sullivan CB, Diggle MA, Clarke SC. Multilocus sequence typing – data analysis in clinical microbiology and public health. *Molecular Biotechnology* 2005; **29**: 245–254.
25. Urwin R, Maiden MCJ. Multilocus sequence typing: a tool for global epidemiology. *Trends in Microbiology* 2003; **11**: 479–487.
26. Mullner P, *et al.* Assigning the source of human campylobacteriosis in New Zealand: a comparative genetic and epidemiological approach. *Infection, Genetics and Evolution* 2009 (in press).
27. Wilson DJ, *et al.* Tracing the source of campylobacteriosis. *PLoS Genetics* 2008; **4**: e1000203.
28. Statistics New Zealand. Population Census 2006 (<http://www.stats.govt.nz/Census/2006CensusHomePage.aspx>). Accessed 16 September 2009.
29. Institute of Environmental Science and Research Limited (ESR). Public Health Surveillance – EpiSurv (<http://www.surveys.crinz.episurv/index/php>). Accessed on 16 January 2009.
30. Lake R, Whyte R, Kliem C. Evaluation of foodborne disease outbreaks/human health surveillance interface. New Zealand Food Safety Authority (NZFSA), 2005.
31. R Development Core Team. R: A Language and Environment for Statistical Computing, version 2.7.0. Vienna: R Foundation for Statistical Computing, 2005.
32. French N, Molecular Epidemiology and Veterinary Public Health Group Hopkirk Institute. Enhancing surveillance of potentially foodborne enteric diseases in New Zealand: human campylobacteriosis in the Manawatu (2008) ([http://www.nzfs.govt.nz/science/research-projects/Campy\\_Attribution\\_Manawatupdf](http://www.nzfs.govt.nz/science/research-projects/Campy_Attribution_Manawatupdf)). Accessed 16 March 2009.
33. Statistics New Zealand. Defining urban and rural New Zealand (<http://www.stats.govt.nz/sitecore/content/statistics/Home/Publications/BusinessPerformanceEnergyAndAgriculture/urban-rural-profile/defining-urban-rural-nz.aspx#profile>). Accessed 10 November 2009.
34. Besag JE, York JC, Mollié A. Bayesian image restoration with two applications in spatial statistics (with discussion). *Annals of the Institute of Statistical Mathematics* 1991; **43**: 1–59.
35. Strachan NJC, *et al.* Attribution of *Campylobacter* infections in Northeast Scotland to specific sources by use of multilocus sequence typing. *Journal of Infectious Diseases* 2009; **199**: 1–4.
36. Australian Government. Department of Health and Ageing. Australia's notifiable diseases status, 2006: Annual report of the National Notifiable Diseases Surveillance System. *Communicable Diseases Intelligence* 2008; **32**.
37. Centre for Disease Control and Prevention. FoodNet Report 2006 (<http://www.cdc.gov/foodnet/reports.htm>). Accessed 25 January 2009.
38. European Food Safety Authority. Community summary report on trends and sources of zoonoses, zoonotic agents, antimicrobial resistance and foodborne outbreaks in the European Union in 2006

- ([http://www.efsa.europa.eu/efsa-locale-1178620753812\\_1178671312912.htm](http://www.efsa.europa.eu/efsa-locale-1178620753812_1178671312912.htm)). Accessed 25 January 2009.
39. **Baker MG, Sneyd E, Wilson NA.** Is the major increase in notified campylobacteriosis in New Zealand real? *Epidemiology and Infection* 2007; **135**: 163–170.
  40. **Nylen G, et al.** The seasonal distribution of *Campylobacter* infection in nine European countries and New Zealand. *Epidemiology and Infection* 2002; **128**: 383–390.
  41. **Sheppard SK, et al.** *Campylobacter* from retail poultry: MLST analysis and the origin of human infection. *Zoonoses and Public Health* 2007; **54**: 36–36.
  42. **Gormley FJ, et al.** Has retail chicken played a role in the decline of human campylobacteriosis? *Applied and Environmental Microbiology* 2008; **74**: 383–390.
  43. **Sopwith W, et al.** *Campylobacter jejuni* multilocus sequence types in humans, Northwest England, 2003–2004. *Emerging Infectious Diseases* 2006; **12**: 1500–1507.
  44. **Gilpin BJ, et al.** Survival of *Campylobacter* spp. in bovine faeces on pasture. *Letters in Applied Microbiology* 2009; **48**: 162–166.
  45. **Gilpin BJ, et al.** The transmission of thermotolerant *Campylobacter* spp. to people living or working on dairy farms in New Zealand. *Zoonoses and Public Health* 2008; **55**: 352–360.
  46. **Forbes KJ, et al.** *Campylobacter* immunity and coinfection following a large outbreak in a farming community. *Journal of Clinical Microbiology* 2009; **47**: 111–116.
  47. **Havelaar AH, et al.** Immunity to *Campylobacter*: its role in risk assessment and epidemiology. *Critical Reviews in Microbiology* 2009; **35**: 1–22.
  48. **Dingle KE, et al.** Sequence typing and comparison of population biology of *Campylobacter coli* and *Campylobacter jejuni*. *Journal of Clinical Microbiology* 2005; **43**: 340–347.
  49. **Best EL, et al.** Specific detection of *Campylobacter jejuni* from faeces using single nucleotide polymorphisms. *Epidemiology and Infection* 2007; **135**: 839–846.
  50. **Clark CG, et al.** Use of the Oxford multilocus sequence typing protocol and sequencing of the flagellin short variable region to characterize isolates from a large outbreak of waterborne *Campylobacter* sp. strains in Walkerton, Ontario, Canada. *Journal of Clinical Microbiology* 2005; **43**: 2080–2091.
  51. **Pope C, et al.** Epidemiology, relative invasive ability, molecular characterization, and competitive performance of *Campylobacter jejuni* strains in the chicken gut. *Applied and Environmental Microbiology* 2007; **73**: 7959–7966.
  52. **Taboada EN, et al.** Comparative genomic assessment of multi-locus sequence typing: rapid accumulation of genomic heterogeneity among clonal isolates of *Campylobacter jejuni*. *BMC Evolutionary Biology* 2008; **8**.
  53. **Hearnden M, et al.** The regionality of campylobacteriosis seasonality in New Zealand. *International Journal of Environmental Health Research* 2003; **13**: 337–348.