


Editorial

New guidance for self-harm: an opportunity not to be missed

Faraz Mughal, Fiona M. Burton, Harriet Fletcher, Karen Lascelles, Rory C. O'Connor, Sarah Rae, Alex B. Thomson and Nav Kapur



Summary

In this editorial we, as members of the 2022 NICE Guideline Committee, highlight and discuss what, in our view, are the key guideline recommendations (generated through evidence synthesis and consensus) for mental health professionals when caring for people after self-harm, and we consider some of the implementation challenges.

Keywords

Self-harm; NICE; suicide; psychosocial interventions; psychiatric nursing

Copyright and usage

© The Author(s), 2023. Published by Cambridge University Press on behalf of the Royal College of Psychiatrists.

Faraz Mughal (pictured) is a practising NHS general practitioner and NIHR Doctoral Fellow who researches the assessment, treatment and prevention of mental health conditions, self-harm and suicide in primary care. **Fiona Marie Burton** is a consultant in emergency medicine and major trauma based in the Queen Elizabeth University Hospital Glasgow. **Harriet Fletcher** is a consultant psychiatrist in psychotherapy at Leeds and York Partnership NHS Foundation Trust. **Karen Lascelles** is a mental health nurse consultant specialising in the areas of self-harm and suicide. **Rory O'Connor**, PhD is Professor of Health Psychology, President of the International Association for Suicide Prevention and he leads the Suicidal Behaviour Research Laboratory at Glasgow University. **Sarah Rae** has experience of co-occurring mental health conditions and self-harming behaviours. She draws on this lived experience to improve care through her involvement in research. **Alex B. Thomson** is a consultant liaison psychiatrist at Northwick Park Hospital, Central and North West London NHS Foundation Trust. He is Interim Chair of the Royal College of Psychiatrists' Faculty of Liaison Psychiatry. **Nav Kapur** is Professor of Psychiatry and Population Health and an honorary consultant at the University of Manchester and Mersey Care NHS Foundation Trust, undertaking self-harm and suicide research for 25 years.

Self-harm is a common problem in patients who seek help from clinical services and is associated with a heightened risk of suicide.¹ In recent years, there have been increases in self-harm particularly in young people, but also in other groups, such as middle-aged men.² Reducing rates remains a key national suicide prevention priority in the UK.² Although there has been some progress in the quality of mental healthcare people receive for self-harm in the past few decades, there is still considerable room for improvement.

The National Institute for Health and Care Excellence (NICE) has updated its guidance for self-harm with the publication of the new 2022 *Self-Harm: Assessment, Management and Preventing Recurrence* (NG225) guideline.³ The new guidance incorporates the latest evidence and combines the previous self-harm guidelines (CG16 and CG133, on short- and long-term management respectively) so that the recommendations are available in a single publication for the first time. The guideline also considers the role of the wider system (primary care, emergency care, education and criminal justice settings) because people who harm themselves can present to a range of settings.

In this editorial we, as clinical, academic and lived-experience members of the 2022 NICE guideline committee, highlight and discuss what, in our view, are the key guideline recommendations (generated through evidence synthesis and consensus) for mental health professionals when caring for people after self-harm. We also consider some of the implementation challenges. Our aim is not to summarise the entire guidance, and we acknowledge that our approach may have led to some bias in the recommendations

included here. The full guideline is readily accessible via the NICE website.

The need for empathy

Mental health professionals work across a diverse range of service settings and will need to approach and interpret the guidance in different ways, but much of the guideline is concerned with fundamental principles of care. A consistent theme throughout is the need to provide compassionate, non-judgemental and empathic care to people who have self-harmed. That is not to say all care for self-harm is currently lacking in these elements. There are examples of excellent care but equally examples of care that is suboptimal. These overarching principles are perhaps even more important to implement for people who frequently self-harm. Such individuals describe negative experiences of receiving mental health treatment which can have a harmful impact on help-seeking and risk of suicide.⁴ All staff should have access to appropriate supervision, and mental health services need to ensure that the training for practitioners is developed with and co-delivered by people with lived experience, to emphasise sensitive person-centred care. Family involvement is key and people who have self-harmed and their family members or carers should receive information about self-harm, care and treatment options, support agencies and resources, as well as self-care, tailored to individual needs.

No place for aversive treatments

It is regrettable that past approaches to treatment of self-harm have included shaming, punitive treatment or withholding care, based on misguided theories related to eliciting attention. Although these approaches have no place in 21st-century practice, contemporary lived-experience accounts demonstrate that they still occur. The guideline restates the principle that there is no place for aversive or punitive treatment, excluding patients or withholding the care that they need, and this extends to the use of criminal punishment as a deterrent.³

Psychosocial assessment, not risk assessment

The guideline states that all people who have self-harmed should receive a psychosocial assessment at the earliest opportunity from a trained mental health professional to identify the circumstances

of the episode and strengths and needs of the person. This assessment should be conducted with hope and optimism, and should not be unduly delayed for medical treatment, with liaison psychiatry teams pivotal in delivering this in emergency departments within general hospitals. Some patients have described not being offered an assessment, which is alarming.⁴ The guideline includes detail of what the assessment should cover, but perhaps even more important than the ‘what’ (the content of the assessment) is the ‘how’: it should be carried out using a compassionate and collaborative approach to develop a shared understanding of why self-harm has occurred.

Mental health professionals are urged to not use risk assessment tools and scales and to not stratify risk into low, medium or high to predict future self-harm repetition or suicide. The guidance also clearly states that risk tools or risk stratification should not determine who should or should not be discharged or offered treatment. The positive predictive values of these tools and categories are poor: they provide false clinical reassurance and result in people in the low and medium risk groups being excluded from care.³ Instead, age-appropriate mental health professionals should focus on the person’s individual circumstances, strengths and needs and develop a risk formulation. This drawing together of the clinical narrative will directly inform care and treatment rather than assessment being an end in itself. Although it is potentially a huge implementation challenge, clinical leaders could help to facilitate a move from a ‘risk-focused’ to a ‘safety-focused’ culture in self-harm services by recognising the therapeutic benefit of sensitive comprehensive assessments rather than approaches that reduce individual experience to a risk category; moving from tick-box approaches to holistic risk formulation; and addressing patients’ needs rather than treating their risk scores.

Aftercare and intervention

Aftercare

Where the clinician feels there are ongoing safety concerns after the initial assessment, prompt aftercare should be arranged within 48 h of assessment, with the same professional if possible. This is the time when repeat self-harm is most likely and early follow-up should enhance service engagement, reduce hopelessness and decrease repeat self-harm and suicide. Continuity of care is valued by clinicians and patients, and it improves safety.³ The role of in-patient psychiatric admission was not specifically considered in this guideline. Although it can be helpful, even life-saving for some patients, views on the relative benefits versus harms vary.

There are significant workforce and resource constraints that affect how the recommendations on aftercare might practically be delivered in real-world mental health services. Closer working between liaison psychiatry, psychological therapy services and primary care teams is critical to develop new models of aftercare, including the consideration of digital delivery.

Interventions

The Cochrane review on the effectiveness of interventions for self-harm repetition was updated in 2021 and this informed NICE treatment recommendations.³ There are no pharmacotherapies recommended for self-harm *per se*, although pharmacological treatment can be used to treat underlying conditions. The guideline did not consider how treatment for conditions related to self-harm might be prioritised. Where patients have a specific mental illness associated with self-harm, treatment for that condition should be arranged according to clinical judgement and associated guidance. The guideline recommends that ‘cognitive-behavioural therapy (CBT)-

informed’ psychological treatments are offered for adults: they should be person-centred and be 4–10 sessions in duration. The scope of ‘CBT informed’ is broad and it encompasses a wide variety of therapeutic approaches, including problem-solving, interpersonal and cognitive approaches. In young people who frequently self-harm dialectical behaviour therapy adapted for adolescents (DBT-A) can be considered. The evidence for DBT-A, however, was limited by a lack of long-term follow-up data and participant samples.³

A challenge for mental health services is having the workforce to deliver these treatments when patients need them. The upskilling of nurses or other practitioners in mental health teams, collaborative working with third-sector agencies and building on existing provision of psychological therapies in primary care could provide some solutions. The guideline highlights the cost-effectiveness of these approaches – they will actually save money in the long term. Psychological treatments should not be withheld from people because of diagnosis, coexisting conditions or substance misuse.

Safety plans

The development of safety plans should be considered for people who self-harm. The fact that the NICE recommendation was ‘consider’ rather than a stronger recommendation reflected the lack of high-quality evidence from randomised controlled trials (RCTs); however, they are in widespread use, and recent reviews have highlighted potential benefits. Again the ‘how’ is at least as important as the ‘what’. Safety plans should be created collaboratively between the person who is self-harming and mental health professionals, with input from family members or carers where appropriate. They should be accessible to the person, mental health professionals and others who may be named as a source of support, such as the general practitioner, and regularly reviewed. They are dynamic documents that benefit from ongoing updates and should be adapted as needs and circumstances change. There is a role for mental health professionals beyond psychiatric practice: the guidance recommends that professionals should help schools in developing support plans for students after self-harm.

Harm minimisation approaches designed to avoid, delay and reduce future self-harm episodes or complications are extremely contentious. The guideline steers clear of recommendations on ‘safer self-harm’ and suggests that professionals should always instil hope that a person can move away from self-harm as a coping strategy. The guideline does acknowledge the lack of evidence in this area. However, distraction techniques, wound hygiene or simple educational advice, such as on the influence of alcohol and illicit substances on self-harm, may be appropriate in the overall approach to the person’s ongoing recovery, after being agreed with the person and family.

Research recommendations

The guideline recommends five areas that would benefit from further research to identify effective interventions. These include models of care for young people; approaches to assessment in non-specialist settings; admission to hospital after self-harm; and psychological interventions and harm minimisation approaches.

Our business and everyone’s business

All mental health professionals will encounter people after self-harm in their daily practice and mental health services remain a key setting for intervention and prevention. However, the updated NICE self-harm guideline reminds us that self-harm can present anywhere across the health and social care system: in education,

in criminal justice and in community settings. This is a public mental health concern that not only has an impact on mental health services but also wider society.

These recommendations emphasise the importance of compassionate person-centred care, which all patients deserve. They highlight the need for a psychosocial assessment for each self-harm episode, the lack of usefulness of risk tools and scales, and the importance of considering the individual needs and safety of the patient and how safety can be optimised.

Implementation of the guideline is likely to require training across services (perhaps based on existing or updated competency frameworks), protected and regular supervision, and new models of care. These models need to be appropriately tested and quality improvement (QI) approaches may be helpful. The recent Emergency Department Safety Assessment and Follow-up Evaluation 2 (ED-SAFE 2) cluster RCT, published since the NICE guideline, suggested that QI with an emphasis on safety planning led to almost a halving in acute presentations for suicidal behaviour.⁵

Adequate resources, preferably ring-fenced funding, will be required for services to develop new ways of managing self-harm, and in all such developments, the involvement of people with lived experience and the third sector is crucial. In the context of a 'permacrisis' where we are emerging from a pandemic into an ongoing economic storm and pressured healthcare systems, implementation will be challenging to say the least. Acting on, and promoting, the new NICE guidance for self-harm will improve care for patients who do not always get the standard of treatment they deserve. But the question is, will we realise this opportunity, or will it be lost?

Faraz Mughal , MPhil, FRCGP, School of Medicine, Keele University, Keele, UK; **Fiona M. Burton**, MRCS(Glas), FRCEM, Department of Emergency Medicine, Queen Elizabeth University Hospital, Glasgow, UK; **Harriet Fletcher**, MRCPsych, Medical Psychotherapy Service, Leeds and York Partnership NHS Foundation Trust, Leeds, UK; **Karen Lascelles** , RMN, MSc, Oxford Health NHS Foundation Trust, Oxford, UK; **Rory C. O'Connor** , PhD, Suicidal Behaviour Research Laboratory, School of Health and Wellbeing, University of Glasgow, Glasgow, UK; **Sarah Rae**, Norfolk and Suffolk NHS Foundation Trust, Norwich, UK; **Alex B. Thomson** , FRCPsych, Department of Liaison Psychiatrist, Northwick Park Hospital, Central and North West London NHS Foundation Trust, London, UK; **Nav Kapur** , MD, FRCPsych, Centre for Mental Health and Safety, Manchester Academic Health Sciences Centre, University of Manchester, Manchester, UK; Mersey Care NHS Foundation Trust, Manchester, UK; and NIHR Greater Manchester Patient Safety Research Collaboration, University of Manchester, Manchester, UK

Correspondence: Faraz Mughal. Email: f.mughal@keele.ac.uk

First received 10 Mar 2023, final revision 21 Jul 2023, accepted 5 Aug 2023

Data availability

Data availability is not applicable to this article as no new data were created or analysed in this study.

Author contributions

F.M. and N.K. conceived the idea for the article. F.M. drafted the first version of the manuscript. All authors substantially commented on drafts and approved the final version for submission.

Funding

F.M. is funded by a National Institute for Health and Care Research (NIHR) Doctoral Fellowship (NIHR300957). N.K. is funded by the NIHR Greater Manchester Patient Safety Research Collaboration (NIHR204295). The views expressed in this article are those of the authors and not necessarily those of NICE, NHS, NIHR or the Department for Health and Social Care (DHSC).

Declaration of interest

All authors were members of the Guideline Committee for the NICE 2022 self-harm guideline, and N.K. was the expert topic advisor. N.K. is a member of the DHSC National Suicide Prevention Strategy Advisory Group, and a member of the *BJPsych* editorial board and did not take part in the review or decision-making process of this paper. R.O.C. is a Co-Chair of the Academic Advisory Group to the Scottish Government's National Suicide Prevention Leadership Group; an advisor on the development of the new Scottish self-harm strategy; a Trustee and Science Council Member of MQ Mental Health Research; President of International Association for Suicide Prevention; and a Trustee of James' Place. A.B.T. is Interim Chair of the Royal College of Psychiatrists' Faculty of Liaison Psychiatry.

References

- 1 Knipe D, Padmanathan P, Newton-Howes G, Chan LF, Kapur N. Suicide and self-harm. *Lancet* 2022; **399**: 1903–16.
- 2 Department of Health and Social Care. *Preventing Suicide in England: Fifth Progress Report of the Cross-Government Outcomes Strategy to Save Lives*. GOV.UK, 2021.
- 3 National Institute for Health and Care Excellence. *Self-Harm: Assessment, Management and Preventing Recurrence (NICE Guideline NG225)*. NICE, 2022 (<https://www.nice.org.uk/guidance/ng225>).
- 4 Quinlivan L, Gorman L, Littlewood DL, Monaghan E, Barlow SJ, Campbell S, et al. 'Wasn't offered one, too poorly to ask for one' – reasons why some patients do not receive a psychosocial assessment following self-harm: qualitative patient and carer survey. *Aust N Z J Psychiatry* 2022; **56**: 398–407.
- 5 Boudreaux ED, Larkin C, Vallejo Sefair A, Ma Y, Li YF, Ibrahim AF, et al. Effect of an emergency department process improvement package on suicide prevention: the ED-SAFE 2 cluster randomized clinical trial. *JAMA Psychiatry* 2023; **80**: 665–74.

Psychiatry in sacred texts

Equanimity for anger: creating space for difficult emotions

Michael Uebel 

The American novelist and journalist Christopher Morley (1890–1957) once declared that 'The size of a man is measured by the size of what makes him angry.' How anger affects us is proportional to the mental space in which we hold it. The more mentally capacious and elastic we are, the more anger, or any difficult emotion, loses its force. So, when asked what size he would be, Rabbi Pinchas of Koretz, in philosopher Martin Buber's account, responded to the question this way: 'Long ago,' he said, 'I conquered my anger and placed it in my pocket. When I have need of it, I take it out.' In his equanimity, Rabbi Pinchas does not banish anger, but acts always in such a way that he is bigger than it and can appropriately and flexibly access it.

© The Author(s), 2023. Published by Cambridge University Press on behalf of the Royal College of Psychiatrists

The British Journal of Psychiatry (2023)
223, 503. doi: 10.1192/bjp.2023.88