

## Research Article

**Cite this article:** Sbatella GM, Adjesiwor AT, Kniss AR, Stahlman PW, Westra P, Moechnig M, Wilson RG (2019) Herbicide options for glyphosate-resistant kochia (*Bassia scoparia*) management in the Great Plains. Weed Technol. doi: [10.1017/wet.2019.48](https://doi.org/10.1017/wet.2019.48)

Received: 7 March 2019

Revised: 20 May 2019

Accepted: 22 May 2019

**Associate Editor:**

Amit Jhala, University of Nebraska, Lincoln

**Nomenclature:**

Glyphosate; kochia, *Bassia scoparia* (L.) A. J. Scott, KCHSC; corn, *Zea mays* L. ZEAMX; soybean, *Glycine max* (L.) Merr. GLXMA; sugar beet, *Beta vulgaris* L. BEAVX; wheat, *Triticum aestivum* L. TRZAX.

**Keywords:**

Crop rotation; environment; herbicide variability.

**Author for correspondence:**

Andrew Kniss, Email: [akniss@uwyo.edu](mailto:akniss@uwyo.edu)

†Deceased.

# Herbicide options for glyphosate-resistant kochia (*Bassia scoparia*) management in the Great Plains

Gustavo M. Sbatella <sup>1†</sup>, Albert T. Adjesiwor <sup>2</sup>, Andrew R. Kniss <sup>3</sup>, Phillip W. Stahlman <sup>4</sup>, Phil Westra <sup>5</sup>, Michael Moechnig <sup>6</sup> and Robert G. Wilson <sup>7</sup>

<sup>1</sup>Assistant Professor, Department of Plant Sciences, Powell Research and Extension Center, University of Wyoming, Powell, WY, USA; <sup>2</sup>Postdoctoral Research Associate, Department of Plant Sciences, University of Wyoming, Laramie, WY, USA; <sup>3</sup>Professor, Department of Plant Sciences, University of Wyoming, Laramie, WY, USA; <sup>4</sup>Professor, Agricultural Research Center–Hays, Kansas State University, Hays, KS, USA; <sup>5</sup>Professor, Department of Bioagricultural Sciences and Pest Management, Colorado State University, Fort Collins, CO, USA; <sup>6</sup>Associate Professor, South Dakota State University, Brookings, SD, USA; current: Corteva Agrisciences, Toronto, SD, USA and <sup>7</sup>Professor, Panhandle Research and Extension Center, University of Nebraska–Lincoln, Scottsbluff, NE, USA

**Abstract**

Kochia is one of the most problematic weeds in the United States. Field studies were conducted in five states (Wyoming, Colorado, Kansas, Nebraska, and South Dakota) over 2 yr (2010 and 2011) to evaluate kochia control with selected herbicides registered in five common crop scenarios: winter wheat, fallow, corn, soybean, and sugar beet to provide insight for diversifying kochia management in crop rotations. Kochia control varied by experimental site such that more variation in kochia control and biomass production was explained by experimental site than herbicide choice within a crop. Kochia control with herbicides currently labeled for use in sugar beet averaged 32% across locations. Kochia control was greatest and most consistent from corn herbicide programs (99%), followed by soybean (96%) and fallow (97%) herbicide programs. Kochia control from wheat herbicide programs was 93%. With respect to the availability of effective herbicide options, glyphosate-resistant kochia control was easiest in corn, soybean, and fallow, followed by wheat; and difficult to manage with herbicides in sugar beet.

**Introduction**

Kochia is an invasive annual forb native to Asia, introduced into the United States at the end of the 1800s as an ornamental from Europe (Friesen et al. 2009; Whitson et al. 2000). Kochia is widespread throughout the northwestern United States, where it is found growing in grasslands, pastures, prairies, roadsides, ditch banks, wastelands, floodplains, riparian habitats, and cultivated fields (Casey 2009; Ou et al. 2018; Stubbendieck et al. 2003). Kochia was ranked among the most serious weed species in the northwest United States due to its high rate of spread (Forcella 1985). Factors contributing to the distribution and troublesome nature of kochia include seed germination characteristics, high seed production, and an effective tumbleweed dispersal mechanism (Beckie et al. 2016; Dille et al. 2017; Everitt et al. 1983).

Kochia is a major concern across the western United States, because it is competitive with many crop species. Features such as early-season emergence, rapid growth, and drought tolerance confer upon kochia a unique competitive ability in many western U.S. cropping systems (Friesen et al. 2009; Schwinghamer and Van Acker 2008). In surveys by Weed Science Society of America members, kochia ranked as one of the top six most troublesome weeds of alfalfa (*Medicago sativa* L.), canola (*Brassica napus* L.), pulse crops, sugar beet, corn, and spring grains (Van Wychen 2016, 2017). Yield loss from kochia competition has been documented in many crops, including oats (*Avena sativa* L.), wheat, sugar beet, sunflower (*Helianthus annuus* L.), and alfalfa (Calpouzou et al. 1980; Friesen et al. 1993; Lewis and Gulden 2014; Manthey et al. 1996; Mesbah et al. 1994). In addition to these competitive traits, kochia is adapted to a variety of common agricultural practices. For example, kochia is very responsive to nitrogen and thrives under different tillage practices and crop rotations (Anderson et al. 2007; Friesen et al. 2009; Lugg et al. 1983; Miller 1986; Watson et al. 2001). Therefore, effective kochia management is a priority in most crop production systems.

Herbicides are a major component of kochia control programs, particularly in commercial crop production system. The outcome of relying on herbicides, combined with kochia characteristics such as high genetic diversity among and within populations, has led to weed population shifts and to the evolution of herbicide-resistant populations (Friesen et al. 2009; Godar et al. 2015; Mengistu and Messersmith 2002; Sbatella and Wilson 2010; Wilson et al. 2007). Kochia has evolved resistance to at least four herbicide sites of action, including acetolactate

© Weed Science Society of America 2019. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.



synthase (ALS) inhibitors, synthetic auxins, photosystem II (PSII) inhibitors, and 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) inhibitors (Bell et al. 1972; Foes et al. 1999; Heap 2019; Saari et al. 1990). Kochia populations resistant to PSII herbicides have been reported in Colorado, Illinois, Indiana, Iowa, Kansas, Montana, Nebraska, North Dakota, and Wyoming (Heap 2019). EPSPS-inhibitor-resistant populations were also found in Colorado, Idaho, Kansas, Montana, Nebraska, Oklahoma, Oregon, and Wyoming. In addition, populations resistant to synthetic auxins have been reported in Colorado, Idaho, Kansas, Montana, Nebraska, and North Dakota (Heap 2019). ALS-inhibitor-resistant kochia populations have been reported in all the states listed above and in Michigan, New Mexico, South Dakota, Texas, Utah, Wisconsin, and Washington. In addition to this widespread resistance, resistance to herbicides with multiple sites of action has been reported for kochia populations in several states (Heap 2019; Kumar et al. 2019). At least one population in Kansas has been shown to be resistant to all four of these herbicide sites of action (Varanasi et al. 2015).

Glyphosate-resistant (GR; EPSP synthase inhibitor) kochia is widespread in the Great Plains (Kumar et al. 2019). Surveys conducted from 2011 to 2014 showed that 39% to 60% of kochia populations in eastern Colorado were resistant to glyphosate (Westra 2016). Although it is unclear what proportion of crop fields in each state have confirmed cases of GR kochia, there have been 13 cases of GR kochia in cereals, including corn, wheat, and sorghum [*Sorghum bicolor* (L.) Moench] in 12 states, 4 cases in soybean in 4 states, and 3 cases in sugar beet in 3 states (Heap 2019). GR kochia is especially concerning to farmers in the Central and Northern Great Plains region, because glyphosate is relied upon heavily in many cropping systems. GR corn and soybean are common rotational crops in the eastern and southern part of the region (Kansas, Nebraska, and South Dakota) for both irrigated and rainfed conditions. Glyphosate-resistant sugar beet is grown under irrigation in several states in the United States, including Wyoming, Colorado, and Nebraska, and kochia is among the most troublesome and difficult to control weeds in sugar beet. Winter wheat–fallow is a common rotation in nonirrigated portions of this region. Glyphosate remains the primary weed control tool in GR crops (Kniss 2018) and in the fallow and postharvest phase of the wheat–fallow rotation; therefore, alternatives to glyphosate are needed to manage GR kochia populations in all of these cropping systems. The objective of this study was to evaluate non-glyphosate herbicide options for kochia control that are available in commonly grown crops and fallow fields in the Central and Northern Great Plains.

## Materials and Methods

Field studies (a total of 10 sites) were conducted across five states (Lingle, WY; Fort Collins, CO; Hays, KS; Scottsbluff, NE; and Brookings, SD) over 2 yr (2010 and 2011). Soils of the experimental sites were loam in Brookings and Fort Collins, silty clay loam in Hays, silt loam in Lingle, and sandy loam in Scottsbluff. The pH of the soils ranged from 6.8 to 8.2. Kochia populations at all study sites were glyphosate susceptible. Herbicide treatments were applied to a fallow site with no crop established to evaluate herbicide efficacy while avoiding the confounding factor of differences in crop competition; kochia control was a function of herbicide treatments only. Three non-glyphosate herbicide treatments were chosen for each of five common cropping scenarios in the region: winter

wheat, fallow, corn, soybean, and sugar beet. Herbicide treatments were chosen based on predicted efficacy on GR kochia, cost, and crop-rotation concerns. Although not exhaustive, the chosen herbicide programs were a representative sample of possible herbicide combinations that would provide the best kochia control within each crop of interest (Table 1). A glyphosate treatment and a nontreated control were also included at each experimental site for a total of 17 treatments per site.

Visual estimates of kochia control were made approximately 30 d after the final POST herbicide was applied using a scale of 0 (no visible kochia injury or apparent density reduction) to 100 (complete death or absence of kochia in the plot). Aboveground kochia biomass production was measured by cutting all kochia at the soil surface in a 1- to 3-m<sup>2</sup> area per plot (depending on location) approximately 60 d after the final POST herbicide was applied at all sites, except Kansas in 2011 or South Dakota in both years (biomass was not collected at those three sites).

Data analyses were performed in R statistical software v. 3.4.4 (R Core Team 2018) using the *lmer* and *glmer* functions in the *lme4* package (Bates et al. 2015). Visual estimates and biomass production were analyzed using a mixed-effects model, in which crop scenario was considered a fixed effect, and experimental site (combination of state and year) and herbicide treatment within a crop scenario were considered random effects. This approach was used because the herbicide treatments (Table 1) were considered a representative sample of kochia control that could be possible within each crop scenario of interest, and the main interest was whether GR kochia could be controlled adequately with herbicides within each crop scenario. Herbicide treatments within each crop scenario were then compared to determine whether there were differences between herbicide programs within a crop scenario. For this comparison, a similar mixed-effects model was used, but herbicide treatment was considered a fixed effect. Post hoc Tukey-adjusted pairwise treatment comparisons were performed at an alpha of 0.05 using the *emmeans* package (Lenth 2018).

## Results and Discussion

Experimental site explained more variation in control estimates compared with herbicide treatment within the crop scenario; the crop scenario for which herbicide treatments were registered explained the greatest amount of kochia control variance (Table 2). This suggests that the environment (and possibly kochia biotype) could have a great influence over kochia control. Experimental site explained more of the variance in kochia biomass production than either crop or herbicide within a crop (Table 2). Kochia biomass production is limited in nonirrigated fallow systems by soil water availability, which was expected to differ between sites.

The five states (Wyoming, Colorado, Kansas, Nebraska, and South Dakota) where the studies were conducted were expected to have substantially different environmental conditions, especially temperature and humidity, which are known to affect weed control with herbicides. For example, warmer temperatures can improve herbicide absorption and efficacy (Ganie et al. 2017; Schultz and Burnside 1980). However, kochia control with glyphosate and dicamba can be reduced at elevated (32.5/22.5 C day/night) temperatures (Ou et al. 2018). Ou et al. (2018) recommended that for best control, glyphosate and dicamba must be applied at day/night temperatures around 25/15 C. Herbicide tolerance can also differ among susceptible

**Table 1.** Herbicide treatments used in this study.

Herbicide	Trade name	Rate	Timing	Registered crop
None	—			
Glyphosate	Roundup PowerMax <sup>a</sup>	860 ae	5-cm kochia	
Ethofumesate phenmedipham + desmedipham + ethofumesate	Nortron <sup>b</sup> SC <sup>b</sup> Progress <sup>b</sup>	1,120	PRE 5-cm kochia	Sugar beet treatment 1
Ethofumesate phenmedipham + desmedipham + ethofumesate + triflurosulfuron	Nortron <sup>b</sup> SC <sup>b</sup> Progress <sup>b</sup>	1,120	PRE 5-cm kochia	Sugar beet treatment 2
Ethofumesate phenmedipham + desmedipham + ethofumesate + triflurosulfuron	UpBeet <sup>c</sup> Nortron <sup>b</sup> SC <sup>b</sup> Progress <sup>b</sup>	17 1,120	PRE 3 applications: 2.5-cm kochia, 7 d later, 14 d later	Sugar beet treatment 3
Dicamba	UpBeet <sup>c</sup> Clarity <sup>d</sup>	9 280 ae	5-cm kochia	Fallow treatment 1
Saflufenacil	Sharpen <sup>d</sup>	25	5-cm kochia	Fallow treatment 2
2,4-D + carfentrazone	Rage™ D-Tech <sup>e</sup>	550 ae 18	5-cm kochia	Fallow treatment 3
Isoxaflutole	Balance <sup>b</sup> Flex <sup>b</sup>	9	PRE	Corn
tembotrione + atrazine	Laudis <sup>b</sup> AAtrex <sup>f</sup>	92 280	5-cm kochia	treatment 1
Acetochlor + atrazine	Degree Xtra <sup>g</sup>		PRE	Corn treatment 2
topramezone Saflufenacil + dimethenamid-P dicamba + diflufenzopyr	Impact <sup>g</sup> Verdict <sup>d</sup>	18	5-cm kochia PRE	Corn treatment 3
Sulfentrazone + imazethapyr	Status <sup>d</sup>		5-cm kochia	
Authority Assist <sup>g</sup>	Authority Assist <sup>g</sup>		PRE	Soybean treatment 1
fluthiacet	Cadet <sup>g</sup>	6	5-cm kochia	
Flumioxazin	Valor <sup>g</sup> SX <sup>h</sup>	90	PRE	Soybean
lactofen	Cobra <sup>h</sup>	218	5-cm kochia	treatment 2
Metribuzin + S-metolachlor	Boundary <sup>g</sup> (2 pt)		PRE	Soybean treatment 3
lactofen	Cobra <sup>h</sup>	218	5-cm kochia	
Pyrasulfatole + bromoxynil	Huskie <sup>b</sup>		5-cm kochia	Wheat treatment 1
Fluroxypyr + bromoxynil	Starane <sup>g</sup> NXT <sup>i</sup>	ae	5-cm kochia	Wheat treatment 2
Dicamba + thifensulfuron + tribenuron + metsulfuron	Agility <sup>g</sup> SG <sup>c</sup>		5-cm kochia	Wheat treatment 3

<sup>a</sup> Monsanto Company, St Louis, MO.<sup>b</sup> Bayer CropScience, Research Triangle Park, NC.<sup>c</sup> DuPont, Wilmington, DE.<sup>d</sup> BASF, Research Triangle Park, NC.<sup>e</sup> FMC, Philadelphia, PA.<sup>f</sup> Syngenta, Greensboro, NC.<sup>g</sup> AMVAC Chemical Corporation, Los Angeles, CA.<sup>h</sup> Valent, Walnut Creek, CA.<sup>i</sup> Dow AgroSciences, Indianapolis, IN.**Table 2.** Variance components for crop, herbicide within crop, and experimental site for kochia control and kochia biomass production.

Source of variation	Estimated SD of effects	
	Control	Biomass production
Herbicide treatment within crop	0.475	15.63
Crop	1.818	76.49
Experimental site	1.143	133.26

weed populations. For example, a 2.8- to 4.2-fold difference in glyphosate susceptibility has been demonstrated among non-resistant common lambsquarters (*Chenopodium album* L.) accessions (Coburn 2017).

One of the challenges in weed management is dealing with the often-unpredictable variability caused by environmental or biological differences between sites. With the herbicide programs selected, those registered for corn and soybean demonstrated less variability in kochia control compared with fallow, wheat, and sugar beet (Figure 1). This suggests that crop choice is at least as important as herbicide choice within a crop when developing recommendations for GR kochia control; and in fact, crop explained more variance in kochia control compared with either experimental site or herbicide within a crop (Table 2). The potential for kochia control failure was relatively low for corn, regardless of herbicide program (median kochia control = 98%), whereas there was no herbicide program evaluated in sugar beet that provided greater than 86% control at any site, with a median kochia control of 39% across all sites (Figure 1). Thus, effective long-term GR kochia management in sugar beet will likely depend on rotating with crops such as corn and soybean, for which effective herbicides are available. Long crop-rotation restrictions will prove challenging, as only one corn herbicide program and one soybean herbicide program evaluated in this study would allow planting of sugar beet the following year.

Herbicide choice within a crop influenced visually assessed kochia control in soybean, wheat, and fallow, but not in corn or sugar beet (Supplementary Table 1; Figure 2). Kochia control with herbicides currently labeled for use in sugar beet averaged 32% across locations (Table 3), and the biomass produced was 2- to 8-fold greater following sugar beet herbicides than the biomass measured in other crop herbicide treatments (Table 3). This was expected, because sugar beet is one of the crops with the fewest herbicides available, and poor kochia control in conventional sugar beet was a primary factor in the rapid and widespread adoption of GR sugar beet (Fernandez-Cornejo et al. 2016; Kniss 2010). Thus, the evolution of GR kochia populations is likely to have the greatest impact on sugar beet production among the cropping systems considered in this research.

Conversely, there are diverse groups of herbicides registered for use in the other crop scenarios in this analysis, making it possible to select herbicides with better kochia control efficacy. This was especially true for corn and soybean herbicide treatments, for which median kochia control across experimental sites was greater than 90% (Figure 1), and kochia biomass was reduced by more than 84% (Table 3) compared with the nontreated. The corn and soybean herbicide treatments included combinations of PRE plus POST herbicide applications. The PRE herbicides used in corn and soybean have been shown to provide good kochia control. For example, acetochlor + atrazine provided 100% kochia control for up to 8 wk after treatment (Kumar and Jha 2015), and PRE application of sulfentrazone and flumioxazin in GR soybean provided 81% to 92% and 88% to 98% kochia control, respectively, within 10 wk after treatment (Hulse 2012).

This work has shown that the selected herbicides for the corn scenario provided the most efficacious and consistent control of kochia. But it is important to note the presence of crops would have further reduced kochia density and biomass production in most of the crop scenarios we evaluated, especially corn, soybean, and wheat. Kochia is an early-emerging annual weed, so crop planting date influences kochia population and density. A crop such as

winter wheat that is established before kochia emerges will have a competitive advantage, and kochia would likely have been reduced further by wheat herbicides if a wheat crop were present. Previous research in Montana, Wyoming, and


Nebraska has shown crop rotations with wheat can significantly reduce kochia density and seed production (EM Mosqueda et al., unpublished data). Likewise, in corn and soybean, at least some of the kochia would be controlled during field preparation (preplant burndown herbicides or tillage), and this could further reduce kochia density to be controlled with PRE and POST herbicides.

In contrast, a majority of kochia seeds emerge after sugar beet is planted, and sugar beet is a poor competitor with kochia, because it is a low-stature rosette-forming crop, exacerbating poor kochia control provided by available herbicides. Corn, soybean, and wheat have much denser canopies forming early in the growing season, making them good competitors with kochia. In a previous study, inclusion of competitive crops (corn or barley [*Hordeum vulgare* L.]) and a late-planted crop (dry bean [*Phaseolus vulgaris* L.]) reduced kochia density by 98% (Lim et al. 2018). Thus, in managing GR kochia, the choice of crop is very important. This explains why diverse crop rotations and effective herbicide mixtures are a good strategy for kochia control.


**Table 3.** Kochia control and biomass production estimated marginal means as affected by herbicide treatments registered for use in five different crop scenarios.<sup>a</sup>

Crop	Control	Biomass production
	%	g m <sup>-2</sup>
Corn	99 a	29 a
Soybean	96 ab	54 a
Sugar beet	32 c	230 b
Wheat	93 b	95 a
Fallow	97 ab	93 a
Nontreated	—	340 c

<sup>a</sup> Estimated marginal means within a column followed by the same letter are not significantly different according to Tukey-adjusted pairwise comparisons ( $\alpha = 0.05$ ).



**Figure 1.** Kochia control ratings 30 d after final application of herbicide treatments labeled for corn, soybean, fallow, wheat, and sugar beet. Each point represents one plot at one site in one year. Numbers next to box plots are the median kochia control from herbicide treatments registered for each crop. Individual herbicide treatments are described in Table 1.



**Figure 2.** Kochia control provided by each herbicide treatment registered for use in five different crop scenarios (corn, soybean, fallow, wheat, sugar beet) across five states and 2 yr. Each point represents the visible kochia control estimate 30 d after final herbicide application in one plot. Individual herbicide treatments are described in Table 1.

**Supplementary material.** To view supplementary material for this article, please visit <https://doi.org/10.1017/wet.2019.48>

**Author ORCIDs.** Andrew R. Kniss,  0000-0003-2551-4959  
Albert T. Adjesiwor,  0000-0002-0047-8771

**Acknowledgments.** Partial funding for this research was provided by Monsanto Company, and Monsanto employees were involved in the design of this research. In addition, funding has been previously provided to the University of Wyoming, Colorado State University, University of Nebraska–Lincoln, South Dakota State University, and Kansas State University in support of the authors' research programs by one or more of the following companies that produce and have financial interests in the herbicides used in this research: Bayer CropScience, DuPont, BASF, FMC, Syngenta, AMVAC Chemical Corporation, Valent, and Dow AgroSciences; these companies were not involved in this research.

## References

- Anderson RL, Stymiest CE, Swan BA, Rickertsen JR (2007) Weed community response to crop rotations in western South Dakota. *Weed Technol* 21: 131–135
- Bates D, Mächler M, Bolker B, Walker S (2015) Fitting linear mixed-effects models using lme4. *J Stat Softw* 67:1–48
- Beckie HJ, Blackshaw RE, Hall LM, Johnson EN (2016) Pollen- and seed-mediated gene flow in kochia (*Kochia scoparia*). *Weed Sci* 64:624–633
- Bell AR, Nalewaja JD, Schooler AB (1972) Response of kochia selections to 2, 4-D, dicamba, and picloram. *Weed Sci* 20:458–462
- Calpouzos CA, Lee GA, McNeal CD (1980) Survey of weed complexes in alfalfa seed fields in Idaho. Pages 127–132 in *Proceedings of the Western Society of Weed Science*. Salt Lake City, UT: Western Society of Weed Science.
- Casey PA (2009) Plant Guide for Kochia (*Kochia scoparia*). Manhattan, KS: USDA–Natural Resources Conservation Service, Kansas Plant Materials Center. 5 p
- Coburn CW (2017) Influence of Experimental Methods on Herbicide Resistance Confirmation. Ph.D dissertation. Laramie, WY: University of Wyoming. 145 p
- Dille JA, Stahlman PW, Du J, Geier PW, Riffel JD, Currie RS, Wilson RG, Sbatella GM, Westra P, Kniss AR, Moechnig MJ, Cole RM (2017) Kochia (*Kochia scoparia*) emergence profiles and seed persistence across the central Great Plains. *Weed Sci* 65:614–625
- Everitt JH, Alaniz MA, Lee JB (1983) Seed germination characteristics of *Kochia scoparia*. *J Range Manage* 36:646–648
- Fernandez-Cornejo J, Wechsler SJ, Milkove DL (2016) The Adoption of Genetically Engineered Alfalfa, Canola and Sugarbeets in the United States. Washington, DC: U.S. Department of Agriculture, Economic Research Service Economic Information Bulletin Number 163. 22 p
- Foes MJ, Liu L, Vigue G, Stoller EW, Wax LM, Tranel PJ (1999) A kochia (*Kochia scoparia*) biotype resistant to triazine and ALS-inhibiting herbicides. *Weed Sci* 47:20–27
- Forcella F (1985) Final distribution is related to rate of spread in alien weeds. *Weed Res* 25:81–191
- Friesen LF, Beckie HJ, Warwick SI, Van Acker RC (2009) The biology of Canadian weeds. 138. *Kochia scoparia* (L.) Schrad. *Can J Plant Sci* 89:141–167
- Friesen LF, Morrison IN, Rashid A, Devine MD (1993) Response of a chlorsulfuron-resistant biotype of *Kochia scoparia* to sulfonylurea and alternative herbicides. *Weed Sci*. 41:100–106
- Ganie ZA, Jugulam M, Jhala AJ (2017) Temperature influences efficacy, absorption, and translocation of 2, 4-D or glyphosate in glyphosate-resistant and glyphosate-susceptible common ragweed (*Ambrosia artemisiifolia*) and giant ragweed (*Ambrosia trifida*). *Weed Sci* 65:588–602
- Godar AS, Stahlman PW, Jugulam M, Dille JA (2015) Glyphosate-resistant kochia in Kansas: EPSPS gene copy number in relation to resistance levels. *Weed Sci* 63:587–595
- Heap I (2019) The International Survey of Herbicide Resistant Weeds. <http://www.weedscience.org>. Accessed: April 15, 2019
- Hulse BM (2012) Kochia Control with Preemergence Herbicides in Soybeans; Dose Response of Three Kochia Populations to Glyphosate; and Response of Corn, Soybean, and Grain Sorghum to Saflufenacil. MS thesis. Manhattan, KS: Kansas State University. 58 p
- Kniss AR (2010) Comparison of conventional and glyphosate-resistant sugar-beet the year of commercial introduction in Wyoming. *J Sugar Beet Res* 47:127–134
- Kniss AR (2018) Genetically engineered herbicide-resistant crops and herbicide-resistant weed evolution in the United States. *Weed Sci* 66:260–273
- Kumar V, Jha P (2015) Effective preemergence and postemergence herbicide programs for kochia control. *Weed Technol* 29:24–34
- Kumar V, Jha P, Jugulam M, Yadav R, Stahlman PW (2019) Herbicide-resistant kochia (*Bassia scoparia*) in North America: a review. *Weed Sci* 67:4–15
- Lenth RV (2018) emmeans: Estimated Marginal Means, aka Least-Squares Means. R Package v. 1.2.3. <https://CRAN.R-project.org/package=emmeans>. Accessed: November 15, 2018.
- Lewis DW, Gulden RH (2014) Effect of kochia (*Kochia scoparia*) interference on sunflower (*Helianthus annuus*) yield. *Weed Sci* 62:158–165
- Lim CA, Mosqueda EG, Jha P, Kniss A, Sbatella GM, Lawrence NC (2018) Optimizing crop rotations to best utilize crop canopy effects on kochia seed bank reduction: a multi-site-year study. In *Proceedings of the 58<sup>th</sup> Annual Meeting*. Arlington, VA: Weed Science Society of America. <http://www.wssaabstracts.com/public/54/proceedings.html>. Accessed: June 5, 2019
- Lugg DG, Cuesta PA, Norcross GY (1983) Effect of nitrogen and phosphorus fertilization on yield and quality of *Kochia scoparia* grown in the greenhouse. *Commun Soil Sci Plant Anal* 14:859–876
- Manthey FA, Hareland GA, Zollinger RK, Huseby DJ (1996) Kochia (*Kochia scoparia*) interference with oat (*Avena sativa*). *Weed Technol* 10:522–525
- Mengistu LW, Messersmith CG (2002) Genetic diversity of kochia. *Weed Sci* 50:498–503.
- Mesbah A, Miller SD, Fornstrom KJ, Legg DE (1994) Kochia (*Kochia scoparia*) and green foxtail (*Setaria viridis*) interference in sugarbeets (*Beta vulgaris*). *Weed Technol* 8:754–759
- Miller SD (1986) Evaluation of Post-harvest Herbicide Treatments for Weed Control in Fallow. Western Society of Weed Science Research Report. San Diego, California. P 287
- Ou J, Thompson CR, Stahlman PW, Bloedow N, Jugulam M (2018) Reduced translocation of glyphosate and dicamba in combination contributes to poor control of kochia scoparia: evidence of herbicide antagonism. *Scientific Rep* 8:5330
- R Core Team (2018) R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Consulting. <http://www.R-project.org>. Accessed: November 15, 2018.
- Saari LL, Cotterman JC, Primiani MM (1990) Mechanism of sulfonylurea resistance in the broadleaf weed, *Kochia scoparia*. *Plant Physiol* 93:55–61
- Sbatella GM, Wilson RG (2010) Isoxaflutole shifts kochia (*Kochia scoparia*) populations in continuous corn. *Weed Technol* 24:392–396
- Schultz ME, Burnside OC (1980) Absorption, translocation and metabolism of 2, 4-D and glyphosate in hemp dogbane (*Apocynum cannabinum*). *Weed Sci* 28:13–20
- Schwinghamer TD, Van Acker RC (2008) Emergence timing and persistence of kochia (*Kochia scoparia*). *Weed Sci* 56:37–41
- Stubbendieck J, Coffin MJ, Landholt LM (2003) Weeds of the Great Plains. Lincoln, NE: Nebraska Dept of Agriculture. 605 p
- Van Wychen L (2016) 2016 Survey of the Most Common and Troublesome Weeds in Broadleaf Crops, Fruits & Vegetables in the United States and Canada. Weed Science Society of America National Weed Survey Dataset. [http://wssa.net/wp-content/uploads/2016-Weed-Survey\\_Broadleaf-crops.xlsx](http://wssa.net/wp-content/uploads/2016-Weed-Survey_Broadleaf-crops.xlsx). Accessed: June 15, 2018
- Van Wychen L (2017) 2017 Survey of the Most Common and Troublesome Weeds in Grass Crops, Pasture and Turf in the United States and Canada. Weed Science Society of America National Weed Survey Dataset. [http://wssa.net/wp-content/uploads/2017-Weed-Survey\\_Grass-crops.xlsx](http://wssa.net/wp-content/uploads/2017-Weed-Survey_Grass-crops.xlsx). Accessed: June 15, 2018

- Varanasi V, Godar A, Currie R, Dille JA, Thompson C, Stahlman P, Jugulam M (2015) Field evolved resistance to four modes of action of herbicides in a single kochia (*Kochia scoparia* L. Schrad.) population. *Pest Manag Sci* 71:1207–1212
- Watson PR, Derksen DA, Thomas AG, Turnbull GT, Blackshaw RE, Leeson JY, Legere A, Van Acker RC, Brandt SA, Johnston AM, Lafond GP, McConkey BG (2001) Weed Management and Ecology in Conservation Tillage Systems: Determination of Weed Community Changes in Conservation-Tillage Systems. Brandon, MB, Canada: Agriculture and Agri-Food Canada Weed Community Analysis Series Publ. Dow-2001-01. 229 p
- Westra EP (2016) Glyphosate-Resistant Kochia (*Kochia scoparia*) Management in the Central Great Plains and Western Canada. Ph.D dissertation. Fort Collins, CO: Colorado State University. 99 p
- Whitson TD, Burrill LC, Dewey SA, Cudney DW, Nelson BE, Lee RD, Parker R (2000) Weeds of the West. Laramie, WY: Western Society of Weed Science and University of Wyoming. 630 p
- Wilson RG, Miller SD, Westra P, Kniss AR, Stahlman PW, Wicks GW, Kachman SD (2007) Glyphosate induced weed shifts in glyphosate-resistant corn or a rotation of glyphosate-resistant corn, sugarbeet, and spring wheat. *Weed Technol* 21:900–909