November 1962

The Journal of Laryngology and Otology

EDITED BY G. H. BATEMAN

ASSISTANT EDITOR LIONEL TAYLOR

Contents

ON THE AETIOLOGY OF OTOSCLEROSIS .

R. F. OGILVIE and I. S. HALL

C. GLEDHILL

REFLECTIONS ON MY FIRST HUNDRED STAPEDECTOMIES

THE CLINICAL SIGNIFICANCE OF HISTOLOGICAL CHANGES IN THE OTOSCLEROTIC STAPES FOOT-PLATE . .

REPORT ON 102 CASES OF STAPEDECTOMY

CLINICAL RECORDS-

THREE CASES OF AURAL PATHOLOGY OF ANGLO-SAXON DATE

THREE CASES OF RHINITIS CASEOSA . . DERMOT J. DURCAN

A CASE OF TETANUS CAUSED BY A CHRONIC EAR INFECTION

FIBRO-SQUAMOUS EPITHELIAL PAPILLOMA FROM

GENERAL NOTES-

London Headley Brothers 109 Kingsway WC2

Annual Subscription £4/4/0 net, U.S.A. \$13 https://doi.org/10.1017/S0022215100060084 Published online by Cambridge University Press H. LUDMAN

J. D. K. DAWES and J. McG. JACKSON

CALVIN WELLS

P. V. WADSWORTH

Montbly, 10/- net post free

ADVERTISEMENTS

The Journal of Laryngology and Otology

(FOUNDED in 1887 by MORELL MACKENZIE and NORRIS WOLFENDEN)

EDITED BY G. H. BATEMAN

> ASSISTANT EDITOR LIONEL TAYLOR

1. Original articles which have not been published elsewhere are invited and should be sent to the Editor. They are considered for publication on the understanding that they are contributed to this Journal solely. Reproduction elsewhere, in whole or in part, is not permitted without the previous written consent of the Author and Editor and the customary acknowledgement must be made.

z. Manuscripts should be typewritten, on one side only of the paper, and well spaced. Captions to illustrations should be typed on a separate sheet and sent at the same time as original photographs, etc.

The Harvard system of recording references should be used, e.g. GREEN, C., and BROWN, D. (1951) J. Laryng., 65, 33. Abbreviations of Journals should follow the style recommended in World Medical Periodicals, published by World Health Organization, 1952.

It is most important that authors should verify personally the accuracy of

every reference before submitting a paper for publication. 3. Galley proofs and engraver's proofs of illustrations are sent to the author. Corrections, which should be kept to a minimum, must be clearly marked, and no extra matter added. Proofs should be returned within 5 days.

4. Illustration blocks are provided free up to the limit of f to per article; beyond this authors are expected to pay half the cost. Coloured illustrations will be charged in full to authors.

Blocks will normally be held by the Printers for three years after which they will be destroyed. Any author who has borne a part of the cost of his blocks is entitled to have these returned to him, but a request for this must be sent within three years of the appearance of the article, to HEADLEY BROTHERS, 109 Kingsway, London, WC2.

5. Orders for reprints must be sent when returning galley proofs, and for this purpose special forms are supplied.

6. Authors of original communications on Oto-Laryngology in other journals are invited to send a copy, or two reprints, to the *Journal of Laryngology*. If they are willing, at the same time, to submit their own abstract (in English, French, Italian, or German) it will be welcomed.

7. Editorial communications may be addressed to THE EDITOR, Journal of Laryngology, c/o HEADLEY BROTHERS, 109 Kingsway, London, WC2.

8. The annual subscription is four guineas sterling (U.S.A. \$13) post free,

and is payable in advance. 9. Single copies will be on sale at 10s. od. each; copies of parts up to Vol. LXIII which are available may be purchased at 7s. 6d. each.

10. All subscriptions, advertising and business communications should be sent to the publishers, HEADLEY BROTHERS, 109 KINGSWAY, LONDON, WC2.

United States of America

Orders for this Journal may be sent through local bookseller, or to STECHERT-HAFNER, INC., 31-33 East 10th Street, New York, or direct to the publishers, HEADLEY BROTHERS, 109 KINGSWAY, LONDON, WC2, England.

Journal of Laryngology and Otology, 1962.

CONTENTS

PA	GE
ON THE AETIOLOGY OF OTOSCLEROSIS. R. F. Ogilvie and I. S. Hall (Edinburgh)	41
Reflections on my First Hundred Stapedectomies. C. Gledhill (Northampton) 8	97
THE CLINICAL SIGNIFICANCE OF HISTOLOGICAL CHANGES IN THE OTOSCLEROTIC STAPES FOOT-PLATE. H. Ludman (London). 9	905
REPORT ON 102 CASES OF STAPEDECTOMY. J. D. K. Dawes and J. McG. Jackson (Newcastle-upon-Tyne))2 I
CLINICAL RECORDS Three Cases of Aural Pathology of Anglo-Saxon Date. Calvin Wells (Norwich)	931
Three Cases of Rhinitis Caseosa. Dermot J. Durcan (London) . $\boldsymbol{\varsigma}$	934
A Case of Tetanus Caused by a Chronic Ear Infection. P. Wadsworth (Hove)	939
Fibro-squamous Epithelial Papilloma from Upper Alveolus. A. Mackenzie Ross (Bournemouth)	940
General Notes	943
For Advertisement space in this Journal apply to:	

HEADLEY BROTHERS Ltd, 109 Kingsway London WC 2

ROYAL EAR HOSPITAL PATTERN ULTRASONIC EQUIPMENT

for the conservative surgery of Ménière's Disease. (Arslan Technique.) U.K. and U.S.A. patents applied for. Now available with the following special advantages:

- TRANSDUCER and cable will withstand sterilization in autoclave.
- Low operating voltage of transducer.
- APPLICATORS can be interchanged to suit anatomy of middle ear.
- Applicator tip normally $2\frac{1}{2}$ mm. in diameter.
- Intensities up to 100 Watts per sq. cm., giving ample power for pituitary irradiation in carcinoma.

GENERATOR operates on 90-250 volts 50 or 60 cycles.

Weight 50 lb. Size 21 by 14 by 12 in.

Tuning and intensity can be checked continuously.

REFERENCES:

Arslan. Proc. Fifth Int. Congress Oto-Rhino-Laryngol., Amsterdam, 1953.

Proc. roy. Soc. Med., 1958, 51, 617.

J. Laryng., 1960, LXXIV, 628.

Ibid., 1960, LXXIV, 730.

Ibid., 1961, LXXV, 345.

Ibid., 1962, LXXVI, 299.

Made in England by FRISTON ELECTRONICS LTD., 25 CHURCH ROAD, RICHMOND, SURREY.

'Diconal' Tablets for relief of intractable pain

Diconal' Tablets provide analgesia comparable with that produced by morphine. They are therefore of great value in carcinoma. Because the tablets are given by mouth and not by injection, the patient can be kept oblivious of the fact that his condition calls for so potent an analgesic.

'Diconal' Tablets contain dipipanone and cyclizine. Dipipanone relieves severe pain; cyclizine appreciably reduces the incidence of nausea and vomiting, even in patients new to this type of therapy.



'Diconal' Tablets are:

Of high value in relief of pain due to neoplastic disease, and other conditions where narcotic drugs would be employed.



A balanced formulation which permits flexibility in dosage, according to the needs of each patient.



Often effective in morphine- and pethidine-resistant cases.

'Diconal' Tablets are subject to the D.D.A.



Each tablet contains dipipanone hydrochloride, B.P.C. 10 mgm., and cyclizine hydrochloride, B.P.C. 30 mgm. Coloured red and scored for easy division.



6 NEW BOOKS . . . FOR LARYNGOLOGISTS AND OTOLOGISTS

APPLIED AUDIOLOGY FOR CHILDREN by D. M. C. Dale, School for Deaf and Partially Deaf Children, Kelston, Auckland, New Zealand. This is a down-to-earth, practical book with much to interest otologists and all who come in contact with children who have defective hearing. The author considers the relationship between the frequency components of speech sounds, the audiogram of the patient, and the frequency response of the hearing aid. He deals with factors which affect the reception of speech by hearing aid users and suggests practical measures by which children can be enabled to hear the most intelligible and continuous pattern of sound that their residual capacity to hear permits. Sept. '62, about 164 pp., about 79 il.

RADIATION THERAPY IN THE MANAGEMENT OF CANCERS OF THE ORAL CAVITY AND OROPHARYNX by Gilbert H. Fletcher and William S. MacComb. Physics Section by Robert J. Shalek and Marilyn Stovall. All of Univ. of Texas M. D. Anderson Hospital and Tumor Institute. The authors cover first principles underlying management with special emphasis on clinical varieties, integration of radiation and surgery, patient care, and management of complications. Applications of principles of radiotherapy to the specific management of squamous cell carcinomas are discussed in detail. Jan. '62, 408 pp., 346 il., \$16.50

RHINOSCLEROMA by Francisco Kerdel-Vegas and Jacinto Convit, both of Universidad Central de Venezuela; Bernard Gordon and Mauricio Goihmann, both of Vargas Hospital. All of Caracas, Venezuela. Rhinoscleroma is a chronic, infectious disease with a slow, non-fatal but often incapacitating evolution. It is important from a social as well as a scientific point of view. The information in this study from an endemic area of rhinoscleroma will be of value to all physicians. It covers every facet of this destructive condition including results of modern methods of treatment. Dec. '62

THE NOSE, PARANASAL SINUSES, AND EARS IN CHILDHOOD by Donald F. Proctor, The Johns Hopkins Univ. This second volume in Doctor Proctor's series will serve to clarify a great many controversial issues for both otolaryngologist and pediatrician. Recommended operative treatments are discussed in detail and fully illustrated. Hearing and speech problems are approached from the viewpoint of preventive medicine, treatment of diseases amenable to cure, and assistance to those children whose irremediable handicap leaves them with defective ability to communicate. Nov. '62

RECRUITING TEACHERS FOR THE DEAF by Joseph S. Rudloff, San Diego, Calif. Mr. Rudloff's work will be a boon to the administrator or supervisor whose difficult task it is to staff programmes for the deaf with qualified teachers. First, pertinent and basic considerations are reviewed. For what type of child do you recruit a teacher? What does the education of a deaf child mean and entail? In the second section the author presents results of a detailed study of teachers already in the field. Why did they enter? What is their educational background? Finally, logical and helpful conclusions are drawn from the study—guidelines that lend fact to an area previously left to opinion. July '62, 76 pp., 30 tables, \$4.25

SPEECH REHABILITATION OF THE LARYNGECTOMIZED by John C. Snidecor, Univ. of Calif. With a broad view of the laryngectomee as a total person, the authors consider the nature of the operation and its impact on the patient, personality in reference to rehabilitation, and the nature of the new speech and its limitations as contrasted with normal speech. Each contributor is well known for substantial research in his own area of specialization. March '62, 204 pp., 34 il., \$8.00

CHARLES C. THOMAS PUBLISHER 301-327 East Lawrence (Avenue · SPRINGFIELD · ILLINOIS · U.S.A.

Please mention The Journal of Laryngology and Otology when replying to advertisements

iv



Please mention The Journal of Laryngology and Otology when replying to advertisements

'PETERS' NEW CLINIC AUDIOMETER FOR PURE-TONE AND SPEECH



CONSIDER THESE FEATURES :

- A continuous frequency range with continuous automatic threshold compensation.
- Twin continuously variable attenuators accurate to I db.
- Narrow band masking of pure tones plus insert receiver.
- Manual or automatic completely silent interruption.
- Loudness Balance and Difference Limen Recruitment Tests.
- High Fidelity Speech Circuit with Accessories for Free Field Speech Audiometry.
- Full Spectrum of White Noise for Masking Speech.
- Completely Transistorised for reliability.

The Westrex Company Limited, who are leading suppliers of Deaf Education Equipment, have a nationwide servicing organisation. In the immediate future their engineers in the field will be co-operating in the speedy servicing of all "Peters" audiometers in Great Britain.

Write for a descriptive brochure giving full details of this magnificent instrument and ask for a demonstration from either:

WESTREX COMPANY Ltd. 152 COLES GREEN ROAD, LONDON, N.W.2

ALFRED PETERS & SONS Ltd. victoria works, gell street, sheffield 3



ORBENIN is indicated for immediate use in known or suspected resistant staphylococcal infections, and in mixed Gram-positive infections (e.g. infected burns) where resistant staphylococci may be present.

• Orbenin is stable to staphylococcal penicillinase and highly effective against 'resistant' staphylococci. Its activity is four times that of methicillin.

• Orbenin is also highly effective against other Gram-positive organisms, including streptococci, pneumococci and penicillin-sensitive staphylococci.


• Orbenin is bactericidal; it is acid-stable and well absorbed by either the oral or intramuscular routes.

Dosage by mouth is 500 mg. 6 hourly. Dosage by intramuscular injection is 250 mg. four to six hourly.



ORBENIN

(3-0. chlorophenyl-5-methyl-4-isoxazolylpenicillin sodium salt monohydrate) is a product of British research at the BEECHAM RESEARCH LABORATORIES LTD BRENTFORD · ENGLAND · *TELEPHONE*; ISLeworth 4111



And the reply was $\mu\alpha\rho\nu\delta\iota\alpha$ which is hardly surprising when you consider that in that helmet he could hardly have heard him anyway.

Fortunately, few, if any, patients wear Grecian helmets these days and most are delighted to have the name of an independent hearing aid consultant of impeccable reputation murmured tactfully into their ears.

Today more people than ever before can benefit from a carefully selected aid. But careful selection is important. INGRAMS, the Independent Suppliers, have twenty-six different makes – the largest selection in the country – for patients to choose from. Couple this with expert impartial advice and a high reputation and you have the reason why so many E.N.T. Specialists refer their patients to Ingrams.



Domiciliary visits in any part of the country

First clinical audiometer to provide 100% accurate audiograms



AMPLIVOX MODEL 82 world's most fully-developed clinical audiometer

It is now known that the disc type of earphone, used in conventional audiometers for air conduction testing and the application of masking noise, transfers the sound by bone conduction to the opposite ear at a level only 40 dB below the signal level. This means that up to 50% of audiograms taken at the present time with other audiometers may be invalid.

The Amplivox Model 82 Audiometer provides, for the first time, the means of taking accurate audiograms in cases of severe unilateral deafness through the provision of narrow band masking applied by insert earphone. This new masking technique is of special significance in bone conduction audiometry.

In addition to the exclusive narrow band masking feature, the Model 82 provides loudness balance test facilities, and *all* the other tests essential to proved pure tone audiometry.

AMPLIVOX LTD Medical Acoustic Division 80 New Bond St, London W.1.Tel: HYDe Park 9888



SEND FOR THIS DESCRIPTIVE BROCHURE Containing full information on the latest audiometric techniques in air and bone conduction.

In acute or chronic sinusitis, secondary infection in influenza and the common cold, and in sore throat

Sulfex Intranasal Drops

meet every therapeutic requirement

rapid concentration at the site of infection prolonged bacteriostasis safe, effective vasoconstriction

> Formula: 'Mickraform' sulphathiazol 5% in an isotonic aqueous medium with 'Paredrinex' (hydroxyamphetamine hydrobromide) 1 per cent.

SMITH KLINE & FRENCH LABORATORIES LTD Welwyn Garden City, Herts



*'Sulfex', 'Mickraform' and 'Paredrinex' are trade marks SX:PA22 (Col)