

# Effective Insomnia Treatments: Investigation of Processes in Mindfulness and Cognitive Therapy

Christopher William Lee,<sup>1</sup> Melissa J. Ree<sup>2</sup> and Mei Yin Wong<sup>3</sup>

<sup>1</sup> Faculty of Health and Medical Sciences, The University of Western Australia, Perth, Western Australia, Australia

<sup>2</sup> The Marian Centre, Perth, Western Australia, Australia

<sup>3</sup> School of Psychology and Exercise Sciences, Murdoch University, Perth, Western Australia, Australia

Understanding the underlying mechanisms of recovery from insomnia is an important goal for improving existing treatments. In a randomised controlled trial, 57 participants with insomnia disorder were given either cognitive therapy (CT) or mindfulness-based therapy (MBT) following 4 sessions of CBT. Each participant was assessed on process measures related to CT and MBT. MBT resulted in improvement on mindfulness process measures and the size of the improvement was significantly greater than achieved in the CT condition. Interestingly, CT and MBT both resulted in significant improvement on the cognitive process measures. Treatment outcome on the primary outcome measure (Insomnia Severity Index) was not predicted by type of treatment but was predicted by posttreatment scores on the cognitive process measures. The results suggest that changes in cognitive processes are especially important in treating insomnia, and that there are different therapeutic modalities through which this can be achieved.

■ **Keywords:** insomnia, mindfulness-based therapy, cognitive therapy, insomnia treatment, unhelpful beliefs about sleep

The economic burden of sleep disorders in Australia is estimated to be \$5.1 billion per year (Hillman & Lack, 2013), and insomnia is by far the most prevalent sleep disorder, representing 8 out of 10 sleep disorders managed in primary care (Charles, Harrison & Britt, 2009). Insomnia is a highly prevalent condition, with one in three people regularly having difficulty with their sleep and approximately 10% of the adult population suffering from the clinical disorder of insomnia (e.g., American Psychiatric Association, 2013; Singareddy et al., 2012). Insomnia disorder is defined as chronic difficulty falling asleep or returning to sleep following night-time awakening, at least three nights a week, with a significant impact on daytime functioning. Research has suggested that insomnia increases the risk of physical health problems, mood disorders, accidents, and poorer quality of life (Roth, 2007). Insomnia also results in impaired occupational performance (Daley et al., 2009) and higher healthcare usage and costs (Siversten, Krokstad, Mykletun, & Øverland, 2009). There is also evidence that treating insomnia can: (1) improve treatment outcome for comorbid psychiatric conditions such as depression (Perlis, Giles, Buysse, Tu, & Kupfer, 1997), and (2) protect against their development (e.g., Harvey et al., 2015, in the case of

Address for correspondence: Dr Melissa Ree, 11 Hamilton St, Subiaco WA 6008, Australia.

Email: [melissa.ree@optusnet.com.au](mailto:melissa.ree@optusnet.com.au)

**Behaviour Change** | Volume 35 | Number 2 | 2018 | pp. 71–90 | © The Author(s) 2018  
doi [10.1017/bec.2018.10](https://doi.org/10.1017/bec.2018.10)

bipolar disorder; and Franzen & Buysee 2008, in the case of depressive relapse). Given insomnia's high prevalence, disease burden and role in comorbid conditions, research to maximise effectiveness of treatment is an important priority.

While acute insomnia may be caused by life events, illness or distress about a current situation (Ellis, Gehrman, Espie, Riemann, & Perlis, 2012), there is general agreement that persistent insomnia is perpetuated, at least in part, by an interplay of cognitive and behavioural factors that are related to sleep, such as unhelpful sleep habits, worry about sleep, and unhelpful sleep-related beliefs. Unhelpful beliefs and thoughts perpetuate both anxiety about poor sleep and unhelpful behaviours that make poor sleep more likely. For example, if a person believes that they must attain 8 hours sleep every night and has the thought 'I won't cope tomorrow if I don't fall asleep soon', they will likely experience increased arousal that interferes with sleep. Further, they may sleep in longer after a poor night of sleep, then use extra caffeine, withdraw from daytime activities, or nap, contributing to poor sleep the following night. Cognitive Behavioural Therapy for Insomnia (CBT-I) is a multi-component treatment that includes a combination of psychoeducation, behavioural interventions and cognitive interventions, and seeks to address sleep-interfering thoughts, beliefs, and behaviours. The conclusion from recent meta-analyses is that CBT-I has an extensive evidence base attesting to its effectiveness (e.g., Okajima, Komada, & Inoue, 2011; Trauer, Qian, Doyle, Rajaratnam, & Cunnington, 2015), including in the case of insomnia comorbid with other conditions (see Geiger-Brown, Rogers, Liu, Ludeman, Downton, & Diaz-Abad, 2015). Trials also indicate that CBT-I is more effective than pharmacotherapy in the long term (Morin et al., 2009; Riemann & Perlis, 2009; Sivertsen et al., 2006). The weight of evidence for its efficacy, along with its preferable side-effect profile (compared to medication), has led to CBT-I being recommended as the first-line treatment of insomnia (e.g., Brasure et al., 2016; Qaseem, Kansagara, Forcica, Cooke, & Denberg, 2016). Robust research findings and clinical guidelines (e.g., Ree, Junge, & Cunnington, 2017) support the use of CBT-I as a first-line treatment; however, Australian research suggests that only 2% of patients seeking treatment for insomnia from their general practitioner (GP) will be referred for CBT-I (Charles, Harrison, & Britt, 2009).

Despite recognition of the effectiveness of CBT-I, there is also appreciation that not all individuals with insomnia achieve an adequate treatment response to this approach and that research into improving treatment outcomes is important. Research suggests that 20–30% of patients do not achieve clinically significant improvement following CBT-I (e.g., Bélanger et al., 2016; Morin & Benca, 2012). Further, only approximately 40% of CBT-I recipients achieve clinical remission (Morin & Benca, 2012). One promising way to maximise treatment effectiveness is via improved understanding of the mechanisms of change within treatments. Once mechanisms of change are well understood, the most critical treatment targets can be identified.

Two developments in psychologically based insomnia treatments are cognitive therapy (CT) and mindfulness-based therapy (MBT). Each have evidence bases suggesting that they offer promise for treating insomnia in their own right and/or by enhancing traditional CBT-I. These two treatments are reviewed below.

In the past, CBT-I was predominantly a behaviour therapy, with the explicit cognitive therapy element often being delivered in a single session (e.g., Carney & Edinger, 2008; Jacobs, 1998; Morin & Espie, 2003). However, following the recognition of the role of cognitive processes in insomnia (Harvey, 2002, 2005; Hiller, Johnston, Dohnt, Lovato, & Gradisar, 2015), a cognitive model of insomnia and its treatment has been

developed (Harvey, 2002). Harvey's (2002) cognitive model describes five maintaining processes that operate both day and night, namely: (1) worry that triggers arousal and distress (e.g., 'I won't cope tomorrow if I don't sleep well'), (2) selective attention to and monitoring for sleep-related threats (e.g., paying attention to signs of poor sleep such as daytime fatigue or night-time arousal), (3) overestimation of sleep and daytime deficits (perceiving any mistake made during the day as a result of poor sleep), (4) dysfunctional thoughts and beliefs about sleep (e.g., 'I must sleep 8 hours every night in order to function well'), and (5) counterproductive safety behaviours (e.g., increased time spent in bed, daytime napping, underexertion, and social withdrawal). The five cognitive processes mutually influence each other, thus creating a vicious cycle that maintains insomnia. Research has found that increased worry, dysfunctional beliefs, selective attention, and safety behaviours each distinguish normal sleepers from those with insomnia (Norell-Clarke, Jansson-Fröjmark, Tillfors, Harvey, & Linton, 2014). Additionally, there is evidence that CT designed to modify the cognitive maintaining factors is effective both as a stand-alone treatment (Harvey et al., 2014; Harvey, Sharpley, Ree, Stinson, & Clark, 2007) and as an addition to four sessions of CBT-I (Wong, Ree, & Lee, 2016). Indeed, the Harvey et al. (2014) trial, which compared CBT, CT, and behaviour therapy (BT) for insomnia, found that while BT produced faster treatment gains, CT produced longer lasting effects. CBT was found to be the treatment of choice, utilising both the faster impact of BT and the long-lasting impact of CT.

In addition to evidence supporting the effectiveness of CT for insomnia, there is evidence that change in *specific* cognitive maintaining factors relates to treatment outcome. For example, several studies suggest that the modification of dysfunctional sleep beliefs in the treatment of insomnia has beneficial effects for sleep improvement (e.g., Carney & Edinger, 2006; Eidelman et al., 2016; Morin, Blais, & Savard, 2002). Further, there is evidence that dysfunctional beliefs about sleep may interfere with behavioural treatment. For example, Cvengros, Crawford, Manber, and Ong (2015) found that patients with more dysfunctional beliefs about sleep were less likely to adhere to the recommendation of time in bed restriction, which requires individuals to match their time spent in bed overnight to their average perceived total sleep time. Several studies support that interventions that reduce worry produce favourable outcomes for sleep (e.g., Carney & Waters, 2006; Sunnhed & Jansson-Fröjmark, 2014). With respect to distorted perception of sleep (i.e., underestimation of total sleep time and overestimation of sleep onset latency), there is evidence that its correction in a therapeutic context results in reduced worry about sleep (Tang & Harvey, 2004). In terms of investigation of several cognitive processes within one treatment outcome study, the Harvey et al. (2007) case series of CT found that worry, dysfunctional beliefs, sleep-related monitoring, and safety behaviours all reduced following CT treatment. While these results are consistent with the cognitive model, the study was not designed to investigate the role of process variables, and so firm conclusions regarding causality cannot be made. Harvey, Dong, Belanger, and Morin (2017) conducted analysis that suggested a composite score of cognitive process measures mediated treatment outcome in CT.

As discussed earlier, the second potential innovation in psychological treatments for insomnia is MBT. In randomised controlled trials targeting insomnia, MBT was found to lead to greater symptom reduction compared to sleep hygiene education (Black, O'Reilly, Olmstead, Breen, & Irwin, 2015; Nakamura, Lipschitz, Kuhn, Kinney, & Donaldson, 2013), self-monitoring (Ong et al., 2014), and a waitlist control.

### Behaviour Change

MBT has also been compared to active treatments in two randomised controlled trials (Garland et al., 2014; Wong et al., 2016). In both cases, each treatment was found to result in significant improvement over time with no significant differences between MBT and CBT-I (Garland et al., 2014) and between MBT and CT (Wong et al., 2016).

Studies investigating the mechanisms by which MBT decreases insomnia symptoms have begun to appear in the literature (e.g., Garland, Zhou, Gonzalez, & Rodriguez, 2016). Since mindfulness has been found to be of benefit in general stress reduction and results in reduced arousal (Chiesa & Serretti, 2010), and because stress and physiological arousal are associated with insomnia (Hall et al., 2007), it is plausible that mindfulness may assist with insomnia by reducing the arousal associated with everyday stress.

In addition, mindfulness may also reduce the arousal associated with not sleeping. Ong, Ulmer, and Manber (2012) described how arousal can increase when someone has negative cognitions about their inability to sleep. They also describe that insomnia can be associated with metacognitive processes that further increase arousal via thoughts. This secondary arousal arises from the emotional valence, interpreted meaning, and level of attachment someone has with their sleep-related thoughts (Ong et al., 2012). MBT may be effective in reducing this metacognitive or secondary arousal. As Ong et al. (2012) note, MBT as a method for reducing arousal can be traced to Buddhist practices that encourage the notion that all things (including thoughts and emotions) come and go. Practice therefore encourages letting thoughts and feelings go, as opposed to forming a strong attachment to them (Ong et al., 2012). Accordingly, Shapiro, Carlson, Astin, and Freedman (2006) suggest that mindfulness helps people to observe thoughts rather than being involved with them. MBT therefore might help people by disengaging from ruminative processing and facilitate distancing from unhelpful cognitions (Garland et al., 2016).

An essential element of mindfulness is that attention is focused on an immediate experience, and in this way it can work to control attention biases (Larouche, Cote, Belisle, & Lorrain, 2014). This can be advantageous, as mindfulness can help redirect the person's attention to a neutral subject — for example, breathing. With practice, Larouche et al. (2014) suggest that it becomes easier to control attentional biases. Focusing on neutral stimuli rather than cues that can increase secondary anxiety, such as watching the clock or focusing on cues of sleepiness, has been proposed to facilitate better sleep (Ong et al., 2012).

In summary, treatment development and outcome research to date suggests that CBT-I, CT, and MBT-I are each effective in the treatment of insomnia. Further, from studies investigating the process of change, there is emerging evidence suggesting that these therapies produce their effects by modifying the theoretically proposed maintaining factors. It is tempting to draw the conclusion therefore that CT works by modifying the proposed cognitive maintaining factors and that MBT works by modifying the proposed mindfulness maintaining factors. However, the role of *both* sets of maintaining processes within each treatment has, to date, not been assessed. Mediation studies that permit investigation of the relationship between change in process measures and treatment outcome across treatment modalities are an important in understanding how these treatments have their effects.

The current study investigates the relationship between process measures and treatment outcome across CT and MBT for insomnia. We examined which interventions (CT vs. MBT) influence which process measures, and which process measures

influence treatment outcome. Specifically, do CT and MBT have differential effects on cognitive and mindfulness process measures? The present investigation was conducted as part of a randomised controlled trial designed to evaluate the relative efficacy of four sessions of CT and MBT following a four-session CBT-I intervention (Wong et al., 2016). Wong et al. (2016) found that outcomes for a standardised four session CBT-I intervention were equally enhanced by the addition of four sessions of CT or MBT. Both CT and MBT resulted in significant improvements in insomnia severity, total sleep time and sleep efficiency, and the effects were maintained at follow-up. In terms of the mediation analysis, it was expected that CT would bring about more change in cognitive process measures relative to MBT, and that MBT would bring about more change in mindfulness process measures relative to CT. It was further expected that greater change in both cognitive and mindfulness process measures would be predictive of greater improvement across treatment.

## Method

### *Study Design*

This study employed a randomised mixed-group design. Participants were individually administered four sessions of CBT for insomnia. They were then randomly allocated to immediate or delayed conditions (see Figure 1). This was to allow determination of whether the second treatment improved outcomes beyond the passage of time alone. In the second treatment phase, participants were also randomly allocated to either four sessions of CT for insomnia (Harvey et al., 2007) or four sessions comprising components of MBT (Segal, Williams, & Teasdale, 2002). Follow-up was at 3 months post-CT/MBT. Further details of the design are described in Wong et al. (2016).

### *Participants*


The sample was recruited from GPs, psychologists, local pharmacies, and advertisements in the local press. Potential participants contacted the researcher to express interest to participate, and underwent telephone screening. A total of 57 participants completed the four sessions of CBT-I and were randomised to receive either CT or MBT. See Figure 1 for participant flow.

### *Selection Criteria*

The inclusion criteria were: (1) 18–65 years of age; (2) complaint of difficulty initiating or maintaining sleep for at least 6 months; (3) mean sleep onset latency or minutes awake after sleep for longer than 30 minutes for at least 3 nights per week; (4) interference in daytime functioning; (5) total sleep time on sleep diary under 6.5 hours for more than 3 nights out of every 7; (6) taking no medication to assist sleep, or be taking a stable dose of medication.

Individuals were excluded if: (1) they presented clear evidence of other sleep disorders; (2) they were receiving treatment of insomnia from another mental health professional; (3) they presented with sleep disturbances due to a medical and/or physical condition as assessed by clinical interview; (4) they had sleep disturbances due to the physiological effect of a substance; (5) the insomnia was assessed as purely secondary to another psychological disorder. Details of the numbers excluded due to these criteria are presented in Figure 1.

### **Behaviour Change**



**FIGURE 1**  
Participants flow.

76

*Primary Outcome Measure: Insomnia Severity*

**Insomnia Severity Index (ISI; Bastien, Vallières, & Morin, 2001).** The ISI is a five-item scale that assesses severity of both nighttime and daytime symptoms of insomnia over the period of the past week. The ISI has adequate internal consistency (Cronbach’s alpha = 0.74– 0.78) with evidence supporting concurrent, predictive, and content validity (Bastien et al., 2001). The total score of the ISI was used to indicate overall severity of insomnia, with higher scores indicating greater severity.

### *CT Process Measures*

The CT process measures were selected to map onto the maintaining factors in the Cognitive Model of Insomnia (Harvey, 2002).

**The Anxiety and Preoccupation about Sleep Questionnaire (APSQ; Tang & Harvey, 2004).** The APSQ is a 10-item scale that measures sleep-related anxiety and preoccupation associated with insomnia. The participants rate on a 10-point scale how true each statement was over the past 2 days (1 = *not true*, 10 = *very true*). The items are then summed to obtain a total score that ranges from 10 to 100. Higher scores indicate higher level of anxiety and preoccupation about sleep. The APSQ has high internal consistency (Cronbach's alpha = 0.92).

**Sleep Associated Monitoring Index (SAMI; Semler & Harvey, 2004).** The SAMI is a valid and reliable instrument that can index monitoring before and after treatment for sleep disturbance. It has 30 items with five subscales assessing presleep monitoring, including: (a) body sensations consistent with falling asleep, (b) body sensations inconsistent with falling asleep, (c) the environment, (d) the clock, and (e) calculation of time. The psychometric properties are adequate (Semler & Harvey, 2004). Internal consistency of the scale is high (Cronbach's alpha = 0.91) and so is that for the subscales (Cronbach's alpha = 0.77–0.87) (Semler & Harvey, 2004).

**Dysfunctional Beliefs and Attitudes about Sleep Scale (DBAS; Morin & Espie, 2003).** The DBAS is a 30-item, self-report scale. It was designed to assess misconceptions about the effects of insomnia, the unpredictability of sleep, daily sleep requirements, causes of insomnia and sleep-promoting habits. The average of all DBAS items serves as the total DBAS score. This instrument has been found to have highly acceptable internal consistency (Cronbach's alpha = 0.86; Edinger & Sampson, 2002).

**Sleep-Related Behaviours Questionnaire (SRBQ; Ree & Harvey, 2004).** The SRBQ is a 32-item, self-report scale designed to assess the use of safety behaviours employed to promote sleep and cope with tiredness. It showed adequate internal consistency (0.92), and discriminates normal sleepers from those with insomnia (Ree & Harvey, 2004).

### *Mindfulness Process Measures*

The following instruments were designed to measure key processes that MBCT was designed to alter.

**The Kentucky Inventory of Mindfulness Skills (KIMS; Baer, Smith, & Allen, 2004).** The KIMS is a 39-item self-report measure (Baer et al., 2004) used for assessing mindfulness skills. The KIMS has adequate to good internal consistency (Cronbach's alpha = 0.83–0.91) and validity (Baer et al., 2004). The subscales of the KIMS are based on an empirically supported factor structure (Baer, Smith, Hopkins, Krietemeyer, & Toney, 2006). The KIMS measures four facets of mindfulness separately and reliably; namely, observing, describing, acting with awareness, and accepting (or allowing) without judgment (Baer et al., 2006). The total score of the KIMS is used to reflect mindfulness skills. Higher scores reflect more mindfulness skills.

**Non-reactivity to inner experience (NRI).** According to Baer and colleagues (2006), factor analyses of a combined pool of items from mindfulness questionnaires showed that there are five clear interpretable facets of mindfulness. Four of the facets are

### **Behaviour Change**

included in the KIMS. Combined items of the Mindfulness Questionnaire (MQ; Chadwick, Hember, Mead, Lilley, & Dagnan, 2005) and Freiburg Mindfulness Inventory (FMI; Buchheld, Grossman, & Walach, 2001) describe a fifth facet of mindfulness, a non-reactive stance toward internal experience, that the KIMS does not include. To measure this other facet of mindfulness, that is, NRI, four items from the MQ and three items from the FMI were administered (Baer et al., 2006). The FMI is a valid and reliable questionnaire that assesses non-judgmental, present-moment observation and openness to negative experience. It has an internal consistency of Cronbach's  $\alpha = 0.93$  (Walach, Buchheld, Büttenmüller, Kleinknecht, & Schmidt, 2006). The MQ assesses a mindful approach to distressing thoughts and images with a good internal consistency (Cronbach's  $\alpha = 0.89$ ; Chadwick et al., 2005).

### *Procedure*

**Pretreatment assessment.** After completion of a phone screen, eligible participants attended a face-to-face clinical interview. The face-to-face clinical interview included questions adapted from the Sleep History Assessment (Morin & Espie, 2003) and an interview guide for assessment of insomnia (Ong, Shapiro, & Manber, 2008). Eligible participants were then asked to complete pencil-and-paper pretreatment questionnaires described above. After completion of the preintervention measures, participants waited 4 weeks without treatment. Participants were then invited to complete the outcome measures described above before commencing CBT treatment.

**Therapist.** The third author, a registered psychologist, delivered all interventions to the participants. Regular clinical supervision by independent experts, both clinical psychologists, was received in CBT, CT, and mindfulness. The psychologist had specific training in CBT, CT, and MBT.

**Cognitive behaviour therapy for insomnia.** All participants received four 1-hour intervention sessions following a standard CBT for primary insomnia protocol (Carney & Edinger, 2008). The majority of this intervention was behaviourally oriented, with one session focused on cognitive strategies, including constructive worry and cognitive restructuring worksheets.

**Cognitive therapy.** The four CT sessions were comprised of CT components as described in Harvey (2005) and Harvey et al. (2007). Individual insomnia formulations, based on the cognitive model, were derived to conceptualise each participant's cognitive maintaining factors (Harvey, 2006). Cognitive restructuring and tailored behavioural experiments (Perlis, Aloia, & Kuhn, 2011; Ree & Harvey, 2004) were developed to assist in reversing the cognitive maintaining factors: unhelpful beliefs about sleep (Morin et al., 2002), worry about sleep (Tang & Harvey, 2004), monitoring for sleep-related threats, and misperception of sleep (Harvey & Tang, 2012).

**Mindfulness-based therapy.** The four MBT sessions focused on the development of mindfulness skills with session content based on the approach described by Segal et al. (2002). In-session meditations were included during each session, and participants were encouraged to listen to mindfulness meditation recordings (e.g., sitting meditation) 6 days out of 7 at times of the day that suited them. Additionally, participants were assigned informal mindfulness tasks each week (e.g., mindful eating and any other activity whereby attention is focused on one activity over a period of time).



The 3-Minute Breathing Space was also introduced for bringing formal meditation practice into daily life. Participants were encouraged to carry out the practice of 3-Minute Breathing Space three times daily at set times and also at any other times when they felt stressed. Barriers and benefits of practice were discussed at each session. A recurrent theme in the MBT sessions was the awareness of present moment experience without judgment.

## Results

### *Demographics of Participants at Pretreatment*

The following analyses were performed on data from the 57 participants who received either CT ( $n = 26$ ) or MBT ( $n = 31$ ). There were no dropouts in either condition. Both treatments had mainly female participants (MBT: 65.6%; CT: 61.3%). The mean (*SD*) age of participants was 49.96 (12.6) years for MBT and 48.97(13.4) years for CT (no significant difference,  $t = -.287$ ,  $p = .775$ ).

### *Treatment Outcome*

As described in detail in Wong et al. (2016), scores on the primary outcome measure (ISI) reduced during CT/MBT treatment significantly more than during the waitlist period; 8.9 at pretreatment, 5.7 at follow-up:  $F(1, 55) = 7.27$ ,  $p < .01$ ,  $\eta^2 = .12$ . There were no significant differences between posttreatment and follow-up, and no significant differences between treatments.

### *Changes in MBT Process Measures*

This analysis explored whether the MBT process measures were affected by treatment and the degree to which these processes were involved in any improvement (see Table 1 for descriptive statistics). A 3 (Time: pretreatment vs. posttreatment vs. follow-up)  $\times$  2 (Intervention: MBT vs. CT) repeated measures MANOVA was computed to compare scores on the KIMS and the NRI. A significant effect for Time on the MBT measures was found, Wilks lambda = .84,  $F(4, 52) = 2.53$ ,  $p = .05$ ,  $\eta^2 = .16$ , as well as a significant interaction for Time by Intervention, Wilks lambda = .84,  $F(4, 52) = 2.5$ ,  $p = .05$ ,  $\eta^2 = .16$ . Univariate tests showed a difference for KIMS over time but not for NRI (see Table 1). This indicates that MBT had a greater impact on KIMS than did CT.

### *Changes in CT Process Measures*

A 3 (Time: pretreatment vs. posttreatment vs. follow-up)  $\times$  2 (Intervention: MBT vs. CT) repeated measures MANOVA was computed to compare scores on the APSQ, DBAS, SAMI, and SRBQ respectively. There was a significant effect of Time for the process measures, Wilks lambda = .41,  $F(8, 48) = 8.59$ ,  $p < .001$ ,  $\eta^2 = .59$ . Univariate tests showed a significant difference for each CT process measure over the treatment period (Table 2). As can be seen in Table 2, for all the CT process measures, there was a trend toward greater reductions for CT posttreatment compared to MBT. However, the size of these differences were not significant, as evident by the Time by Intervention analysis, Wilks lambda = .87,  $F(8, 48) = .98$ ,  $p = .52$ ,  $\eta^2 = .13$ .

### *Predictors of Sleep Improvement*

We tested whether improvement in insomnia severity would be significantly predicted by treatment-related changes for process measures of MBT and CT. Factor analysis

### **Behaviour Change**

**TABLE 1**  
Means and Standard Deviations for Mindfulness Process Measures Over Time for MBT and CT Conditions

Time Intervention Mindfulness measures	Pretreatment		Posttreatment		Follow-up		$F(2,110)$	$\eta^2$
	CT <sup>a</sup>	MBT <sup>b</sup>	CT <sup>a</sup>	MBT <sup>b</sup>	CT <sup>a</sup>	MBT <sup>b</sup>		
KIMS	128.19 (14.83)	121.31 (13.99)	128.52 (17.42)	132.77 (18.79)	128.00 (14.08)	127.15 (19.19)	5.31*	.09
NRI	49.89 (8.48)	50.13 (8.58)	49.09 (8.91)	51.09 (8.02)	50.36(7.64)	49.57 (9.85)	.01 $ns$	0


Note: \* $p < .01$ . CT = cognitive therapy; MBT = mindfulness-based therapy. KIMS = Kentucky Inventory of Mindfulness. NRI = non-reactivity to inner experiences. <sup>a</sup> $n = 31$ , <sup>b</sup> $n = 26$ .

**TABLE 2**

Means and Standard Deviations for CT Process Measures Over Time for MBT and CT

Time Intervention CT measures <i>M (SD)</i>	Pretreatment		Posttreatment		Follow-up		Univariate <i>F</i> (2,110) for time, $\eta^2$ in brackets**
	CT <sup>a</sup>	MBT <sup>b</sup>	CT <sup>a</sup>	MBT <sup>b</sup>	CT <sup>a</sup>	MBT <sup>b</sup>	
APSQ	34.81 (18.2)	33.58 (18.42)	23.03 (15.71)	26.19 (13.67)	19.52 (9.84)	25.35 (12.84)	22.62 (.29)
DBAS	2.33 (1.38)	3.04 (1.35)	1.47 (.99)	2.27 (1.33)	1.4 (1.18)	2.1 (1.23)	22.28 (.29)
SAMI	70.23 (22.91)	73.88 (23.3)	57.26 (18.47)	64.96 (18.38)	55 (12.23)	66.54 (19.38)	19.69 (.26)
SRBQ	68.03 (16.21)	68.35 (17.86)	59.35 (13.54)	64.88 (17.17)	55.16 (18.38)	59.73 (16.49)	20.55 (.27)

Note: \*\* $p < .001$  in each case. CT = cognitive therapy; MBT = mindfulness based therapy. APSQ = Anxiety and Preoccupation about Sleep Questionnaire. DBAS = Dysfunctional Beliefs and Attitudes about Sleep Scale. SAMI = Sleep Associated Monitoring Index. SRBQ = Sleep Related Behaviours Questionnaire. <sup>a</sup> $n = 31$ , <sup>b</sup> $n = 26$ .



**FIGURE 2**

Predictors of treatment outcomes.

Note: \*\*\* $p < .001$ , \*\* $p < .01$ , \* $p < .05$ .

**TABLE 3**

The Associations Between the Composite Cognitive Process Measure Posttreatment, Pretreatment Process Scores, and Treatment Condition

Variable		B	SE	$\beta$	$r^2$ (p value)
CT composite change posttreatment	CT composite pretreatment	.73	.08	.73	.50 (.001)
	Mindfulness composite pretreatment	-.12	.09	-.12	.01 (.41)
	Treatment type	.32	.16	.16	.03 (.11)

Note: B = unstandardised coefficient B; SE = standard error;  $\beta$  = beta;  $r^2$  = part correlation coefficient square. ( $n = 57$ ) d

was used to derive a single composite score for the MBT process measures and for the CT process measures. These scores were then computed in regression analyses to determine the extent that MBT process measures or CT process measures predict change in insomnia severity. A diagrammatic summary of all the following findings may be found in Figure 2.

First, we assessed whether the composites of the CT process measures and mindfulness measures at pretreatment predicted change in scores on the composite of CT measures at posttreatment. As can be seen in Table 3, only the composite of CT

**TABLE 4**

The Associations Between the Composite Mindfulness Process Measure Post-MBT or CT Treatment, Pretreatment Process Measures, and Type of Treatment Received

Variable		B	SE	$\beta$	$r^2$ ( $p$ value)
Mindfulness composite change posttreatment	CT composite pretreatment	.08	.10	.08	.01 (.62)
	Mindfulness Composite pretreatment	.72	.10	.72	.48 (.001)
Treatment type		.44	.20	.22	.05 (.04)

Note: B = unstandardised coefficient B; SE = standard error;  $\beta$  = beta;  $r^2$  = part correlation coefficient square. ( $N = 57$ ).

process measures at pretreatment significantly predicted the composite CT process measure change at posttreatment. The composite of mindfulness process measures at pretreatment did not significantly predict composite CT at posttreatment. As can be seen in Figure 2, the relationship between intervention (MBT or CT) and the composite CT process measures approached significance ( $p = .06$ ), with the direction being that CT was more likely to result in higher CT process measure change at the end of treatment.

Similarly, we assessed whether the composites of CT process measures and mindfulness measures at pretreatment predicted change in scores on the composite of mindfulness measures at posttreatment. Table 4 presents the process variables predicting change in composite mindfulness process measures at posttreatment.

As can be seen in Table 4, mindfulness process measures pretreatment significantly predict the composite of mindfulness measures at posttreatment. The composite CT measure at pretreatment was not significantly related to the mindfulness measures at posttreatment. However, treatment type was found to significantly predict the composite of mindfulness at posttreatment.

Further analysis was conducted to determine whether the composite CT and mindfulness process measures at pretreatment, composites of CT and mindfulness process measures posttreatment, ISI pretreatment, and intervention type predicted change in scores on the ISI at posttreatment.

As can be seen in Table 5 the only variables significantly related to ISI at posttreatment were pretreatment ISI and the composite CT measure posttreatment. The total  $R^2$  value for the model is .66,  $F(6, 50) = 15.99$ ,  $p < .001$ . This included the unique variance explained by each variable, with any overlap or shared variance removed or partialled out. Also, as can be seen in Table 5, 14% of the variance in insomnia severity posttreatment was uniquely explained by insomnia severity at pretreatment, while 10% of the variance in insomnia severity posttreatment was uniquely explained by the composite of CT process measures posttreatment.

## Discussion

Two major findings arose from the current study. First, as expected, MBT was associated with significant improvement in mindfulness measures compared to CT. This is in contrast to the CT process measures that were significantly improved

### Behaviour Change

**TABLE 5**

Process Variables predicting Insomnia Severity Index Scores Post MBCT or CT

Variable	B	SE	$\beta$	$r^2$
ISI Pretreatment	.40	.09	.46***	.14***
CT composite pretreatment	-.48	.55	-.18	-.01
CT composite posttreatment	2.14	.57	.56***	.10***
Mindfulness composite pretreatment	.42	.47	.11	.01
Mindfulness Composite posttreatment	-.70	.48	-.18	-.01
Interventions	-.12	.71	-.02	.00

Note: B = unstandardised coefficient B; SE = standard error;  $\beta$  = beta;  $r^2$  = part correlation coefficient square.

\*\*\*  $p < .001$ .

by both CT and mindfulness treatment interventions. Second, the CT composite process measure strongly predicted variance in insomnia severity at posttreatment, but contrary to prediction, this result did not differ significantly across CT and MBT. Unlike the CT process measure composite, the mindfulness process measure composite at posttreatment did not significantly predict change in insomnia severity.

The first finding that MBT produced change in cognitive processes is striking given that the version of MBT in the current study did not incorporate any direct work on modifying unhelpful thoughts, beliefs or behaviours. Others have noted that mindfulness-based approaches raise awareness of thoughts and feelings in ways that allow a disengagement from ruminative processes (Nolen-Hoeksema, 1991), seeing them as 'mental events' and not accurate reflections of reality (Segal et al., 2002). The current results are consistent with the notion that mindfulness-based approaches may not only alter metacognitive processes but cognitive processes, such as monitoring, distorted perception and cognitive content. This is in line with research suggesting cognitive change can result from therapies other than CT. For example, an investigation of MBT in the treatment of insomnia suggested that dysfunctional beliefs about sleep and thought-control strategies change with MBT (Larouche et al., 2014). Additionally, in a trial of CBT and pharmacotherapy, Morin and colleagues (2002) found that the degree of belief change during both treatments was predictive of treatment response, but CBT produced more belief change than did pharmacotherapy. Further, Eidelman et al. (2016) found that dysfunctional beliefs are modified not just by CT but also by BT, a treatment that does not explicitly aim to modify cognitive processes such as unhelpful beliefs. Finally, Garland, Rouleau, Campbell, Samuels, and Carlson (2015) implemented a mindfulness-based cancer recovery intervention and found improvements in insomnia and dysfunctional beliefs about sleep. These studies, together with the current results, are consistent with the notion that change in cognitive processes is important *and* that it is achievable via CT or MBT.

The finding that CT process measures changed in each of CT and MBT is particularly noteworthy, given the second finding that the cognitive process measures independently predicted insomnia severity over and above pretreatment insomnia severity scores. Harvey et al. (2017) found a consistent result in finding that a

composite CT process measure mediated treatment outcome in both CT and behaviour therapy (BT) for insomnia. The results of the current study and that of Harvey et al. support the cognitive model of insomnia in providing evidence for the importance of cognitive change in insomnia treatment (Harvey, 2002, 2005). Importantly, these results suggest that CT is not the only method for treating these cognitive maintaining factors. Indeed, BT and MBT appear to also be effective. The finding that cognitive processes predict treatment outcome suggests that (1) understanding how best to alter cognitive processes may assist optimisation of treatment effectiveness, and (2) that treatment research should include other therapies in addition to CT/CBT in the quest to maximise cognitive change.

Moving now to the prediction that the mindfulness process measure would predict treatment effects. Findings were contrary to this prediction in that the mindfulness process measure composite did not relate to treatment outcome. Research by Ong et al. (2009) examined the effects of an integrated CBT/MBT approach to insomnia treatment, finding that although the treatment was effective, it did not bring about changes in the mindfulness process measure, the Kentucky Inventory of Mindfulness Skills (KIMS). This suggests a lack of relationship between the mindfulness process measure and treatment outcome, and in this way is consistent with the current findings. Ong et al. suggested that the exposure to mindfulness was perhaps not enough to sufficiently develop mindfulness skills. In response to this result, Ong and Sholtes (2010), in a single case study, revised their integrated CBT/MBT intervention by extending the number of sessions from 6 to 8, including an all-day meditation retreat. KIMS scores were increased in the case study. Where the current results differ is that the mindfulness process measure score *did* change with a short, four-session MBT intervention. The fact that process measure change was not associated with an improved treatment outcome is puzzling. In essence, an improvement in mindfulness suggests that participants undergoing MBT had improved skills in observing, describing, acting with awareness, and accepting without judgment. However, it appears that unless this also resulted in a change in cognitions about preoccupation with sleep, dysfunctional thoughts about sleep and/or decreased safety behaviours, insomnia was not likely to change. Further research could therefore focus on what facilitated this cognitive change for the MBT participants.

There are several study limitations to note; the first is that the sample size of the current study may not have had sufficient power to detect significant difference between treatments on the cognitive process measures. There was a trend toward CT being associated with a larger difference on the cognitive process measure compared with MBT, but it only approached statistical significance. Second, it may be that the CT process measures were more sensitive to change in the current study than the MBT process measures. The CT measures were developed for use in insomnia, whereas the MBT measures were developed for transdiagnostic use.

Third, although video recordings were reviewed during regular supervision of the therapist by experts in each approach, no formal fidelity treatment checks were carried out. The design of the study could also be improved by the posttreatment measures being collected by someone independent to the delivery of the treatment. However, the fact we collected an objective physiological measure that showed the same response pattern to the self-report measures suggests this was not a major flaw (see Wong et al., 2016).

In summary, there is evidence that both CT and MBT are effective treatments for insomnia. There is also evidence that CT produces change in cognitive process

measures and that MBT brings about change in both mindfulness and cognitive process measures. The current results support that changing the content of unhelpful thoughts/beliefs (as in CT) or the relationship with unhelpful thoughts/beliefs (as in MBT) results in a reduction in unhelpful cognitive processes. Finally, the results of this study underscore the importance of cognitive factors as mediators in insomnia, given that they were significantly related to treatment outcome regardless of treatment condition.

## Compliance with Ethical Standards

This study received no external funding. The project was approved by the Human Research Committee at Murdoch University and deemed to be consistent with the guidelines of the National Statement on Ethical Conduct involving Human Research. Informed consent was obtained from all individual participants included in the study and the information sheet and consent form was approved by the Human Research Committee.

## Conflict of Interest

None.

## References

- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Washington, DC: Author.
- Baer, R.A., Smith, G.T., & Allen, K.B. (2004). Assessment of mindfulness by self-report the Kentucky Inventory of Mindfulness Skills. *Assessment, 11*, 191–206. doi:10.1177/1073191104268029
- Baer, R.A., Smith, G.T., Hopkins, J., Krietemeyer, J., & Toney, L. (2006). Using self-report assessment methods to explore facets of mindfulness. *Assessment, 13*, 27–45. doi:10.1177/1073191105283504
- Bastien, C.H., Vallières, A., & Morin, C.M. (2001). Validation of the Insomnia Severity Index as an outcome measure for insomnia research. *Sleep Medicine, 2*, 297–307. doi:10.1016/S1389-9457(00)00065-4
- Bélanger, L., Harvey, A.G., Fortier-Brochu, É., Beaulieu-Bonneau, S., Eidelman, P., Talbot, L., . . . Morin, C.M. (2016). Impact of comorbid anxiety and depressive disorders on treatment response to cognitive behavior therapy for insomnia. *Journal of Consulting and Clinical Psychology, 84*, 659–667. doi:10.1037/ccp0000084
- Black, D.S., O'Reilly, G.A., Olmstead, R., Breen, E.C., & Irwin, M.R. (2015). Mindfulness meditation and improvement in sleep quality and daytime impairment among older adults with sleep disturbances: A randomized clinical trial. *JAMA Internal Medicine, 175*, 494–501. doi:10.1001/jamainternmed.2014.8081
- Brasure, M., Fuchs, E., MacDonald, R., Nelson, V.A., Koffel, E., Olson, C.M., . . . Kane, R.L. (2016). Psychological and behavioral interventions for managing insomnia disorder: An evidence report for a clinical practice guideline by the American College of Physicians. *Annals of Internal Medicine*. Retrieved from <http://annals.org/article.aspx?articleid=2519966>
- Buchheld, N., Grossman, P., & Walach, H. (2001). Measuring mindfulness in insight meditation (Vipassana) and meditation-based psychotherapy: The development of the Freiburg Mindfulness Inventory (FMI). *Journal for Meditation and Meditation Research, 1*, 11–34.
- Carney, C.E., & Edinger, J.D. (2006). Identifying critical beliefs about sleep in primary insomnia. *Sleep, 29*, 444–453.
- Carney, C.E., & Edinger, J.D. (2008). *Overcoming insomnia: A cognitive-behavioral therapy approach workbook*. Oxford University Press.



- Carney, C.E., & Waters, W.F. (2006). Effects of a structured problem-solving procedure on pre-sleep cognitive arousal in college students with insomnia. *Behavioral Sleep Medicine*, 4, 13–28. doi:10.1207/s15402010bsm0401\_2
- Chadwick, P., Hember, M., Mead, S., Lilley, D., & Dagnan, D. (2005). *Responding mindfully to unpleasant thoughts and images: Reliability and validity of the Mindfulness Questionnaire*. Unpublished manuscript.
- Charles, J., Harrison, C., Britt, H. (2009). Insomnia. *Australian Family Physician*, 38, 283.
- Chiesa, A., & Serretti, A. (2010). A systematic review of neurobiological and clinical features of mindfulness meditations. *Psychological Medicine*, 40, 1239–1252. doi:10.1017/S0033291709991747
- Cvengros, J.A., Crawford, M.R., Manber, R., & Ong, J.C. (2015). The relationship between beliefs about sleep and adherence to behavioral treatment combined with meditation for insomnia. *Behavioral Sleep Medicine*, 13, 52–63. doi:10.1080/15402002.2013.838767
- Daley, M., Morin, C.M., LeBlanc, M., Gregoire, J.P., Savard, J., & Baillargeon, L. (2009). Insomnia and its relationship to health-care utilization, work absenteeism, productivity and accidents. *Sleep Medicine*, 10, 427–438. doi:10.1016/j.sleep.2008.04.005
- Edinger, J., & Sampson, W. (2002). An abbreviated cognitive-behavioral insomnia therapy designed for primary care settings. *Sleep*, 25, A70–A71.
- Eidelman, P., Talbot, L., Ivers, H., Bélanger, L., Morin, C.M., & Harvey, A.G. (2016). Change in dysfunctional beliefs about sleep in behavior therapy, cognitive therapy, and cognitive-behavioral therapy for insomnia. *Behavior Therapy*, 47, 102–115. doi:10.1016/j.beth.2015.10.002
- Ellis, J.G., Gehrman, P., Espie, C.A., Riemann, D., & Perlis, M.L. (2012). Acute insomnia: Current conceptualizations and future directions. *Sleep Medicine Reviews*, 16, 5–14.
- Franzen, P.L., & Buysse, D.J. (2008). Sleep disturbances and depression: risk relationships for subsequent depression and therapeutic implications. *Dialogues in clinical neuroscience*, 10(4), 473.
- Garland, S.N., Carlson, L.E., Stephens, A.J., Antle, M.C., Samuels, C., & Campbell, T.S. (2014). Mindfulness-based stress reduction compared with cognitive behavioral therapy for the treatment of insomnia comorbid with cancer: A randomized, partially blinded, noninferiority trial. *Journal of Clinical Oncology*, 32, 449–457. doi:10.1200/JCO.2012.47.7265
- Garland, S.N., Zhou, E.S., Gonzalez, B.D., & Rodriguez, N. (2016). The quest for mindful sleep: A critical synthesis of the impact of mindfulness-based interventions for insomnia. *Current Sleep Medicine Reports*, 2, 142–151. doi:10.1007/s40675-016-0050-3
- Garland, S.N., Rouleau, C.R., Campbell, T., Samuels, C., & Carlson, L.E. (2015). The comparative impact of mindfulness-based cancer recovery (MBCR) and cognitive behavior therapy for insomnia (CBT-I) on sleep and mindfulness in cancer patients. *Explore: The Journal of Science and Healing*, 11, 445–454. doi:10.1016/j.explore.2015.08.004
- Geiger-Brown, J.M., Rogers, V.E., Liu, W., Ludeman, E.M., Downton, K.D., & Diaz-Abad, M. (2015). Cognitive behavioral therapy in persons with comorbid insomnia: A meta-analysis. *Sleep Medicine Reviews*, 23, 54–67.
- Hall, M., Thayer, J.F., Germain, A., Moul, D., Vasko, R., Puhl, M., . . . Buysse, D.J. (2007). Psychological stress is associated with heightened physiological arousal during NREM sleep in primary insomnia. *Behavioral Sleep Medicine*, 5, 178–193. doi:10.1080/15402000701263221
- Harvey, A.G. (2002). A cognitive model of insomnia. *Behaviour Research and Therapy*, 40, 869–893. doi:10.1016/S0005-7967(01)00061-4
- Harvey, A.G. (2005). A cognitive theory and therapy for chronic insomnia. *Journal of Cognitive Psychotherapy*, 19, 41–59. doi:10.1891/jcop.19.1.41.66332
- Harvey, A.G. (2006). What about patients who can't sleep? Case formulation for insomnia. In N. Tarrrier, (Ed.), *Case formulation in cognitive behavior therapy: The treatment of challenging and complex cases* (pp. 293–311). New York, NY: Routledge/Taylor & Francis Group.
- Harvey, A.G., Bélanger, L., Talbot, L., Eidelman, P., Beaulieu-Bonneau, S., Fortier-Brochu, É., . . . Morin, C.M. (2014). Comparative efficacy of behavior therapy, cognitive therapy, and cognitive

- behavior therapy for chronic insomnia: A randomized controlled trial. *Journal of Consulting and Clinical Psychology*, 82, 670–683. doi:10.1037/a0036606
- Harvey, A.G., Dong, L., Bélanger, L., & Morin, C.M. (2017). Mediators and treatment matching in behavior therapy, cognitive therapy and cognitive behavior therapy for chronic insomnia. *Journal of Consulting and Clinical Psychology*, 85, 975–987. doi:10.1037/ccp0000244
- Harvey, A.G., Sharpley, A.L., Ree, M.J., Stinson, K., & Clark, D.M. (2007). An open trial of cognitive therapy for chronic insomnia. *Behaviour Research and Therapy*, 45, 2491–2501. doi:10.1016/j.brat.2007.04.007
- Harvey, A.G., Soehner, A.M., Kaplan, K.A., Hein, K., Lee, J., Kanady, J., ... Buysse, D.J. (2015). Treating insomnia improves mood state, sleep, and functioning in bipolar disorder: A pilot randomized controlled trial. *Journal of Consulting and Clinical Psychology*, 83, 564–577. doi:10.1037/a0038655
- Harvey, A.G., & Tang, N.K. (2012). (Mis) perception of sleep in insomnia: A puzzle and a resolution. *Psychological Bulletin*, 138, 77–101. doi:10.1037/a0025730
- Hiller, R.M., Johnston, A., Dohnt, H., Lovato, N., & Gradisar, M. (2015). Assessing cognitive processes related to insomnia: A review and measurement guide for Harvey's cognitive model for the maintenance of insomnia. *Sleep Medicine Reviews*, 23, 46–53. doi:10.1016/j.smrv.2014.11.006
- Jacobs, G.D. (1998). *Say good night to insomnia: The 6-week solution*. New York, NY: Henry Holt.
- Larouche, M., Cote, G., Belisle, D., & Lorrain, D. (2014). Kind attention and non-judgment in mindfulness-based cognitive therapy applied to the treatment of insomnia: state of knowledge. *Pathologie Biologie*, 62, 284–291. doi:10.1016/j.patbio.2014.07.002
- Morin, C.M., & Benca, R. (2012). Chronic insomnia. *The Lancet*, 379, 1129–1141. doi:10.1016/S0140-6736(11)60750-2
- Morin, C.M., Blais, F., & Savard, J. (2002). Are changes in beliefs and attitudes about sleep related to sleep improvements in the treatment of insomnia? *Behaviour Research and Therapy*, 40, 741–752. doi:10.1016/S0005-7967(01)00055-9
- Morin, C.M., & Espie, C.A. (2003). *Insomnia: A clinical guide to assessment and treatment*. New York, NY: Kluwer Academic/Plenum Publishers.
- Morin, C.M., Vallières, A., Guay, B., Ivers, H., Savard, J., Mérette, C., ... Baillargeon, L. (2009). Cognitive behavioral therapy, singly and combined with medication, for persistent insomnia: A randomized controlled trial. *Journal of the American Medical Association*, 301, 2005–2015. doi:10.1001/jama.2009.682
- Nakamura, Y., Lipschitz, D.L., Kuhn, R., Kinney, A.Y., & Donaldson, G.W. (2013). Investigating efficacy of two brief mind–body intervention programs for managing sleep disturbance in cancer survivors: A pilot randomized controlled trial. *Journal of Cancer Survivorship*, 7, 165–182. doi:10.1007/s11764-012-0252-8
- Nolen-Hoeksema, S. (1991). Responses to depression and their effects on the duration of depressive episodes. *Journal of Abnormal Psychology*, 100, 569–582. doi:10.1037/0021-843X.100.4.569
- Norell-Clarke, A., Jansson-Fröjmark, M., Tillfors, M., Harvey, A.G., & Linton, S.J. (2014). Cognitive processes and their association with persistence and remission of insomnia: Findings from a longitudinal study in the general population. *Behaviour Research and Therapy*, 54, 38–48. doi:10.1016/j.brat.2014.01.002
- Okajima, I., Komada, Y., & Inoue, Y. (2011). A meta-analysis on the treatment effectiveness of cognitive behavioral therapy for primary insomnia. *Sleep and Biological Rhythms*, 9, 24–34. doi:10.1111/j.1479-8425.2010.00481.x
- Ong, J.C., Manber, R., Segal, Z., Xia, Y., Shapiro, S., & Wyatt, J.K. (2014). A randomized controlled trial of mindfulness meditation for chronic insomnia. *Sleep*, 37, 1553–1563. doi:10.5665/sleep.4010
- Ong, J.C., Shapiro, S.L., & Manber, R. (2008). Combining mindfulness meditation with cognitive-behavior therapy for insomnia: A treatment-development study. *Behavior Therapy*, 39, 171–182. doi:10.1016/j.beth.2007.07.002

- Ong, J.C., Shapiro, S.L., & Manber, R. (2009). Mindfulness meditation and cognitive behavioral therapy for insomnia: A naturalistic 12-month follow-up. *Explore: The Journal of Science and Healing*, 5, 30–36. doi:10.1016/j.explore.2008.10.004
- Ong, J.C., Ulmer, C.S., & Manber, R. (2012). Improving sleep with mindfulness and acceptance: A metacognitive model of insomnia. *Behaviour Research and Therapy*, 50, 651–660. doi:10.1016/j.brat.2012.08.001
- Ong, J., & Sholtes, D. (2010). A mindfulness-based approach to the treatment of insomnia. *Journal of Clinical Psychology*, 66, 1175–1184. doi:10.1002/jclp.20736
- Perlis, M., Aloia, M., & Kuhn, B. (Eds.) (2011). *Behavioral treatments for sleep disorders*. San Diego, CA: Academic Press.
- Perlis, M.L., Giles, D.E., Buysse, D.J., Tu, X., & Kupfer, D.J. (1997). Self-reported sleep disturbance as a prodromal symptom in recurrent depression. *Journal of Affective Disorders*, 42, 209–212.
- Qaseem, A., Kansagara, D., Forcica, M.A., Cooke, M., & Denberg, T.D. (2016). Management of chronic insomnia disorder in adults: A clinical practice guideline from the American College of Physicians. *Annals of Internal Medicine*. Retrieved from <http://annals.org/article.aspx?articleid=2518955>
- Ree, M.J., & Harvey, A.G. (2004). Investigating safety behaviours in insomnia: The development of the Sleep-Related Behaviours Questionnaire (SRBQ). *Behaviour Change*, 21, 26–36. doi:10.1375/bech.21.1.26.35971
- Ree, M., Junge, M., & Cunnington, D. (2017). Australasian Sleep Association position statement regarding the use of psychological/behavioral treatments in the management of insomnia in adults. *Sleep Medicine*, 36, S43–S47.
- Riemann, D., & Perlis, M.L. (2009). The treatments of chronic insomnia: A review of benzodiazepine receptor agonists and psychological and behavioral therapies. *Sleep Medicine Reviews*, 13, 205–214. doi:10.1016/j.smrv.2008.06.001
- Roth, T. (2007). Insomnia: Definition, prevalence, etiology, and consequences. *Journal of Clinical Sleep Medicine*, 3, S7.
- Segal, Z.V., Williams, J.M.G., & Teasdale, J.D. (2002). *Mindfulness-based cognitive therapy for depression: A new approach to preventing relapse*. New York, NY: Guilford Press.
- Semler, C.N., & Harvey, A.G. (2004). Monitoring for sleep-related threat: A pilot study of the Sleep Associated Monitoring Index (SAMI). *Psychosomatic Medicine*, 66, 242–250.
- Shapiro, S.L., Carlson, L.E., Astin, J.A., & Freedman, B. (2006). Mechanisms of mindfulness. *Journal of Clinical Psychology*, 62, 373–386. doi:10.1002/jclp.20237
- Singareddy, R., Vgontzas, A.N., Fernandez-Mendoza, J., Liao, D., Calhoun, S., Shaffer, M.L., & Bixler, E.O. (2012). Risk factors for incident chronic insomnia: A general population prospective study. *Sleep Medicine*, 13, 346–353.
- Sivertsen, B., Krokstad, S., Mykletun, A., & Øverland, S. (2009). Insomnia symptoms and use of health care services and medications: The HUNT-2 study. *Behavioral Sleep Medicine*, 7, 210–222. doi:10.1080/15402000903190199
- Sivertsen, B., Omvik, S., Pallesen, S. ale, Bjorvatn, B., Havik, O.E., Kvale, G., . . . Nordhus, I.H. (2006). Cognitive behavioral therapy vs zopiclone for treatment of chronic primary insomnia in older adults: A randomized controlled trial. *Journal of the American Medical Association*, 295, 2851–2858. doi:10.1001/jama.295.24.2851
- Sunnhed, R., & Jansson-Fröjmark, M. (2014). Are changes in worry associated with treatment response in cognitive behavioral therapy for insomnia? *Cognitive Behaviour Therapy*, 43, 1–11. doi:10.1080/16506073.2013.846399
- Tang, N.K., & Harvey, A.G. (2004). Correcting distorted perception of sleep in insomnia: A novel behavioural experiment? *Behaviour Research and Therapy*, 42, 27–39. doi:10.1016/S0005-7967(03)00068-8
- Trauer, J.M., Qian, M.Y., Doyle, J.S., Rajaratnam, S.M., & Cunnington, D. (2015). Cognitive behavioral therapy for chronic insomnia: A systematic review and meta-analysis. *Annals of Internal Medicine*, 163, 191–204. doi:10.7326/M14-2841

- Walach, H., Buchheld, N., Buttenmüller, V., Kleinknecht, N., & Schmidt, S. (2006). Measuring mindfulness — The Freiburg Mindfulness Inventory (FMI). *Personality and Individual Differences*, 40, 1543–1555. doi:[10.1016/j.paid.2005.11.025](https://doi.org/10.1016/j.paid.2005.11.025)
- Wong, M.Y., Ree, M.J. and Lee, C.W. (2016). Enhancing CBT for chronic insomnia: A randomised clinical trial of additive components of mindfulness or cognitive therapy. *Clinical Psychology & Psychotherapy*, 23. 377–385. doi:[10.1002/cpp.1980](https://doi.org/10.1002/cpp.1980)