

NECESSARY AND SUFFICIENT CONDITIONS  
FOR OSCILLATIONS OF DELAY DIFFERENTIAL INEQUALITIES

JURANG YAN

A necessary and sufficient condition is obtained for a first order linear delay differential inequality to be oscillatory. Our main result improves and extends some known results.

1. INTRODUCTION

The oscillation theory of first order linear delay differential inequalities and equations has been extensively developed by many authors. We mention in particular the papers by Koplatadze and Chanturia [1], Ladas [2], and Ladas and Stavroulakis [3]. for a survey of results concerned with oscillations we refer to Zhang [5] and the references therein. Most of the known results, however, give sufficient conditions for oscillations.

The aim of this paper is to obtain necessary and sufficient conditions under which the first order delay differential inequality

$$(1) \quad x(t)[x'(t) + p(t)x(t - \tau(t))] \leq 0$$

is oscillatory, where:

- (a)  $p(t) \in C[[t_0, \infty), \mathbf{R}]$  and  $p(t) \geq 0$ ;
- (b)  $\tau(t) \in C[[t_0, \infty), \mathbf{R}]$ ,  $\tau(t) \geq 0$  and  $\lim_{t \rightarrow \infty} [t - \tau(t)] = \infty$ .

Conditions (a) and (b) are assumed to hold throughout this paper.

As is customary, a solution is said to be *oscillatory* if it has arbitrarily large zeros and *nonoscillatory* if it is eventually positive or eventually negative. Inequality (1) is said to be *oscillatory* if every solution of (1) is oscillatory.

Our main result is the following:

**THEOREM.** *A necessary and sufficient condition for inequality (1) to be oscillatory is that for every sufficiently large  $T$  there exists a  $\tilde{t} > T$  such that*

$$(2) \quad \int_{\tilde{t}-\tau(\tilde{t})}^{\tilde{t}} p(s) ds > \frac{1}{e}.$$

---

Received 15 April, 1988

Copyright Clearance Centre, Inc. Serial-fee code: 0004-9729/89 \$A2.00+0.00.

As an immediate consequence of the theorem we obtain the following corollary for the inequalities

$$(3) \quad x'(t) + p(t)x(t - \tau(t)) \leq 0,$$

$$(4) \quad x'(t) + p(t)x(t - \tau(t)) \geq 0$$

and the equation

$$(5) \quad x'(t) + p(t)x(t - \tau(t)) = 0.$$

**COROLLARY.** *A necessary and sufficient condition for (3) to have no eventually positive solution, (4) to have no eventually negative solution and (5) to be oscillatory, is that condition (2) is satisfied.*

### 2. PROOF OF THE THEOREM

Consider the integral inequality

$$(6) \quad u(t) \geq p(t) \exp \left( \int_{t-\tau(t)}^t u(s) ds \right).$$

The function  $u(t)$  said to be a nonnegative solution on  $[t_1, \infty)$  ( $t_1 \geq t_0$ ) if  $u(t) \geq 0$  is continuous and bounded on  $[t_1, t_2)$ , continuous on  $[t_2, \infty)$  and satisfies (6) for all  $t \geq t_2$ , where  $t_2 = \inf\{t \mid t - \tau(t) \geq t_1\}$ .

Using an argument similar to that given by the author ([4]), the following lemma is easily established:

**LEMMA 1.** *Inequality (1) has a nonoscillatory solution if and only if inequality (6) has a nonnegative solution on  $[t_1, \infty)$  for some large  $t_1$ .*

For some  $t_1 \geq t_0$ , we define the following sequence of functions on  $[t_1, \infty)$ .

$$(7) \quad \begin{aligned} u_0(t) &= 0, & t &\geq t_2 \\ u_k(t) &= \begin{cases} p(t) \exp \left( \int_{t-\tau(t)}^t u_{k-1}(s) ds \right), & t \geq t_2, \\ 0, & t_1 \leq t < t_2 \end{cases} \\ & k = 1, 2, \dots \end{aligned}$$

where  $t_2 = \inf\{t \mid t - \tau(t) \geq t_1\}$ . By a simple induction, we have that

$$(8) \quad 0 \leq u_k(t) \leq u_{k+1}(t), \quad t \geq t_1, \quad k = 1, 2, \dots$$

LEMMA 2. Inequality (6) has a nonnegative solution on  $[t_1, \infty)$  if and only if the sequence (7) is convergent for any  $t \geq t_1$ .

PROOF: Suppose that  $u(t)$  is a nonnegative solution on  $[t_1, \infty)$ . Thus  $u_0(t) \leq u(t)$ ,  $t \in [t_1, \infty)$ . Then, it follows by induction that

$$u_0(t) \leq u_k(t) \leq u(t), \quad t \geq t_1, \quad k = 1, 2, \dots$$

Hence, from (8), we see that the sequence (7) converges to a finite limit  $\bar{u}(t)$ ,  $t \geq t_1$ .

Conversely, suppose that the sequence (7) converges to a finite limit  $u(t)$  for any  $t \geq t_1$ . Hence  $u_k(t) \leq u(t)$ ,  $t \geq t_1$ , and, by the monotone convergence theorem,

$$\lim_{k \rightarrow \infty} \int_{t-\tau(t)}^t u_k(s) ds = \int_{t-\tau(t)}^t u(s) ds, \quad t \geq t_2.$$

From (7), we have that

$$u(t) = \begin{cases} p(t) \exp \left( \int_{t-\tau(t)}^t u(s) ds \right), & t \geq t_2, \\ 0 & t_1 \leq t < t_2, \end{cases}$$

which implies that (6) has a nonnegative solution on  $[t_1, \infty)$ . ■

By combining Lemma 1 and Lemma 2 we have the following result:

LEMMA 3. Inequality (1) is oscillatory if and only if, for every large  $T$ , there exists a  $\bar{t} > T$  such that

$$\lim_{k \rightarrow \infty} u_k(\bar{t}) = \infty.$$

PROOF OF THEOREM: Let

$$(9) \quad \int_{\bar{t}-\tau(\bar{t})}^{\bar{t}} p(s) ds = \bar{\alpha}, \quad \bar{t} > T \geq t_3,$$

where  $t_3 = \inf\{t \mid t - \tau(t) \geq t_2\}$ . From (9), we are able to choose a constant  $\alpha$  with  $\frac{1}{e} < \alpha < \bar{\alpha}$ . Now, choose an interval  $\Delta = (\bar{t} - \delta, \bar{t} + \delta)$  ( $\bar{t} - \delta \geq T$ ) such that

$$(10) \quad \int_{t-\tau(t)}^t p(s) ds \geq \alpha \quad t \in \Delta.$$

Consider the sequence (7) on the interval  $\Delta$ ,

$$u_0(t) = 0, \quad t \in \Delta,$$

$$u_1(t) = \xi_1 p(t) \quad t \in \Delta, \text{ where } \xi_1 = 1,$$

$$\begin{aligned} u_2(t) &= p(t) \exp \left( \int_{t-\tau(t)}^t p(s) ds \right) \geq p(t) e^{\xi_1 \alpha} \\ &= \xi_2 p(t), \text{ where } \xi_2 = e^{\xi_1 \alpha}. \end{aligned}$$

By a simple induction, it is easy to show that

$$(11) \quad u_{k+1}(t) \geq \xi_{k+1}p(t), \quad t \in \Delta$$

where

$$(12) \quad \xi_{k+1} = e^{\xi_k \alpha}.$$

Obviously,  $\{\xi_n\}_{n=1}^\infty$  is increasing. Let  $\lim_{n \rightarrow \infty} \xi_n = \xi$ , where  $\xi$  is a positive constant or infinite. We claim that  $\xi$  is infinite. Indeed, if this were false, the sequence would converge to some finite limit  $\xi > 0$ . Then, in view of (12), it follows that  $\xi = e^{\alpha \xi}$  and hence we derive that  $(\ln \xi)/\xi = \alpha$ . But  $\max_{\xi > 0} \{(\ln \xi)/\xi\} = \frac{1}{e}$ . This implies  $\alpha \leq \frac{1}{e}$ , which is impossible. Therefore  $\lim_{n \rightarrow \infty} \xi_n = \infty$ .

From (11), we find that

$$\lim_{k \rightarrow \infty} u_k(t) = \infty, \quad t \in \Delta.$$

Thus, by Lemma 3, inequality (1) is oscillatory.

Conversely, suppose that there exists a large  $T \geq t_3$ , where  $t_3 = \inf\{t \mid t - \tau(t) \geq t_2\}$  such that

$$(13) \quad \int_{t-\tau(t)}^t p(s)ds \leq \frac{1}{e}, \quad t \geq T.$$

Consider the sequence (7) on  $[T, \infty)$ ,

$$\begin{aligned} u_0(t) &= 0, & t \geq T, \\ u_1(t) &= p(t) \leq ep(t), & t \geq T. \end{aligned}$$

From (7) and (13), we have that

$$u_2(t) = p(t) \exp\left(\int_{t-\tau(t)}^t ep(s)ds\right) \leq ep(t), \quad t \geq T.$$

By induction, we have that

$$u_k(t) \leq ep(t), \quad t \geq T, \quad k = 1, 2, \dots$$

Consequently,  $u_n(t)$  converges pointwise to a finite limit for each  $t \geq T$ . By Lemma 3, (1) has a nonoscillatory solution. The proof of the Theorem is complete. ■

## 3. REMARKS

1. The Theorem in this paper improves and extends the previous results of Koplatdze and Chanturia [1] and Ladas [2].

2. One easily generalises the results in this paper to inequalities of the form

$$x(t)[x(t) + a(t)x(t) + p(t)x(t - \tau(t))] \leq 0,$$

where  $p(t)$  and  $\tau(t)$  satisfy conditions (a) and (b). The function  $a(t) \in C[[t_0, \infty), \mathbb{R})$  and is allowed to change sign on  $[t_0, \infty)$ . Hence our results improve and extend the results of Ladas and Stravroulakis, [3].

3. Our techniques can be applied to advanced differential inequalities of the form

$$x(t)[x'(t) + a(t)x(t) - p(t)x(t + \tau(t))] \leq 0,$$

where  $p(t)$ ,  $\tau(t)$  and  $a(t)$  satisfy the above conditions.

## REFERENCES

- [1] R.G. Koplatadze and T.A. Chanturia, 'On oscillatory and monotonic solutions of first-order differential equations with retarded arguments', Russian, *Differentsial'nye Uravneniya* **8** (1982), 1463—1465.
- [2] G. Ladas, 'Sharp conditions for oscillations caused by delays', *App. Anal.* **9** (1979), 93–98.
- [3] G. Ladas and I.P. Stavroulakis, 'On delay differential inequalities of first order', *Funkcial Ekvac.* **25** (1982), 105–113.
- [4] J. Yan, 'Oscillation of solutions of first order delay differential equations', *Nonlinear Anal.* **11** (1987), 1279–1287.
- [5] B.G. Zhang, 'A survey of the oscillation of solutions of first order differential equations with deviating argument', *Proc. International Conference on Trends in Nonlinear Analysis, Arlington, TX June 12–22* (1984).

Department of Mathematics,  
Shanzi University,  
Taiyuan, Shanxi,  
People's Republic of China.