

Summer Meeting, 28 June–1 July 2010, Nutrition and health: cell to community

Who performs better? An objective assessment of consumers ability to perform food label-based, nutrition information assessments

D. Mackison¹, A. S. Anderson² and W. L. Wrieden³

¹Department of Nursing and Midwifery, University of Stirling, Stirling FK9 4LA, UK, ²Centre for Public Health Nutrition Research, University of Dundee, Dundee DD1 9SY, UK and ³School of Pharmacy and Life Sciences, Robert Gordon University, Aberdeen AB25 IHG, UK

In order to use nutrition label information to guide healthy food choices, consumers require a basic understanding of nutrition communications. Reviews of the literature highlight gaps in understanding (e.g. desirable and less desirable levels of intake) and competence (ability to identify differences in nutrient content)^(1,2). The current study objectively examines the consumers' ability to assess nutrition label information.

A short postal survey (which underwent reliability, validity, item discrimination and item difficulty assessment)⁽³⁾ was distributed to a nation-wide sample representative of gender, age and socio-economic (SES) position⁽⁴⁾ of the UK population. The survey included a 10-item measurement tool on ability to perform food label-based, nutrition information assessments. The questions comprised numerical (*n* 3) and comprehension tasks (*n* 7) with a possible performance score of 0 to 10. Kruskal Wallis and Mann–Whitney *U* tests were used to test for differences between overall performance score and demographic variables (gender, age, educational attainment and SES). Completed responses for 786 UK adults (35.5% male and 64.5% female) aged 18 upwards and representative of SES for the UK population were obtained.

Overall, the mean performance score was 6.7 (SD±2.3) and 77% obtained >5. No significant difference was detected by gender. Respondents <50 years were significantly more likely to obtain a high score than those aged ≥50 years (*P*<0.05). Kruskal Wallis tests indicated that score increased with higher levels of educational attainment (*P*<0.001) and SES (e.g. more affluent) (*P*<0.001).

Question focus	Question type	% of respondents answering each question correctly						
		Overall sample (<i>n</i> 786)	SES			Educational attainment		
			Affluent (<i>n</i> 294)	Comfortable (<i>n</i> 193)	Non-affluent (<i>n</i> 288)	Post school (<i>n</i> 448)	School (<i>n</i> 152)	None (<i>n</i> 179)
Sugar in 2 servings	N*	60.9	76.2	66.3	41.0	73.2	59.2	31.8
Fat in half the pack	N*	70.1	80.3	74.1	55.9	80.8	68.4	45.3
Servings in the product	N*	78.3	86.7	81.9	66.3	85.9	78.3	59.2
Fat content in product 1	C†	63.0	66.3	61.7	58.7	65.2	66.4	53.1
Fat content in product 2	C†	41.8	45.9	44.0	36.5	46.0	36.2	37.4
Sugar content product 3	C†	68.8	75.5	66.8	62.5	75.7	69.7	52.5
Comparison of sat fat	C†	87.0	91.2	92.7	78.1	90.8	90.8	75.4
Lowest sat fat content	C†	25.8	29.9	26.4	21.2	30.1	24.3	16.2
Energy content	C†	85.6	89.5	90.2	78.1	89.3	90.8	72.1
Reducing salt intake	C†	87.5	91.2	91.7	80.2	92.6	86.8	76.0

*N = numerical question; †C = comprehension question.

In conclusion, current nutrition information presentation is unlikely to provide optimal assistance for consumers from more disadvantaged backgrounds and further work is needed to enhance existing nutrition communication for vulnerable groups.

Funding provided by the Food Standards Agency Postgraduate Scholarship Scheme.

- Cowburn G & Stockley L (2005) Consumer understanding and use of nutrition labelling: a systematic review. *Public Health Nutr* 8, 21–28.
- Grunert KJ & Wills JM (2007) A review of European research on consumer response to nutrition information on food labels. *J Public Health* 15, 385–399.
- Mackison D, Wrieden WL & Anderson AS (2010) Validity and reliability testing of a short questionnaire developed to assess consumers' use, understanding and perception of food labels. *Eur J Clin Nutr* 64, 210–217.
- CACI (2006) Acorn the Smarter Consumer Classification User Guide. <http://www.caci.co.uk/brochures.aspx>