

a-T-MENABILITY OF GROUPS ACTING ON TREES

ŚWIATOSŁAW R. GAL

We present some partial results concerning a-T-menability of groups acting on trees.
Various known results are given uniform proofs.

1. DEFINITION OF A-T-MENABILITY

DEFINITION 1.1: Given a metric space (X, ρ) , the action $G \rightarrow \text{Isom}(X)$ is *metrically proper* if, given $x \in X$, the displacement function $G \ni g \mapsto \rho(x, gx) \in \mathbb{R}$ is proper. If the group action is set we say that X is metrically proper.

The above property does not depend on a choice of a point x .

DEFINITION 1.2: (M. Gromov) A locally compact, second countable, compactly generated group G is *a-T-menable* if there exists metrically proper isometric G -action on some affine Hilbert space.

REMARK. a-T-menability is equivalent to the existence of C_0 -approximate unity consisting of positive definite functions. The latter is called in the literature the Approximation Property of Haagerup.

Throughout this paper by representation we mean isometric affine action. By a subgroup we always mean a closed subgroup.

2. MOTIVATION

We are motivated by the following Theorems:

THEOREM 2.1. ([2, Theorem 6.2.8]) *Let Γ be a countable group acting on a tree without inversions, with finite edge stabilisers. If vertex stabilisers in Γ are a-T-menable, then so is Γ .*

THEOREM 2.2. ([2, Example 6.1.6]) *If N is a-T-menable and G/N is amenable then G is a-T-menable.*

On the other hand:

Received 30th September, 2003

Partially supported by a KBN grant 2 P03A 017 25. The author would like to thank Tadeusz Januszkiewicz for his helpful information, Agnieszka for her hospitality in Vienna and finally Jan Dymara for carefully reading preliminary version of this paper and his assistance in improving the presentation.

Copyright Clearance Centre, Inc. Serial-fee code: 0004-9727/04 \$A2.00+0.00.

THEOREM 2.3. ([5, Chapter 8 Lemma 6]) *Every isometric affine representation of $G = SL_2\mathbb{Z} \times \mathbb{Z}^2$ has a \mathbb{Z}^2 -fixed point.*

The group G acts on a tree as it can be decomposed as $(\mathbb{Z}/4\mathbb{Z}^2) *_{\mathbb{Z}/2\mathbb{Z}^2} (\mathbb{Z}/6\mathbb{Z}^2)$. The factors are a-T-menable by Theorem 2.2, since \mathbb{Z}^2 is a-T-menable as it acts on \mathbb{R}^2 . On the other hand $SL_2\mathbb{Z}$ is a-T-menable by the result of Haagerup [4]. Therefore extra assumptions about finiteness of edge stabilisers (in Theorem 2.1) or amenability of the quotients (in Theorem 2.2) cannot be simply weakened to a-T-menability.

We believe that a-T-menability is a property of representations (see next section for definitions), rather than that of groups. Therefore in this paper we shall concentrate on a question: what are the conditions under which given proper affine representations of two groups extend to one of their product with amalgamation, rather than whether there exist some such representation that extend.

The most naïve observation is that if there is a proper representation of a free product with amalgamation, the restrictions are proper representations of the factors that coincide on the common subgroup. It is not known whether the reverse holds, however we shall prove some results in this direction.

The main result of this paper is Theorem 5.4. As a result we strengthen a result from [2] (see Section 6). Constructions, we give, are purely geometric. We shall also give an affirmative answer to the question of A. Valette, whether Baumslag-Solitar groups are a-T-menable. This was originally done (using another approach) in [3].

3. ACTIONS ON TREES IN GENERAL

Before we examine the case of a free product with amalgamation, let us restate a general observation of Haagerup [4] and its easy generalisations.

Let T be a tree. By E we denote a vector space of functions on edges of T with finite support. By V we denote the affine space of functions on vertices of T with finite support and total mass one. The structure of affine space is given as follows: $\delta_v - \delta_w$ is equal to the characteristic function of the segment joining vertices v and w (with the appropriate signs with respect to some auxiliary orientation on the edges of T).

DEFINITION 3.1: Let $U(T)$ be an affine Hilbert space completion of V defined above.

There is an obvious $\text{Aut}(T)$ action of the group of cellular automorphisms of T on $U(T)$, with an $\text{Aut}(T)$ -equivariant isometric embedding of T . An immediate consequence of the construction is

PROPOSITION 3.2: If Γ acts metrically properly on a tree T then Γ is a-T-menable.

In particular $SL_2\mathbb{Z} = \mathbb{Z}/4 *_{\mathbb{Z}/2} \mathbb{Z}/6$ is a-T-menable.

If T is locally finite, the (topological) group of all cellular automorphisms of T is a-T-menable. Even if Γ acts effectively on T , the inclusion $\Gamma \rightarrow \text{Aut}(T)$ in general is not closed (therefore we cannot conclude that Γ is a-T-menable). However we have

PROPOSITION 3.3. If Γ acts on a locally finite tree T and there exists an affine representation of Γ on an affine Hilbert space W , such that stabiliser of any vertex acts properly, then $U(T) \oplus W$ is Γ -proper.

PROOF: Given $g_n \in \Gamma$ with bounded displacement (as acting on T), for any vertex v the distance from g_nv to v is bounded. Since there are only finitely many such vertices, one can take a subsequence such that g_nv is constant. Stabilisers of vertices act properly on W , thus $\{g_n\}$ is relatively compact. \square

PROPOSITION 3.4. If Γ acts on a locally finite tree T and for each vertex v some affine proper representation of Stab_v on W_v extends to a (perhaps non-proper) affine representation of Γ , then Γ is a-T-menable.

PROOF: This is obvious if the quotient $\Gamma \backslash T$ has finitely many vertices, since then the sum of the representations corresponding to any lifts will fulfill the assumptions of Proposition 3.3.

Since there is no infinite sum operation in the category of affine Hilbert spaces, the construction will depend on choices made.

Let $\{v_k\}$ be a sequence of representatives of the vertices of the quotients. Let K_k be an exhausting sequence of compact subsets of Γ , let $x_k \in W_{v_k}$,

$$a_k = k^2 + \sup \{ \|x_k - gx_k\|^2 : g \in K_k \}.$$

Define a norm on $\prod_k W_{v_k}$ in a following way:

$$\|y - z\|^2 := \sum a_k^{-2} \|y_k - z_k\|^2.$$

Let W be the completion of the affine space

$$\left\{ y \in \prod_k W_{v_k} : \|y - x\| < \infty \right\}.$$

The choices are made in a such way that Γ acts diagonally on W and there are Γ -equivariant projections (up to scalar change of norm) to each of W_{v_k} . \square

NOTE. The proof of Proposition 3.4 follows the standard proof of the fact that direct limit of a-T-menable groups is also a-T-menable.

The example of $Sl_2 \mathbb{Z} \times \mathbb{Z}^2$ shows that there are some obstructions for a representation to extend from the subgroup. In the terms of group cohomology, inclusion of groups need not induce epimorphisms on the level of the first cohomology.

4. AFFINE REPRESENTATIONS AND SUBGROUPS

If V is G -invariant subspace of W then W/V is a linear G -representation (the coset V is a fixed point), therefore V is metrically proper if and only if W is. In general, there

is no minimal invariant subspace. Let $H < G$. Assume $V \subset W$ is H -invariant subspace. Fix $x \in W$. Define

$$\psi_{(V \subset W)}: H \backslash G \ni Hg \mapsto \|gx + V\| \in \mathbb{R}$$

(where $gx + V$ is a coset of gx in W/V). If $\psi_{(V \subset W)}$ is proper we say that $(V \subset W)$ is $H \backslash G$ -proper. The definition does not depend on the choice of x . If $H = \{e\} < G$, and V is any point, then $(V \subset W)$ is $H \backslash G$ -proper exactly if W is G -proper.

LEMMA 4.1. *Let $G_1 < G_2 < G_3$. If there are $W_1 \subset W_2 \subset W_3$, such that W_i is G_i invariant and $(W_i \subset W_{i+1})$ is $G_i \backslash G_{i+1}$ -proper then $(W_1 \subset W_3)$ is $G_1 \backslash G_3$ -proper.*

PROOF: If $\psi_{(W_1 \subset W_3)}(G_1 g_n)$ is bounded, then $\psi_{(W_2 \subset W_3)}(G_2 g_n)$ is bounded. Therefore $g_n = g'_n h_n$ where $g'_n \in G_2$ and $\{h_n\}$ is relatively compact. By the triangle inequality

$$\|g'_n x - x\| \leq \|g_n(h_n^{-1}x - x)\| + \|g_n x - x\| + \|h_n^{-1}x - x\|,$$

so $\psi_{(W_1 \subset W_2)}(G_1 g'_n)$ is bounded, therefore $g'_n = g''_n h'_n$, where $g''_n \in G_1$ and $\{h'_n\}$ is relatively compact. Finally $g_n = g''_n(h'_n h_n)$. \square

Unfortunately it is not known whether for any $H < G$ and proper G -representation W there exist H -invariant subspace V , such that $(V \subset W)$ is $H \backslash G$ -proper. The cases when it does happen are discussed in the following sections.

5. FREE PRODUCTS WITH AMALGAMATION

If $\Gamma = G_1 *_{\mathcal{H}} G_2$ then, according to Serre theory [6], a graph T with the set of vertices equal to $\Gamma/G_1 \cup \Gamma/G_2$ and the set of edges equal to Γ/\mathcal{H} , with inclusion as incidence relation, is a tree (on which Γ acts on the left). The representations of G_1 and G_2 on $U(T)$ have global fixed points.

Let H be a common subgroup in G_1 and G_2 . Let W_i be G_i -representations. Let W be their common H -invariant subspace. Define $\mathcal{H}_i := W_i/W$. Inductively decompose $\uparrow_H^{G_i} \mathcal{H}_{\omega}$ (where ω is a sequence of 1s and 2s) with respect to H as $\mathcal{H}_{\omega} \oplus \mathcal{H}_{i\omega}$. G_1 acts on

$$\mathcal{H}_2^{\bullet} = (\mathcal{H}_2 \oplus \mathcal{H}_{12}) \oplus (\mathcal{H}_{212} \oplus \mathcal{H}_{1212}) \oplus \dots$$

and

$$\mathcal{H}_1^{\circ} = (\mathcal{H}_{21} \oplus \mathcal{H}_{121}) \oplus \dots,$$

G_2 acts on $\mathcal{H}_1^{\bullet} = (\mathcal{H}_1 \oplus \mathcal{H}_{21}) \oplus \dots$ and $\mathcal{H}_2^{\circ} = (\mathcal{H}_{12} \oplus \mathcal{H}_{212}) \oplus \dots$. Both representations of H on \mathcal{H}_i^{\bullet} coincide.

DEFINITION 5.1. Let

$$W_{\Gamma} = W \oplus \mathcal{H}_1^{\bullet} \oplus \mathcal{H}_2^{\bullet} = W_1 \oplus \mathcal{H}_1^{\circ} \oplus \mathcal{H}_2^{\bullet} = W_2 \oplus \mathcal{H}_1^{\bullet} \oplus \mathcal{H}_2^{\circ}.$$

An immediate consequence from the construction is

THEOREM 5.2. Let $H < G_i$, $i = 1, 2$. Let W_i be G_i -representations. Let W be their common H -invariant subspace. Let $\Gamma = G_1 *_{\mathbb{H}} G_2$. Then W_i and W are respectively G_i - and H -invariant subspaces of W_{Γ} .

NOTE. Although there is no way to induce an affine representation, W_{Γ} is morally equal to $\uparrow_{G_1}^{\Gamma} W_1 \oplus \uparrow_{G_2}^{\Gamma} W_2 / \uparrow_H^{\Gamma} W$. If $W' \subset W$ is another H -invariant subspace, then W_{Γ} is not a Γ -invariant subspace of W' (constructed from the triple $W_1 \supset W' \subset W_2$).

A straightforward consequence of Proposition 3.3 and Theorem 5.2 is the following.

COROLLARY 5.3. If H is of finite index in G_1 and G_2 and if there are metrically proper representations of G_i that coincide when restricted to H , then $G_1 *_{\mathbb{H}} G_2$ is a-T-menable.

EXAMPLE. ([2, 1].) The torus group $\Gamma_{p,q} = \langle x, y | x^p = y^q \rangle$ is a-T-menable.

THEOREM 5.4. With the assumptions of Theorem 5.2, if W_i is proper affine G_i -representation and $(W \subset W_i)$ is $H \backslash G_i$ -proper for $i = 1, 2$ then

- (1) $W_{\Gamma} \oplus U(T)$ is Γ -proper,
- (2) if $H' < G_1$ and V is proper H' -space such that $(V \subset W_1)$ is $H' \backslash G_1$ -proper, then $(V \subset W_{\Gamma} \oplus U(T))$ is $H' \backslash \Gamma$ -proper.

PROOF: In fact (1) is a special case of (2). Therefore we shall prove (2).

Given $\gamma_n \in \Gamma$ such that $\psi_{(W,V)}(\gamma_n)$ is bounded. Define the length function $\ell: \Gamma \rightarrow \mathbb{N}$ by $\ell|_H \equiv 0$, $\ell(\gamma) = \min\{\ell(\eta) + 1 | \gamma = \eta g, \text{ where } g \in G_1 \cup G_2\}$. This function is equal to the distortion of the action on $U(T)$. Therefore we may find a subsequence such that $\ell(\gamma_n) = k$.

If $k \leq 1$ there is nothing to prove. If all but finitely many $\gamma_n \in G_2$ we have to use Lemma 4.1.

Let $x \in V$. Define $\varphi_x(\gamma)$ to be the component of γx in $\bigoplus_{|\omega|=l(\gamma)} \mathcal{H}_{\omega}$.

LEMMA 5.5. If $\gamma = \gamma_1 \gamma_2$, $\ell(\gamma) = \ell(\gamma_1) + \ell(\gamma_2)$ and $\ell(\gamma_2) \geq 1$, then $\|\varphi_x(\gamma)\| = \|\varphi_x(\gamma_2)\|$.

PROOF: Without loss of generality $\ell(\gamma_1) = 1$. From the definition of induced representation $\gamma_1 \mathcal{H}_{\omega} \perp \mathcal{H}_{\omega}$, therefore $\varphi_x(\gamma) = \gamma_1 \varphi_x(\gamma_2)$. \square

Now we proceed by induction on k as follows. We define η_n such that $\gamma_n = \eta_n g_n$ ($g_n \in G_i$) and $l(\eta_n) = k - 1$.

From Lemma 5.5 we see $\|\gamma_n x - x\| \geq \|\varphi_x(\gamma_n)\| = \|\varphi_x(g_n)\| = \psi_{(W_i,W)}(g_n)$. Therefore $\{g_n\}$ is relatively compact. Since $\eta_n x = \gamma_n x - \eta_n(g_n x - x)$ and, by induction assumption, $\psi_{(V,W_{\Gamma})}$ is proper when restricted to cosets of elements of length smaller than k , we obtain the claim. \square

6. GROUPS THAT ACT ON TREES WITH FINITE EDGE STABILISERS

The first case, where it is easy to fulfill the assumptions of the Theorem 5.4 is when H is finite, since then one can find a fixed point of any H -representation simply taking the centre of mass $*$ of any orbit. The pair $(\{*\} \subset W_i)$ is $H \backslash G_i$ proper if and only if W_i is proper G_i -space. Summarising this:

PROPOSITION. ([2, Proposition 6.2.3 (1)]) *Let G_1, G_2 be two groups containing finite subgroup H , and let $\Gamma = G_1 *_H G_2$ be the corresponding amalgamated product. If G_1 and G_2 are a-T-menable, then so is Γ .*

Theorem 2.1 is an easy consequence of the above [2].

7. BAUMSLAG-SOLITAR GROUPS AND THEIR CERTAIN GENERALISATIONS

The second easy case occurs when H is of finite index in G_i 's. Then any pair is automatically $H \backslash G_i$ proper.


Let us recall some definitions from [3]. Let $G \subset \mathfrak{N}$ be a closed subgroup of a locally compact, compactly generated, topological group \mathfrak{N} . Let $i_k: H \rightarrow G$ ($k = 1, 2$) be two inclusions onto finite index open subgroups, which are conjugated by an automorphism ϕ of \mathfrak{N} .

DEFINITION 7.1: The \mathfrak{N} -BS group Γ is the group derived from (G, H, i_1, i_2) by the (topological) HNN construction.

THEOREM 7.2. ([3].) *If \mathfrak{N} is a-T-menable then \mathfrak{N} -BS groups are a-T-menable.*


PROOF: We mimic the proof [2, 6.2.7] for the case of a HNN extension, where the edge stabiliser is finite. \square

STEP 1. Let Γ_0 be a fundamental group of the following tree:



We have to find consistent representations of different copies of G . The Hilbert space in each case will be the one on which \mathfrak{N} acts properly. The k -th copy acts by $\phi^k(\cdot)$, where ϕ is the automorphism of \mathfrak{N} that conjugates i_1 and i_2 .

By induction, each of



satisfies assumptions of Theorem 5.4 (alternatively: by Theorem 5.2 and induction we construct representation of Γ_0 and then use Proposition 3.4). It is easy to show [2,

Proposition 6.1.1] that an increasing union of open a-T-menable subgroups is again a-T-menable. Therefore Γ_0 is a-T-menable.

STEP 2. $\Gamma = \Gamma_0 \ltimes \mathbb{Z}$, where \mathbb{Z} acts through the shift. Therefore Γ is an extension of an a-T-menable group with amenable quotient, so Γ is a-T-menable by Theorem 2.2. \square

REFERENCES

- [1] C. Béguin and T. Ceccherini-Silberstein, ‘Formes faibles de moyennabilité pour les groupes à un relateur’, *Bull. Belg. Math. Soc. Simon Stevin* **1** (2000), 135–148.
- [2] P-A. Cherix, M. Cowling, P. Jolissaint, P. Julg and A. Valette, *Groups with the Haagerup property (Gromov’s a-T-menability)* (Birkhäuser Verlag, Basel, 2001).
- [3] S.R. Gal and T. Januszakiewicz, ‘New a-T-menable HNN-extensions’, *J. Lie Theory* **13** (2003), 383–385.
- [4] U. Haagerup, ‘An example of non-nuclear C^* -algebra which has the metric approximation property’, *Invent. Math.* **50** (1979), 279–293.
- [5] P. de la Harpe and A. Valette, ‘La propriété (T) de Kazhdan pour les groupes localement compacts’, *Astérisque* **157** (1989), 158.
- [6] J.P. Serre, *Trees* (Springer-Verlag, Berlin, New York, 1980).

Department of Mathematics
Wrocław University
pl. Grunwaldzki 2/4
50-384 Wrocław
Poland
<http://www.math.uni.wroc.pl/~sgal/>