

VLBA Observations of 3C 120

J. L. Gómez¹, A. P. Marscher², A. Alberdi¹, J. M^a. Martí³ & J. M^a. Ibáñez³

Abstract. We present 1.3 cm and 7 mm VLBA observations of the radio galaxy 3C 120 at epochs November 11 and December 22, 1996. The 7 mm maps, with linear resolution of ~ 0.1 pc, show a very rich structure consisting of up to eight superluminal ($\sim 7c$) components.

The nearby ($z=0.033$) radio galaxy 3C 120 shows a core and gently curving jet with structural changes (5 GHz observations by Walker, Benson, & Unwin 1987). We performed 22 and 43 GHz simultaneous polarimetry VLBA observations of 3C 120 (see Fig. 1 for epochs November 11 and December 22, 1996). With only about one month separation between the two epochs, a significant variation in the source structure is clearly visible. The linear resolution at 7 mm is ~ 0.1 pc, one of the highest achieved to date. We distinguish up to eight different moving components (labeled *A* to *H*) in the maps at 7 mm, with proper motions of the order of 3 mas/yr (except components *F* and *G*), giving superluminal motions of about 7 *c* ($H_0=65$ km s⁻¹ Mpc⁻¹).

Comparing the most recently ejected component *H* with the rest of the moving features, and assuming a similar flux density evolution for them, we deduce a rapidly decreasing emission in the first stages of the evolution of the components, later reaching a slower decay. Another possibility is that component *H* represents a stronger event in the source activity. Components *F* and *G* move significantly slower than the rest of components, indicating a different nature. Following the predictions of our numerical simulations (Gómez et al. 1997; see also these Proceedings, p. 49), *F* and *G* may be the result of the interaction of a strong component (*E*) with recollimation shocks in the underlying jet. In that case *F* and *G* should progressively fade and decelerate. Depending on the relative strength of the interacting shocks, the dragged components might even show a brief backward motion before finally fading (see the animated simulations at <http://www.iaa.es/~jlgomez/jets.html>). This is most probably observed in the inner portion of the jet, and may be revealed by upcoming CMVA 3 mm observations. The polarimetry information should also reveal more information on the nature of the multiple components. However a careful flux and proper motion monitoring program is essential for a full interpretation.

Acknowledgments. This research is supported in part by the Spanish DGICYT (PB94-1275, PB94-0973) and by NATO grant SA.5-2-05(CRG.961 228). J. L. Gómez and J. M^a. Martí gratefully acknowledge return grants from the Spanish Ministry of Education. The National Radio Astronomy Observatory is a facility of the National Science Foundation, operated under a cooperative agreement by Associated Universities, Inc.

References

- Gómez, J. L., et al. 1997. *ApJ*, **482**, L33–36.
Walker, R. C., Benson, J. M., & Unwin, S. C. 1987. *ApJ*, **316**, 546–572.

¹Instituto de Astrofísica de Andalucía (CSIC), Granada, Spain

²Department of Astronomy, Boston University, Boston, USA

³Departamento de Astronomía y Astrofísica, Universidad de Valencia, Valencia, Spain




Figure 1. Total intensity radio maps of 3C 120 at epochs November 11, 1996 (top images) and December 22, 1996. Left column maps correspond to 7 mm wavelength observation, whereas right column maps are for 1.3 cm.