

Use of virulence determinants and seropathotypes to distinguish high- and low-risk *Escherichia coli* O157 and non-O157 isolates from Europe

M. F. ANJUM^{1*}, E. JONES¹, V. MORRISON¹, R. TOZZOLI², S. MORABITO²,
I. TOTH³, B. NAGY³, G. SMITH⁴, A. ASPAN⁵, E. M. NIELSEN⁶, P. FACH⁷,
S. HERRERA-LEÓN⁸, M. J. WOODWARD^{1,9} AND R. M. LA RAGIONE^{1,10}

¹ Department of Bacteriology and Food Safety, Animal Health and Veterinary Laboratories Agency, Weybridge, Addlestone, Surrey, UK; ² Department of Veterinary Public Health and Food Safety, Istituto Superiore di Sanità, Viale Regina Elena Rome, Italy; ³ Veterinary Medical Research Institute of the Hungarian Academy of Sciences, Budapest, Hungary; ⁴ Health Protection Agency, London UK; ⁵ SVA, National Veterinary Institute, Uppsala, Sweden; ⁶ Statens Serum Institut, Department of Microbiological Surveillance and Research, Copenhagen, Denmark; ⁷ ANSES, French Agency for Food, Environmental and Occupational Health & Safety, Food Safety Laboratory, Maisons-Alfort, France; ⁸ Institute of Health Carlos III, National Center of Microbiology, Laboratory of Enterobacteriaceae, Campylobacter and Vibrio, Majadahonda, Madrid, Spain; ⁹ Department of Food and Nutritional sciences, University of Reading, Whiteknights, Reading, UK; ¹⁰ Faculty of Health and Medical Sciences, University of Surrey, Guildford, Surrey, UK

Received 7 August 2012; Final revision 1 May 2013; Accepted 18 June 2013;
first published online 18 July 2013

SUMMARY

The presence of 10 virulence genes was examined using polymerase chain reaction (PCR) in 365 European O157 and non-O157 *Escherichia coli* isolates associated with verotoxin production. Strain-specific PCR data were analysed using hierarchical clustering. The resulting dendrogram clearly separated O157 from non-O157 strains. The former clustered typical high-risk seropathotype (SPT) A strains from all regions, including Sweden and Spain, which were homogenous by Cramer's *V* statistic, and strains with less typical O157 features mostly from Hungary. The non-O157 strains divided into a high-risk SPTB harbouring O26, O111 and O103 strains, a group pathogenic to pigs, and a group with few virulence genes other than for verotoxin. The data demonstrate SPT designation and selected PCR separated verotoxigenic *E. coli* of high and low risk to humans; although more virulence genes or pulsed-field gel electrophoresis will need to be included to separate high-risk strains further for epidemiological tracing.

Key words: *Escherichia coli*, Europe, verotoxigenic, virulence.

INTRODUCTION

Escherichia coli are a normal constituent of the microflora of the gastrointestinal tract. However, certain *E. coli* types have been associated with disease in humans and animals. Based on their clinical

features, virulence and adherence properties, *E. coli* strains associated with diarrhoeal disease have been subdivided into six different categories or pathotypes [1]. Verotoxigenic *E. coli* (VTEC) belong to one of the six pathotypes and are responsible for a large number of outbreaks of human diarrhoea ranging in severity from non-bloody diarrhoea to haemorrhagic colitis (HC) and haemolytic uraemic syndrome (HUS), a leading cause of acute renal failure in children [2, 3].

* Author for correspondence: Dr M. F. Anjum, Animal Health and Veterinary Laboratories Agency, Weybridge, Addlestone, Surrey KT15 3NB, UK.
(Email: Muna.Anjum@ahvla.gsi.gov.uk)

VTEC O157:H7 is the most common serotype associated with HUS and HC, but it is far from being the only serotype associated with this disease [3]. Serotypes such as O26:H11, O111:H8, O145:H28 and O103:H2 are among non-O157 serotypes frequently associated with HUS-related illness in humans [1], and are currently among the most prevalent VTEC serogroups in Europe [4]. In fact, over 200 *E. coli* [5] serotypes associated with verocytotoxin production have been isolated from humans. It has been estimated in the USA that 265 000 VTEC infections occur each year, of which only 36% are caused by O157 strains, with the remainder being caused by non-O157 VTEC. However, the numbers are likely to be higher due to the challenge in identifying non-O157 VTEC strains (<http://www.cdc.gov/ecoli/general/index.html>).

Although the main feature of any VTEC strain is the production of verocytotoxins, which are important in pathogenesis, a repertoire of virulence determinants has been shown to be associated with disease [1–3]. A number of studies have used a polymerase chain reaction (PCR)-based binary typing method, whereby the presence of a gene is scored as 1 and absence as 0, to provide a genetic fingerprint of virulence genes present and a potential risk assessment of these strains to humans [6–8]. Others have used information on the ability of particular VTEC serotypes to produce epidemics and/or HUS disease in humans, in combination with the presence of virulence genes, to designate seropathotypes (SPTs) A–E; with strains from SPTA having most severe sequelae of infection for humans while strains from SPTE are usually confined to animals [9]. In this study utilizing PCR we assessed the presence of 10 virulence genes for 365 *E. coli* strains belonging to O157 and non-O157 VTEC serogroups gathered from 11 European countries. The rationale for choosing these genes was based on their function (Table 1), which has been mainly associated with human illness or severity of clinical infections. Although the criteria for choosing these genes were not based on animal disease, virulence profile of VTECs isolated from diarrhoeic calves and other livestock have shown an association between the presence of virulence factors such as verocytotoxin, intimin and enterohaemolysin and disease [10–12]. Therefore, the aim of this work was to use the PCR data and designated SPT to provide a potential molecular risk assessment of the individual strains and their ability to cause disease and/or epidemics.

METHODS

Bacterial strains

A total of 365 strains were characterized. Most were chosen randomly from common serotypes representative of O157 and non-O157 serogroups associated with verotoxin production isolated from a variety of hosts from Denmark, France, Italy, Spain, and the UK; the O157 and non-O157 isolates from Sweden and Hungary, were representative of their prevalence in their respective countries (see Supplementary Appendix 1, available online). The number of isolates from each country were as follows: Austria (2), Croatia (2), Denmark (32), France (15), Germany (11), Hungary (50), Italy (92), Spain (19), Sweden (61) and the UK (England, Scotland, Wales, 81). There were 162 strains from serogroup O157; while the remaining 203 strains were from a number of non-O157 serogroups previously shown to be VTEC [13–19] (Supplementary Appendix 1) but did not include the O104 serotype as this study commenced before the German outbreak [20]. The strains were chosen prior to any knowledge of their virulence profile by PCR.

Growth, PCR amplification of virulence genes and analysis of data

Strains were cultured at 37 °C, aerobically in Luria–Bertani (LB) broth for 16 h. Following culture, 1 ml cell culture was centrifuged at 16 000 *g* and the pellet resuspended in 300 μ l TE buffer [10 mM Tris (pH 8), 0.1 mM EDTA]. For cell lysis the suspension was heated for 10 min at 95–100 °C, cooled immediately and centrifuged at 16 000 *g*. The supernatant was collected and 5 μ l used as a DNA template for PCR assays. The 10 genes included in this study are listed in Table 1; all PCRs were performed as simplex assays. Presence of the full *efal* gene was determined by two separate PCRs (*efal*-5' and *efal*-3').

PCR reactions were performed in a total volume of 25 μ l containing 5 μ l cell supernatant, 0.4 μ M of each primer (MWG Biotech, USA), 200 μ M of each dNTP, 2.5 μ l \times 10 PCR reaction buffer, 1.5 mM MgCl₂, 0.75 U platinum *Taq* DNA polymerase (Invitrogen, USA). All 365 *E. coli* strains were screened with a panel of 11 PCR primer pairs representing 10 virulence genes selected to represent virulence characteristics associated with VTEC (Table 1). Results of the PCR performed for each strain are listed in Supplementary Appendix 1 and are binary coded.

Table 1. Sequence of the PCR primers and function of genes used in this study for virulotyping the *E. coli* strains

Target	Amplicon size (bp)	Forward primer (5'–3')	Reverse primer (5'–3')	Gene function	Ref.
<i>efa1-5'</i>	1205	TAT GAG ACT GCC AGA GAA AGT	GAT GGG TTG TTG TTG TAT TTC TTC	Adherence of host intestinal mucosa and in repression of host lymphocyte activation response	[21]
<i>efa1-3'</i>	713	TGC GCA CAA TTG ACT ACA GAG GAA	ATA CGA CCA TCA GGG GAA TCA C	As above	[21]
<i>adfO</i>	501	TGG TGG CCC GCA TAC AGC	TGC CCA GTC AGC CCA GGT TA	Intestinal colonization factor	[22]
<i>sodCF</i>	467	GCA GAA CAG GAA GTC CCA A	TTA TTG AAT GAT GCC GCA GG	Prophage-encoded superoxide dismutase	[23]
<i>eae</i>	862	CCC GAA TTC GGC ACA AGC ATA AGC	CCC GGA TCC GTC TCG CCA GTA TTC G	Intimin production to bind to host cells	[24]
<i>vt1</i>	285	CCC GGA TCC ATG AAA AAA ACA TTA TTA ATA GC	CCC GAA TCC AGC TAT TCT GAG TCA ACG	Verocytotoxin production	[25]
<i>vt2</i>	260	CCC GGA TCC ATG AAG AAG ATG TTT ATG GCG	CCC GAA TTC TCA GTC ATT ATT AAA CTG CAC	Verocytotoxin production	[25]
<i>ehx</i>	890	GAG CGA GCT AAG CAG CTT G	CCT GCT CCA GAA TAA ACC ACA	Plasmid encoded enterohaemolysin	[25]
<i>tccp</i>	1150	ATG ATT AAC AAT GTT TCT TCA CTT	TCA CGA GCG CTT AGA TGT ATT AAT GCC	Tir-cytoskeleton coupling protein, an effector protein	[26]
<i>ureA</i>	271	GAC TCC AAG AGA AAA AGA CAA ACT A	CAG ATT ATC GGA TTA TGG ACG GTA	Involved in hydrolysis of urea	[27]
<i>terB</i>	286	AGG CCG TGA CGA ACT GAC C	TCG CAA CGG CAA TAC CAACAC G	Involved in tellurite resistance	[28]

To find associations between strains, the binary-converted PCR data for all genes for the 365 strains were analysed in Bionumerics version 5.10 (Applied Maths, Belgium), whereby agglomerative hierarchical clustering was applied to the Euclidean distances between strains. The unweighted pair-group method with arithmetic mean (UPGMA) was used to construct the dendrogram. Strains were assigned to SPTs using previously published criteria [9]; SPTA were O157:H7 or O157:H⁻ serotypes which have caused epidemic HUS disease in humans; SPTB were non-O157 of multiple serotypes which have caused epidemic HUS disease in humans; SPTC were non-O157 of multiple serotypes which have caused sporadic HUS disease in humans; SPTD were non-O157 of multiple serotypes which have caused sporadic non-complicated diarrhoea in humans; SPTE were non-O157 multiple serotypes not associated with disease in humans.

Statistical analysis

Cramer's *V* statistic was used to determine strong association between each pair of genes/PCR for all 11 PCR products for serotypes O157 and O26. The two nominal variables were the genes/PCR products (Table 2) and each was a binary variable, i.e. had two levels, being present or absent. A Cramer's *V* statistic value of ≥ 0.7 was chosen for a gene/PCR pair from within a country considered to be strongly associated. Such pairs were tabulated (Table 2) to compare virulence characteristics of strains for these serotypes.

RESULTS

Differences in virulence genes present in O157 and non-O157 strains

In the present study, the presence of 10 virulence genes (Table 1) were chosen from those known to be involved in pathogenesis in VTEC, particularly O157:H7, encoded on integrated lamboid prophages, O-islands and plasmids [29], for screening the panel of *E. coli* strains. The resulting binary virulence typing or virulotyping data was analysed by hierarchical clustering (Fig. 1) and each of the main clusters was designated a SPT to give insight to the risk posed by these strains.

Cluster E: the main O157 group

Most O157 strains were present in cluster E, which is a subcluster of cluster A of the dendrogram (Fig. 1).

Table 2. Associations between genes in *E. coli* serotypes O157 and O26 from different European countries calculated by Cramer's *V* association matrix. Only associations >0.7 are shown for simplicity*

Gene pair		O157		O26
Nominal variable 1	Nominal variable 2	Italy (N=51)†	Hungary (N=27)	Italy (N=17)
<i>eae</i>	<i>sodC</i>		0.9035	
<i>eae</i>	<i>icf</i>		0.9035	
<i>eae</i>	<i>ehly</i>		0.8238	
<i>eae</i>	<i>efa1-3'</i>			1.000
<i>eae</i>	<i>ureA</i>			1.000
<i>eae</i>	<i>terB</i>			1.000
<i>vt1</i>	<i>vt2</i>		0.9177	-0.8771
<i>adfO</i>	<i>sodC</i>		0.8071	
<i>adfO</i>	<i>ehly</i>		0.9117	
<i>ureA</i>	<i>terB</i>	1.000		1.000
<i>ureA</i>	<i>tccP</i>	1.000		
<i>efa1-3'</i>	<i>ureA</i>			1.000
<i>efa1-3'</i>	<i>terB</i>			1.000
<i>sodC</i>	<i>ehly</i>		0.7266	
<i>tccP</i>	<i>terB</i>	1.000		

* These may be positive or negative associations.

† Numbers in parentheses indicates total number of isolates from each country.

The virulence profile of the O157 strains were not entirely homogenous, but between 78% and 94% showed presence of: *eae*, *vt2*, *efa1-5'*, *adfO*, *sodCF*, *ehx*, *tccP*, *ureA* and *terB* genes; only 2% of strains were *efa1-3'*⁺, indicating the *efa1* gene was mostly truncated; *vt1* was present in 40% of strains, either singly or in addition to *vt2*. In fact, this was the dominant virulence profile of strains within the O157 SPTA group in subcluster E with $>90\%$ of strains harbouring the previously mentioned virulotype (Table 3). This group included strains of bovine and human origin of serotype O157:H7 or H⁻ strains (where the H complex data was available). Nineteen of the 33 human strains were associated with disease. The O157 strains included all Swedish, most Spanish, Italian and UK strains but only five Hungarian strains from this study; one Croatian and one Austrian strain were also present (Supplementary Appendix 2). However, four of the 117 strains within this cluster were non-O157 (i.e. O176:H⁻, O5, O111, O145); but showed typical O157 features such as presence of the *tccP* gene in the former three strains, while the latter strain showed presence of the truncated *efa1-5'* gene. These strains had been isolated from bovine, foodstuff and human strains.




Fig. 1. The virulotype fingerprint for each strain and the resulting seropathotype (SPT) of the major groups identified from clustering of the binary-converted PCR data in Bionumerics (v. 5.0): A, B, C, D, E and F. Subclusters were designated as O157 or non-O157 depending on whether the number of O157 strains were the majority or minority in the cluster; and presence of the different serotypes was reflected in the SPT designation. Based on this there were two O157 subclusters (O157 SPTA and O157 majority SPTA/B/C). Details of the strain characteristics within each group are given in Supplementary Appendix 2.

Strains within the second O157 majority SPTA/B/C subcluster harboured a majority of O157 strains and a few non-O157 strains; together strains within this subcluster corresponded with serogroups belonging to SPTA, B and C [9] and hence was named O157 majority SPTA/B/C (Fig. 1). It contained 34 O157 strains mostly of H7 or H⁻ complex, of which 15 were from Hungary, nine from Italy, and ≤ 5 strains from the UK, Spain, Denmark and Croatia. There were nine non-O157 strains from the following serogroups: O145, O128, O111, O115, O103, O49, and O26. Eight of the 12 human strains had caused disease; five were O157 strains and the remaining three were from serogroups O26, O128 and O145. All eight strains that had caused disease in humans harboured the *vt2* gene, although this gene was also present in two human isolates not associated with disease. In fact there was no discernible pattern in the

virulence genes tested in the eight isolates that had caused disease in humans from the remaining four that had not. This was a general trend that we noted in the virulence profile of all isolates that had caused disease in humans compared to those that had not but were of human origin; there was little difference between each group although the isolates causing disease showed slightly higher percentage of *eae* and *vt2* genes (Supplementary Appendix 2). There was no dominant 'virulence profile' which defined this group although all strains harboured the *eae* gene, >80% harboured *adfO*, *sodC*, *tccP*, *ureA*, and *terB* genes; and the majority (>60%) were *vt2* and *ehx* positive (Table 3). This is in contrast to the preceding group (O157 SPTA; Table 3), where >90% of strains harboured *vt2*, *ehx*, and the truncated *efa1* gene. In fact in several strains both *vt* genes were absent. Within the panel of *vt*-negative strains 11 were O157:H7

Table 3. Prevalence of the virulence genes in each seropathotype (SPT) group identified within clusters of the PCR data

Cluster	SPT group	% Gene presence											
		<i>eae</i>	<i>vt1</i>	<i>vt2</i>	<i>vt1</i> and <i>vt2</i>	<i>efal-5'</i>	<i>efal-3'</i>	<i>adfO</i>	<i>sodC</i>	<i>ehx</i>	<i>tccP</i>	<i>ureA</i>	<i>terB</i>
E	O157 SPTA	99	50	94	44	97	2	92	100	100	96	97	100
E	O157 dom* SPTA/B/C	100	19	60	6	49	14	91	81	70	81	84	84
C	non-O157/O157 SPTA/D/E	86	5	24	5	19	24	71	100	71	19	33	0
F	non-O157 SPTD/E	100	0	5	0	74	100	84	95	0	0	58	68
F	non-O157 SPTB	100	92	20	14	81	92	89	72	95	1	90	88
B	non-O157 SPTD/E	11	43	56	20	5	10	13	10	18	10	15	26
	All O157 strains	94	41	78	37	79	2	86	90	85	89	86	85
	All non-O157 strains	73	59	32	14	52	63	66	59	63	4	61	63

* dom is the abbreviation for group dominantly containing strains of O157 SPTA serogroup and a few strains from non-O157 serogroups belonging to SPTB and C.

strains of bovine origin from Hungary, of which six were also negative for the *efal* gene (Supplementary Appendix 2).

Cluster C harboured strains from serotype/groups belonging to SPTA, D and E [9], with O157 strains being the most numerous serogroup. However, only 7/21 strains were O157 strains. The incidence of disease in humans was low in this cluster with only 4/10 strains causing disease in humans and three of these belonged to the O157 H7/H⁻ complex. All strains in this cluster harboured the *sodCF* gene and >70% harboured the *eae*, *adfO*, and *ehx* genes; none were *terB* positive, and prevalence of *vt* genes was low (Table 3). The non-O157 strains belonged to the following serogroups/types: O146, O145, O108:H9, O103, O56:H7, O28:H28, and O26. Strains from this cluster were of human, porcine or ovine origin with all eight porcine strains from Hungary, four of which had caused disease in pigs.

Cluster B contained only non-O157 strains with one exception; these were six O157 strains from Hungary of bovine origin and various H-types. These O157 strains were all positive for *tccP* and occasionally positive for the *ureA* genes but negative for all other genes including both *vt* genes and *eae* (Supplementary Appendix 2).

Cramer's *V* statistic was used to determine the within-country relationship of the virulence genes for O157 strains where enough data was available. This was also useful to determine genes that are likely to be associated in strains from different countries (Table 2). Cramer's *V* statistic showed that the 51

Italian strains were significantly different but harboured three perfect correlations indicating that Italian strains possessed *ureA* and *terB* in the presence of *tccP*. The 26 strains from Hungary were also very different but were highly likely to harbour *adfO*, *sodCF*, and *ehxA* when *eae* was present. A large proportion of O157 strains from Hungary did not harbour *vt1*, *vt2*, *efal-5'*, *terB* or *ureA* genes. An association statistic could not be calculated for the 35 Swedish and 16 Spanish strains due to the highly similar virulence patterns they harboured. Therefore, based on the virulence genes used in this study, the O157 strains from Spain and Sweden showed both high within- and between-country (i.e. Sweden and Spain) similarity. Whereas the Italian and Hungarian strains showed very little between-country similarity but some within-country similarity.

Clusters B and F: the main non-O157 group

A defining feature of the non-O157 strains was the virtual absence of *tccP* (96% negative) and low prevalence of *vt2* (32% positive); all other genes were present in ~60% of isolates including the *efal-3'* component. All non-O157 strains within cluster F harboured *eae*, and *tccP* was virtually absent (Table 3). These strains were divided into two major subclusters based on the remaining genotype. Most strains (94/104) within non-O157 SPTB cluster F were from serogroups O111, O26 and O103 and of mostly H⁻, H11 and H2 complexes, respectively, collected from different regions in Europe. The remaining strains

belonged to serogroups O145 (6); O157 (2); O121 (1) and O5 (1). More than 80% of strains representing the prevalent serogroups (O111, O26, O103), harboured genes for *vt1*, *efal-5'*, *efal-3'*, *adfO*, *ehx*, *ureA* and *terB* (Table 3). All strains except two were of human or bovine origin; 24/54 human strains and 5/45 bovine strains had been associated with disease.

In contrast <5% of strains within the non-O157 porcine SPTD/E group within cluster F harboured either *vt* or *ehx* genes but all were *efal-3'* and >80% *adfO* and *sodCF* positive. Fourteen of the 19 strains were of porcine origin from Hungary; six belonged to serotype O4/O123:H11, three to O49:H10/H⁻, three to O45:H11, one to O103:H11 and one to O56. Three strains were human (O157:H8, O111:H2, O128:H2) and two of ovine origin (O26, O56). Seventeen of the 19 strains in this group had caused disease in their host animal.

Forty-nine of the 61 strains within cluster B harboured the *vt* genes (one or both), the remaining virulence genes, including *eae*, were present in <26% of strains (Table 3). The SPT of this group was designated SPTD/E based on serotype/group [9]. Twenty-one strains were of human origin and the remaining strains were of bovine (18), foodstuff (8), porcine (3), ovine (2), wild bird (1) and unknown (8) origin. Seven of the human and one of the porcine strains had caused disease.

Cramer's *V* statistic was used to detect frequencies of the virulence genes harboured by *E. coli* O26 strains from Italy and Sweden, as these datasets were large enough. The correlation matrix found six perfectly positive associations for the 17 Italian strains which indicated that in the presence of *eae*, genes such as *ureA*, *terB* and *efal-3'* were always present; these strains were also highly unlikely to harbour *vt1* and *vt2* together (Table 2). In contrast, all Swedish strains again showed nearly identical virulence characteristics (*eae*⁺, *vt1*⁺, *efal*⁺, *adfO*⁺, *ehx*⁺, *ureA*⁺, *terB*⁺) but an association matrix could not be calculated. Therefore, as with the O157 strains, the Swedish O26 strains showed high within-country similarity, which was distinct from the Italian O26 strains for which only a subset of genes was associated with all strains. There was not sufficient data present for the other non-O157 serogroups for Cramer's *V* statistic to be applied.

DISCUSSION

A virulotyping approach based on the presence of 10 virulence genes was used to assess whether this

may indicate the potential risk posed to humans by 365 strains of both O157 and non-O157 serogroups. The cluster analysis clearly divided O157 and non-O157 strains into separate clusters based on the virulotyping data, which matched well with SPT designations, e.g. the 'high-risk' O157 SPTA strains harboured most of the chosen virulence genes except *efal-3'*, while most of these virulence genes were absent from the 'low-risk' non-O157 SPTD/E strains from cluster B.

However, our virulotyping data show that there is a lack of gradation if only the seropathotyping scheme is used. For instance the 'low-risk' non-O157 porcine SPTD/E strains from cluster F were not distinguished from the non-O157 SPTD/E strains in cluster B, which were also identified as low risk, although they harboured far fewer virulence genes with strains from a variety of human and animal sources, of whom only a few were ill. Interestingly, about 90% of strains within the 'low-risk' non-O157 porcine SPTD/E cluster had caused disease in pigs compared to 50% of porcine strains in cluster C, designated as non-O157/O157 SPTA/D/E, which included some 'high-risk' strains from humans and hence received the SPTA designation. Therefore, the SPT designation although relevant to humans may be less relevant to other animal species. Porcine strains from both groups were from Hungary and a feature distinguishing the non-O157 porcine SPTD/E cluster was the presence in most strains of *terB*, part of the tellurite-resistant operon [28] and complete *efal*, the enterohaemorrhagic *E. coli* adherence factor [29]. The *efal* gene is present on the pathogenicity island termed OI-122, in the EHEC O157 reference strain EDL 933, whose presence has been shown to correlate with severe disease in humans [21]; this data indicates it may also affect pigs.

The SPTA group, the main O157 group, contained O157:H7/H⁻ strains gathered from several different regions of Europe which showed very similar virulence profiles. Their genotype was identical to the EHEC O157 strain reference strain EDL 933 [29]. All Swedish and most Spanish O157 strains were within the O157 SPTA cluster; they were also found to be highly homogenous using Cramer's *V* statistic indicating that they share this subset of important virulence genes, which could not be used to distinguish them further. Although most strains included were randomly chosen, the Swedish and Hungarian O157 strains were from previous studies [30, 31], with pulsed-field gel electrophoresis (PFGE) data available, and reflected their prevalence in the respective

countries. The PFGE data available for the Swedish strains showed that they could be divided further into subgroups with different profiles that have been linked with clinical outcomes [30]. In contrast the virulotype data of the Hungarian O157 strains, which were shown to be particularly heterogeneous from the clustering and Cramer's V statistic, showed a good match with the heterogeneity of the PFGE profiles [31]. Clustering from the PCR virulotyping had placed these O157 strains into three different clusters. Strains from cluster E SPTA/B/C could be matched to PFGE cluster A; from cluster E SPTA to PFGE clusters B/C; and cluster B non-O157 SPTD/E to PFGE cluster D. Therefore, although the 10-gene virulotyping scheme was discriminatory for the latter group, its level of resolution will need to be improved for highly homogenous strains, e.g. the Swedish O157. This could be done by increasing the number of target genes, as shown by Brandt *et al.* [7] who used a 41-gene virulotyping scheme. Even though their data also showed PFGE to be more discriminatory, the PFGE clustering generally corresponded with the PCR virulotyping data.

Recent whole genome sequence analysis of selected strains from non-O157 VTEC serogroups (O26, O111, O103), has shown that they share many virulence genes that are found on prophages and plasmids in O157, which are acquired through horizontal transfer of the mobile elements [32]. Our PCR results indicate that presence of these virulence genes is widespread in strains from these serogroups, which are known to be high risk to humans. We were not able to distinguish between strains only on the basis of human disease as strains with the same virulence genes showed different clinical outcomes (Supplementary Appendix 2), possibly indicating that manifestation of human disease is complicated by other factors such as host immune response. However, there was higher prevalence of *vt2* genes than *vt1* in the O157 group compared to the non-O157 strains of SPTB, which was reflected in disease occurrence: ~60% of human strains within O157 (cluster E) caused disease while ~42% of human strains within the non-O157 SPTB cluster caused disease. Another characteristic seen in the non-O157 group was the low prevalence of the *tccP* gene, which was expected, although it was lower than previously reported (14% [26]) and could be due to only the 1014-bp PCR amplicon being considered. We also observed that there were some differences in the virulence genes harboured by non-O157 strains that were present in cluster E (O157

majority SPTA/B/C) compared to those in cluster F (non-O157 SPTB); notably strains in the former group mainly harboured the truncated *efal* gene and none harboured the *vt1* toxin, being more similar to cluster E O157 strains (Supplementary Appendix 2). It should be noted that strains in cluster B also harboured one or both *vt* genes but showed much lower incidence of disease in strains of human origin, indicating that other virulence factors are required in combination with *vt* genes to cause disease in humans.

The routine use of a simple PCR scheme and SPT for surveillance, as presented here, although less sensitive than PFGE, nevertheless correctly distinguished high-risk *E. coli* pathogenic strains from low(er) risk ones and is much simpler to perform. Therefore, on performing PCR, isolates showing virulence profiles similar to those present in cluster E or cluster F non-O157 SPTB (Table 3), would be expected to be high risk to humans while those with similar profile to cluster B isolates would be expected to be low risk to humans. The resolution of the PCR scheme can be improved if larger numbers of genes are used to virulotype isolates, which need to be small enough to be routinely practicable in reference laboratories. An alternative would be to use the 10-gene PCR virulotyping scheme presented here to identify 'high-risk' strains rapidly, which can then be followed by more discriminatory methods such as the use of larger numbers of virulence gene PCRs or the virulence gene array [33], or even performing PFGE for further subtyping of high-risk strains to enable epidemiological tracing and risk analysis following outbreaks. The use of PCR to distinguish between high- and low-risk strains is not new; a pentaplex PCR panel has been successfully established for routine laboratory use as the minimal predictor of avian pathogenic *E. coli* which may be of high risk to both poultry and humans [34].

SUPPLEMENTARY MATERIAL

For supplementary material accompanying this paper visit <http://dx.doi.org/10.1017/S0950268813001635>.

ACKNOWLEDGEMENTS

The authors acknowledge the Med-Vet-Net network of excellence (WP26) for funding the studies. The authors are grateful to Mr Mark Saunders for his contribution in consolidating the PCR data and the *E. coli* Reference Laboratory at the Animal

Health and Veterinary Laboratories Agency for contributing strains for this study.

DECLARATION OF INTEREST

None.

REFERENCES

- Nataro JP, Kaper JB. Diarrheagenic *Escherichia coli*. *Clinical Microbiology Reviews* 1998; **11**: 142–201.
- Griffin PM, Tauxe RV. The epidemiology of infections caused by *Escherichia coli* O157:H7, other enterohemorrhagic *E. coli*, and the associated hemolytic uremic syndrome. *Epidemiologic Reviews* 1991; **13**: 60–98.
- Karmali MA. Infection by verocytotoxin-producing *Escherichia coli*. *Clinical Microbiology Reviews* 1989; **2**: 15–38.
- European Centre for Disease Control and Prevention. ECDC surveillance report, *Quarterly STEC/VTEC Report Q1 2008*, January–March 2008.
- World Health Organisation. Zoonotic non-O157 shiga toxin-producing *Escherichia coli* (STEC). Report of a WHO Scientific Working Group Meeting, Berlin, 23–26 June 1998.
- Coombes BK, et al. Molecular analysis as an aid to assess the public health risk of non-O157 Shiga toxin-producing *Escherichia coli* strains. *Applied and Environmental Microbiology* 2008; **74**: 2153–2160.
- Brandt SM, et al. Molecular risk assessment and epidemiological typing of Shiga toxin-producing *Escherichia coli* by using a novel PCR binary typing system. *Applied and Environmental Microbiology* 2011; **77**: 2458–2470.
- Bugarel M, Beutin L, Fach P. Low-density microarray targeting non-locus of enterocyte effacement effectors (nle genes) and major virulence factors of Shiga toxin-producing *Escherichia coli* (STEC): a new approach for molecular risk assessment of STEC isolates. *Applied and Environmental Microbiology* 2010; **76**: 203–211.
- Karmali MA, et al. Association of genomic O island 122 of *Escherichia coli* EDL 933 with verocytotoxin-producing *Escherichia coli* seropathotypes that are linked to epidemic and/or serious disease. *Journal of Clinical Microbiology* 2003; **41**: 4930–4940.
- Lee JH, Hur J, Stein BD. Occurrence and characteristics of enterohemorrhagic *Escherichia coli* O26 and O111 in calves associated with diarrhea. *Veterinary Journal* 2008; **176**: 205–209.
- Badouei MA, et al. Virulence gene profiles and intimin subtypes of Shiga toxin-producing *Escherichia coli* isolated from healthy and diarrhoeic calves. *Veterinary Record* 2010; **167**: 858–861.
- Hutchinson JP, et al. Verocytotoxin-producing and attaching and effacing activity of *Escherichia coli* isolated from diseased farm livestock. *Veterinary Record* 2011; **168**: 536.
- Scheutz F, et al. Designation of O174 and O175 to temporary O groups OX3 and OX7, and six new *E. coli* O groups that include Verocytotoxin-producing *E. coli* (VTEC): O176, O177, O178, O179, O180 and O181. *Acta Pathologica, Microbiologica et Immunologica Scandinavica* 2004; **112**: 569–584.
- Garabal JI, et al. Serogroups of *Escherichia coli* isolated from piglets in Spain. *Veterinary Microbiology* 1996; **48**: 113–123.
- Renter DG, et al. Prevalence, risk factors, O serogroups, and virulence profiles of Shiga toxin-producing bacteria from cattle production environments. *Journal of Food Protection* 2005; **68**: 1556–1565.
- DebRoy C, et al. Multiplex polymerase chain reaction assay for detection of nonserotypable Shiga toxin-producing *Escherichia coli* strains of serogroup O147. *Foodborne Pathogens and Disease* 2010; **7**: 1407–1414.
- Manna SK, et al. Serogroup distribution and virulence characteristics of sorbitol-negative *Escherichia coli* from food and cattle stool. *Journal of Applied Microbiology* 2010; **108**: 658–665.
- Iyoda S, et al. Inducible *stx2* phages are lysogenized in the enteroaggregative and other phenotypic *Escherichia coli* O86:HNM isolated from patients. *FEMS Microbiology Letters* 2000; **191**: 7–10.
- Orden JA, et al. Verotoxin-producing *Escherichia coli* (VTEC) and *eae*-positive non-VTEC in 1-30-days-old diarrhoeic dairy calves. *Veterinary Microbiology* 1998; **63**: 239–248.
- Mellmann A, et al. Prospective genomic characterization of the German enterohemorrhagic *Escherichia coli* O104:H4 outbreak by rapid next generation sequencing technology. *PLoS One* 2011; **6**: e22751.
- Morabito S, et al. A mosaic pathogenicity island made up of the locus of enterocyte effacement and a pathogenicity island of *Escherichia coli* O157:H7 is frequently present in attaching and effacing *E. coli*. *Infection and Immunity* 2003; **71**: 3343–3348.
- Imamovic L, et al. OI-57, a genomic island of *Escherichia coli* O157, is present in other seropathotypes of Shiga toxin-producing *E. coli* associated with severe human disease. *Infection and Immunity* 2010; **78**: 4697–4704.
- D’Orazio M, et al. Regulatory and structural properties differentiating the chromosomal and the bacteriophage-associated *Escherichia coli* O157:H7 Cu, Zn superoxide dismutases. *BMC Microbiology* 2008; **8**: 166.
- Wieler LH, et al. Shiga toxin-producing *Escherichia coli* strains from bovines: association of adhesion with carriage of *eae* and other genes. *Journal of Clinical Microbiology* 1996; **34**: 2980–2984.
- Russmann H, et al. Genotyping of Shiga-like toxin genes in non-O157 *Escherichia coli* strains associated with haemolytic uraemic syndrome. *Journal of Medical Microbiology* 1995; **42**: 404–410.
- Garmendia J, et al. Distribution of *tccP* in clinical enterohemorrhagic and enteropathogenic *Escherichia coli* isolates. *Journal of Clinical Microbiology* 2005; **43**: 5715–5720.

27. **Friedrich AW, et al.** Distribution of the urease gene cluster among and urease activities of enterohemorrhagic *Escherichia coli* O157 isolates from humans. *Journal of Clinical Microbiology* 2005; **43**: 546–550.
28. **Taylor DE, et al.** Genomic variability of O islands encoding tellurite resistance in enterohemorrhagic *Escherichia coli* O157:H7 isolates. *Journal of Bacteriology* 2002; **184**: 4690–4698.
29. **Perna NT, et al.** Genome sequence of enterohaemorrhagic *Escherichia coli* O157:H7. *Nature* 2001; **409**: 529–533.
30. **Eriksson E, et al.** Genotypic characterization to identify markers associated with putative hypervirulence in Swedish *Escherichia coli* O157:H7 cattle strains. *Journal of Applied Microbiology* 2011; **110**: 323–332.
31. **Toth I, et al.** Virulence genes and molecular typing of different groups of *Escherichia coli* O157 strains in cattle. *Applied and Environmental Microbiology* 2009; **75**: 6282–6291.
32. **Ogura Y, et al.** Comparative genomics reveal the mechanism of the parallel evolution of O157 and non-O157 enterohemorrhagic *Escherichia coli*. *Proceedings of the National Academy of Sciences USA* 2009; **106**: 17 939–1744.
33. **Anjum MF, et al.** Pathotyping *Escherichia coli* by using miniaturized DNA microarrays. *Applied and Environmental Microbiology* 2007; **73**: 5692–5697.
34. **Johnson TJ, et al.** Identification of minimal predictors of avian pathogenic *Escherichia coli* virulence for use as a rapid diagnostic tool. *Journal of Clinical Microbiology* 2008; **46**: 3987–3996.