

ON THE RATIONAL SOLUTIONS OF q -PAINLEVÉ V EQUATION

TETSU MASUDA

Abstract. We give an explicit determinant formula for a class of rational solutions of a q -analogue of the Painlevé V equation. The entries of the determinant are given by the continuous q -Laguerre polynomials.

§1. Introduction

Since the introduction of the singularity confinement criterion as the discrete analogue of the Painlevé test [2], a lot of ordinary difference equations have been proposed as discrete Painlevé equations [12], [1]. It is known that the discrete Painlevé equations possess several properties analogous to the continuous ones such as the coalescence cascade, symmetry as the Bäcklund transformations and particular solutions.

Recently, Kajiwara, Noumi and Yamada have proposed a q -analogue of the Painlevé IV equation [3], and investigated the structure of symmetry and special solutions of the q -P_{IV}. It has been shown that the q -P_{IV} admits two types of special solutions; one is the special function type solutions, which are expressed in terms of the continuous q -Hermite-Weber functions, and another is the rational solutions expressed as the ratio of a q -analogue of Okamoto polynomials [11].

In this paper, we consider the symmetric form of q -P_V

$$\begin{aligned}
 \bar{a}_0 &= a_0, & \bar{a}_1 &= a_1, & \bar{a}_2 &= a_2, & \bar{a}_3 &= a_3, \\
 \bar{f}_0 &= a_0 a_1 f_1 \frac{1 + a_2 f_2 + a_2 a_3 f_2 f_3 + a_2 a_3 a_0 f_2 f_3 f_0}{1 + a_0 f_0 + a_0 a_1 f_0 f_1 + a_0 a_1 a_2 f_0 f_1 f_2}, \\
 \bar{f}_1 &= a_1 a_2 f_2 \frac{1 + a_3 f_3 + a_3 a_0 f_3 f_0 + a_3 a_0 a_1 f_3 f_0 f_1}{1 + a_1 f_1 + a_1 a_2 f_1 f_2 + a_1 a_2 a_3 f_1 f_2 f_3}, \\
 \bar{f}_2 &= a_2 a_3 f_3 \frac{1 + a_0 f_0 + a_0 a_1 f_0 f_1 + a_0 a_1 a_2 f_0 f_1 f_2}{1 + a_2 f_2 + a_2 a_3 f_2 f_3 + a_2 a_3 a_0 f_2 f_3 f_0},
 \end{aligned}
 \tag{1.1}$$

Received July 25, 2001.

2000 Mathematics Subject Classification: 33E17; 34M55.

$$\bar{f}_3 = a_3 a_0 f_0 \frac{1 + a_1 f_1 + a_1 a_2 f_1 f_2 + a_1 a_2 a_3 f_1 f_2 f_3}{1 + a_3 f_3 + a_3 a_0 f_3 f_0 + a_3 a_0 a_1 f_3 f_0 f_1},$$

with

$$(1.2) \quad a_0 a_1 a_2 a_3 = q^{-1},$$

where $\bar{}$ stands for the discrete time evolution. Introducing a variable c by

$$(1.3) \quad f_0 f_2 = f_1 f_3 = c^{-1},$$

we find that c plays a role of the independent variable,

$$(1.4) \quad \bar{c} = qc.$$

Originally, the equation (1.1) is derived as a subsystem of the discrete dynamical systems associated with the extended affine Weyl group symmetry of type $A_{m-1}^{(1)} \times A_{n-1}^{(1)}$ [4]. In the case of $(m, n) = (2, 4)$, by regarding a translation of $\widetilde{W}(A_1^{(1)})$ as the discrete time evolution $\bar{}$, we obtain the system (1.1). The variables q and c are invariant for the action of $\widetilde{W}(A_1^{(1)} \times A_3^{(1)})$ and $\widetilde{W}(A_3^{(1)})$, respectively. The inverse time evolution of (1.1) is given by

$$(1.5) \quad \begin{aligned} \underline{a}_0 &= a_0, & \underline{a}_1 &= a_1, & \underline{a}_2 &= a_2, & \underline{a}_3 &= a_3, \\ \underline{f}_0 &= \frac{f_3}{a_0 a_1} \frac{a_2 a_1 a_0 + a_1 a_0 f_2 + a_0 f_2 f_1 + f_2 f_1 f_0}{a_0 a_3 a_2 + a_3 a_2 f_0 + a_2 f_0 f_3 + f_0 f_3 f_2}, \\ \underline{f}_1 &= \frac{f_0}{a_1 a_2} \frac{a_3 a_2 a_1 + a_2 a_1 f_3 + a_1 f_3 f_2 + f_3 f_2 f_1}{a_1 a_0 a_3 + a_0 a_3 f_1 + a_3 f_1 f_0 + f_1 f_0 f_3}, \\ \underline{f}_2 &= \frac{f_1}{a_2 a_3} \frac{a_0 a_3 a_2 + a_3 a_2 f_0 + a_2 f_0 f_3 + f_0 f_3 f_2}{a_2 a_1 a_0 + a_1 a_0 f_2 + a_0 f_2 f_1 + f_2 f_1 f_0}, \\ \underline{f}_3 &= \frac{f_0}{a_3 a_0} \frac{a_1 a_0 a_3 + a_0 a_3 f_1 + a_3 f_1 f_0 + f_1 f_0 f_3}{a_3 a_2 a_1 + a_2 a_1 f_3 + a_1 f_3 f_2 + f_3 f_2 f_1}. \end{aligned}$$

The reason why we refer to the discrete system (1.1) as the symmetric form of q -P_V is as follows. By the original construction in [4], it is clear that this equation admits the affine Weyl group symmetry of type $A_3^{(1)}$ as the Bäcklund transformation group, which is stated in Section 3 precisely. Moreover, the system (1.1) reduces to the symmetric form of the Painlevé V equation in the continuum limit. We set

$$(1.6) \quad q = e^{\frac{\varepsilon^2}{2}}, \quad a_i = e^{-\frac{\varepsilon^2}{2} \alpha_i}, \quad f_i = -e^{-\varepsilon \varphi_i}, \quad c = e^{\varepsilon \gamma},$$

and define the derivation $\frac{d}{ds}$ by

$$(1.7) \quad \frac{dz}{ds} = \lim_{\varepsilon \rightarrow 0} \frac{\bar{z} - z}{\varepsilon},$$

for a function z in α_i and φ_i . Then, we get from (1.1) and (1.4)

$$(1.8) \quad \frac{d\varphi_0}{ds} = \frac{1}{\gamma} \left[\varphi_0\varphi_2(\varphi_1 - \varphi_3) + \left(\frac{1}{2} - \alpha_2\right)\varphi_0 + \alpha_0\varphi_2 \right], \quad \frac{d\gamma}{ds} = \frac{1}{2}.$$

Introducing the variable t and derivation $'$ as

$$(1.9) \quad \gamma = \sqrt{t}, \quad ' = t \frac{d}{dt},$$

we have

$$(1.10) \quad \begin{aligned} \alpha'_0 &= 0, & \alpha'_1 &= 0, & \alpha'_2 &= 0, & \alpha'_3 &= 0, \\ \varphi'_0 &= \varphi_0\varphi_2(\varphi_1 - \varphi_3) + \left(\frac{1}{2} - \alpha_2\right)\varphi_0 + \alpha_0\varphi_2, \\ \varphi'_1 &= \varphi_1\varphi_3(\varphi_2 - \varphi_0) + \left(\frac{1}{2} - \alpha_3\right)\varphi_1 + \alpha_1\varphi_3, \\ \varphi'_2 &= \varphi_2\varphi_0(\varphi_3 - \varphi_1) + \left(\frac{1}{2} - \alpha_0\right)\varphi_2 + \alpha_2\varphi_0, \\ \varphi'_3 &= \varphi_3\varphi_1(\varphi_0 - \varphi_2) + \left(\frac{1}{2} - \alpha_1\right)\varphi_3 + \alpha_3\varphi_1. \end{aligned}$$

The normalization conditions (1.2) and (1.3) reduce to

$$(1.11) \quad \alpha_0 + \alpha_1 + \alpha_2 + \alpha_3 = 1,$$

and

$$(1.12) \quad \varphi_0 + \varphi_2 = \varphi_1 + \varphi_3 = \sqrt{t},$$

respectively. This differential system (1.10)–(1.12) is nothing but the symmetric form of P_V [10].

On the other hand, it has been revealed that a family of the rational solutions of P_V , which exists on the barycenters of Weyl chambers, has a determinant formula whose entries are the Laguerre polynomials [8]. This determinant expression is regarded as a specialization of the universal characters [7]. The aim of this paper is to present an explicit determinant formula for a class of the rational solutions of q - P_V .

This paper is organized as follows. In Section 2, we give the main result of this paper. In Section 3, we describe the affine Weyl group symmetry and derive a set of bilinear equations for the τ -functions of q -PV. In Section 4, we construct the rational solutions of q -PV. Proof of our result is given in Section 5. Section 6 is devoted to some remarks.

§2. Main result

DEFINITION 2.1. Let $p_k^{(b)} = p_k^{(b)}(y|q)$ and $q_k^{(b)} = q_k^{(b)}(y|q)$, $k \in \mathbb{Z}$, be two sets of polynomials defined by

$$(2.1) \quad \sum_{k=0}^{\infty} p_k^{(b)} \lambda^k = \frac{(q^{\frac{1}{4}} b \lambda, q^{\frac{3}{4}} b \lambda; q)_{\infty}}{(-q^{\frac{1}{4}} x \lambda, -q^{\frac{3}{4}} x^{-1} \lambda; q)_{\infty}}, \quad p_k^{(b)} = 0 \text{ for } k < 0,$$

$$\sum_{k=0}^{\infty} q_k^{(b)} \lambda^k = \frac{(-q^{\frac{1}{4}} x \lambda, -q^{\frac{3}{4}} x^{-1} \lambda; q)_{\infty}}{(q^{\frac{1}{4}} b^{-1} \lambda, q^{\frac{3}{4}} b^{-1} \lambda; q)_{\infty}}, \quad q_k^{(b)} = 0 \text{ for } k < 0,$$

with

$$(2.2) \quad y = -\frac{1}{2}(q^{-1/4}x + q^{1/4}x^{-1}).$$

For $m, n \in \mathbb{Z}_{\geq 0}$, we define a family of polynomials $R_{m,n}^{(b)} = R_{m,n}^{(b)}(y|q)$ by

$$(2.3) \quad R_{m,n}^{(b)} = \begin{vmatrix} q_1^{(b)} & q_0^{(b)} & \cdots & q_{-m+2}^{(b)} & q_{-m+1}^{(b)} & \cdots & q_{-m-n+3}^{(b)} & q_{-m-n+2}^{(b)} \\ q_3^{(b)} & q_2^{(b)} & \cdots & q_{-m+4}^{(b)} & q_{-m+3}^{(b)} & \cdots & q_{-m-n+5}^{(b)} & q_{-m-n+4}^{(b)} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots \\ q_{2m-1}^{(b)} & q_{2m-2}^{(b)} & \cdots & q_m^{(b)} & q_{m-1}^{(b)} & \cdots & q_{m-n+1}^{(b)} & q_{m-n}^{(b)} \\ p_{n-m}^{(b)} & p_{n-m+1}^{(b)} & \cdots & p_{n-1}^{(b)} & p_n^{(b)} & \cdots & p_{2n-2}^{(b)} & p_{2n-1}^{(b)} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots \\ p_{-n-m+4}^{(b)} & p_{-n-m+5}^{(b)} & \cdots & p_{-n+3}^{(b)} & p_{-n+4}^{(b)} & \cdots & p_2^{(b)} & p_3^{(b)} \\ p_{-n-m+2}^{(b)} & p_{-n-m+3}^{(b)} & \cdots & p_{-n+1}^{(b)} & p_{-n+2}^{(b)} & \cdots & p_0^{(b)} & p_1^{(b)} \end{vmatrix}.$$

For $m, n \in \mathbb{Z}_{< 0}$, we define $R_{m,n}^{(b)}$ by

$$(2.4) \quad R_{m,n}^{(b)} = (-1)^{m(m+1)/2} R_{-m-1,n}^{(b)}, \quad R_{m,n}^{(b)} = (-1)^{n(n+1)/2} R_{m,-n-1}^{(b)}.$$

Remark 2.2. The polynomials $p_k^{(b)}$ and $q_k^{(b)}$ are essentially the continuous q -Laguerre polynomials $P_k^{(\alpha)}(y|q)$, which is defined by [6]

$$(2.5) \quad \sum_{k=0}^{\infty} P_k^{(\alpha)}(y|q) \lambda^k = \frac{(q^{\alpha+\frac{1}{2}} \lambda, q^{\alpha+1} \lambda; q)_{\infty}}{(q^{\frac{1}{2}\alpha+\frac{1}{4}} e^{i\theta} \lambda, q^{\frac{1}{2}\alpha+\frac{1}{4}} e^{-i\theta} \lambda; q)_{\infty}},$$

$$P_k^{(\alpha)}(y|q) = 0 \text{ for } k < 0, \quad y = \cos \theta.$$

In fact, denoting as

$$(2.6) \quad L_k(y, b|q) = P_k^{(\alpha)}(y|q), \quad b = q^{\frac{1}{2}\alpha + \frac{1}{4}},$$

we see that $p_k^{(b)}$ and $q_k^{(b)}$ are expressed as

$$(2.7) \quad \begin{aligned} p_k^{(b)}(y|q) &= (q^{\frac{3}{4}}b^{-1})^k L_k(y, q^{-\frac{1}{4}}b|q), \\ q_k^{(b)}(y|q) &= (q^{-\frac{3}{4}}b)^k L_k(y, q^{\frac{1}{4}}b^{-1}|q^{-1}), \end{aligned}$$

respectively.

Our main result is stated as follows.

THEOREM 2.3. *We set*

$$(2.8) \quad S_{m,n}(x, a) = R_{m,n}^{(b)}(y), \quad y = -\frac{1}{2}(q^{-1/4}x + q^{1/4}x^{-1}), \quad b = q^{-\frac{1}{2}(m-n)}a.$$

Then, for the parameters

$$(2.9) \quad (a_0, a_1, a_2, a_3) = (q^{n-\frac{1}{2}}a, a^{-1}, q^{-m-\frac{1}{2}}a, q^{m-n}a^{-1}),$$

we have the following rational solutions of q -P_V,

$$(2.10) \quad \begin{aligned} &1 + q^{\frac{1}{2}(2n-1)}af_0(x, a) \\ &= q^{\frac{1}{2}m}(1 + q^{-\frac{1}{2}(m-n+1)}ax^{-1}) \frac{S_{m,n}(x, a)S_{m-1,n-1}(q^{\frac{1}{2}}x, q^{-1}a)}{S_{m,n-1}(q^{\frac{1}{2}}x, a)S_{m-1,n}(x, q^{-1}a)}, \\ &1 + a^{-1}f_1(x, a) \\ &= q^{\frac{1}{2}n}(1 + q^{\frac{1}{2}(m-n)}a^{-1}x^{-1}) \frac{S_{m-1,n}(x, q^{-1}a)S_{m,n-1}(q^{\frac{1}{2}}x, qa)}{S_{m,n}(q^{\frac{1}{2}}x, a)S_{m-1,n-1}(x, a)}, \\ &1 + q^{-\frac{1}{2}(2m+1)}af_2(x, a) \\ &= q^{-\frac{1}{2}m}(1 + q^{-\frac{1}{2}(m-n+1)}ax^{-1}) \frac{S_{m-1,n-1}(x, a)S_{m,n}(q^{\frac{1}{2}}x, q^{-1}a)}{S_{m-1,n}(q^{\frac{1}{2}}x, q^{-1}a)S_{m,n-1}(x, a)}, \\ &1 + q^{m-n}a^{-1}f_3(x, a) \\ &= q^{-\frac{1}{2}n}(1 + q^{\frac{1}{2}(m-n)}a^{-1}x^{-1}) \frac{S_{m,n-1}(x, a)S_{m-1,n}(q^{\frac{1}{2}}x, a)}{S_{m-1,n-1}(q^{\frac{1}{2}}x, a)S_{m,n}(x, a)}, \end{aligned}$$

with $x^2 = c$. Moreover, the above solutions admit the other expressions as

$$\begin{aligned}
 (2.11) \quad & 1 + q^{-\frac{1}{2}(2n-1)}a^{-1}f_0(x, a) \\
 &= q^{-\frac{1}{2}n}(1 + q^{\frac{1}{2}(m-n+1)}a^{-1}x^{-1}) \frac{S_{m,n}(q^{\frac{1}{2}}x, a)S_{m-1,n-1}(x, q^{-1}a)}{S_{m,n-1}(q^{\frac{1}{2}}x, a)S_{m-1,n}(x, q^{-1}a)}, \\
 & 1 + af_1(x, a) \\
 &= q^{\frac{1}{2}m}(1 + q^{-\frac{1}{2}(m-n)}ax^{-1}) \frac{S_{m-1,n}(q^{\frac{1}{2}}x, q^{-1}a)S_{m,n-1}(x, qa)}{S_{m,n}(q^{\frac{1}{2}}x, a)S_{m-1,n-1}(x, a)}, \\
 & 1 + q^{\frac{1}{2}(2m+1)}a^{-1}f_2(x, a) \\
 &= q^{\frac{1}{2}n}(1 + q^{\frac{1}{2}(m-n+1)}a^{-1}x^{-1}) \frac{S_{m-1,n-1}(q^{\frac{1}{2}}x, a)S_{m,n}(x, q^{-1}a)}{S_{m-1,n}(q^{\frac{1}{2}}x, q^{-1}a)S_{m,n-1}(x, a)}, \\
 & 1 + q^{-m+n}af_3(x, a) \\
 &= q^{-\frac{1}{2}m}(1 + q^{-\frac{1}{2}(m-n)}ax^{-1}) \frac{S_{m,n-1}(q^{\frac{1}{2}}x, a)S_{m-1,n}(x, a)}{S_{m-1,n-1}(q^{\frac{1}{2}}x, a)S_{m,n}(x, a)}.
 \end{aligned}$$

Remark 2.4. The rational solutions of q - P_V in Theorem 2.3 are a q -analogue of those to the Painlevé V equation [8]. See Appendix C in detail.

§3. Weyl group symmetry and bilinear relations

As we mentioned in Section 1, the symmetric form of q - P_V (1.1) admits the symmetry of the extended affine Weyl group $\widetilde{W} = \langle s_0, s_1, s_2, s_3, \pi \rangle$ of type $A_3^{(1)}$ as a group of Bäcklund transformations. The action of s_i and π on the variables a_i and f_i is given by

$$(3.1) \quad s_i(a_j) = a_j a_i^{-a_{ij}}, \quad \pi(a_j) = a_{j+1},$$

$$(3.2) \quad s_i(f_j) = f_j \left(\frac{a_i + f_i}{1 + a_i f_i} \right)^{u_{ij}}, \quad \pi(f_j) = f_{j+1},$$

where $A = (a_{ij})_{i,j=0}^3$ is the generalized Cartan matrix of type $A_3^{(1)}$ and $U = (u_{ij})_{i,j=0}^3$ is an orientation matrix of the corresponding Dynkin diagram

$$(3.3) \quad A = \begin{pmatrix} 2 & -1 & 0 & -1 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -1 \\ -1 & 0 & -1 & 2 \end{pmatrix}, \quad U = \begin{pmatrix} 0 & 1 & 0 & -1 \\ -1 & 0 & 1 & 0 \\ 0 & -1 & 0 & 1 \\ 1 & 0 & -1 & 0 \end{pmatrix},$$

and indices are understood as elements of $\mathbb{Z}/4\mathbb{Z}$. These transformations commute with the time evolution and satisfy the fundamental relations

$$(3.4) \quad s_i^2 = 1, \quad s_i s_j = s_j s_i \quad (j \neq i, i \pm 1), \quad s_i s_j s_i = s_j s_i s_j \quad (j = i \pm 1), \\ \pi^4 = 1, \quad \pi s_j = s_{j+1} \pi.$$

Let us introduce τ -functions τ_i as solutions of the following equations [5],

$$(3.5) \quad \bar{\tau}_i = g_i \frac{\bar{\tau}_i \bar{\tau}_{i+1}}{\tau_{i+1}},$$

where g_i is given by

$$(3.6) \quad g_i = 1 + a_{i+1} f_{i+1} + a_{i+1} a_{i+2} f_{i+1} f_{i+2} + a_{i+1} a_{i+2} a_{i+3} f_{i+1} f_{i+2} f_{i+3}.$$

The inverse transformations are given as,

$$(3.7) \quad \underline{\tau}_i = h_i \frac{\tau_{i-1} \tau_i}{\bar{\tau}_{i-1}},$$

with

$$(3.8) \quad h_i = 1 + \frac{f_{i-1}}{a_{i-1}} + \frac{f_{i-1} f_{i-2}}{a_{i-1} a_{i-2}} + \frac{f_{i-1} f_{i-2} f_{i-3}}{a_{i-1} a_{i-2} a_{i-3}}.$$

The Bäcklund transformations can be lifted on the τ -functions as follows:

$$(3.9) \quad s_i(\tau_i) = \left(1 + \frac{f_i}{a_i}\right) \frac{\bar{\tau}_{i-1} \tau_{i+1}}{\bar{\tau}_i}, \quad s_i(\bar{\tau}_i) = (1 + a_i f_i) \frac{\bar{\tau}_{i-1} \tau_{i+1}}{\tau_i}, \\ s_i(\tau_j) = \tau_j, \quad s_i(\bar{\tau}_j) = \bar{\tau}_j, \quad (i \neq j), \\ \pi(\tau_j) = \tau_{j+1}, \quad \pi(\bar{\tau}_j) = \bar{\tau}_{j+1}.$$

The fundamental relations (3.4) are preserved in this lifting. Note that we have the multiplicative formulas

$$(3.10) \quad 1 + \frac{f_i}{a_i} = \frac{\bar{\tau}_i s_i(\tau_i)}{\bar{\tau}_{i-1} \tau_{i+1}}, \quad 1 + a_i f_i = \frac{\tau_i s_i(\bar{\tau}_i)}{\bar{\tau}_{i-1} \tau_{i+1}},$$

for the independent variables f_i in terms of τ -functions.

PROPOSITION 3.1. *We have the following bilinear equations:*

$$\begin{aligned}
 \tau_0 s_0 s_1(\bar{\tau}_1) &= a_0^2 s_0(\tau_0) s_1(\bar{\tau}_1) + (1 - a_0^2) \tau_2 \bar{\tau}_3, \\
 \bar{\tau}_1 s_1 s_0(\tau_0) &= a_1^{-2} s_0(\tau_0) s_1(\bar{\tau}_1) + (1 - a_1^{-2}) \tau_2 \bar{\tau}_3, \\
 \tau_1 s_1 s_2(\bar{\tau}_2) &= a_1^2 s_1(\tau_1) s_2(\bar{\tau}_2) + (1 - a_1^2) \tau_3 \bar{\tau}_0, \\
 \bar{\tau}_2 s_2 s_1(\tau_1) &= a_2^{-2} s_1(\tau_1) s_2(\bar{\tau}_2) + (1 - a_2^{-2}) \tau_3 \bar{\tau}_0, \\
 \tau_2 s_2 s_3(\bar{\tau}_3) &= a_2^2 s_2(\tau_2) s_3(\bar{\tau}_3) + (1 - a_2^2) \tau_3 \bar{\tau}_1, \\
 \bar{\tau}_3 s_3 s_2(\tau_2) &= a_3^{-2} s_2(\tau_2) s_3(\bar{\tau}_3) + (1 - a_3^{-2}) \tau_0 \bar{\tau}_1, \\
 \tau_3 s_3 s_0(\bar{\tau}_0) &= a_3^2 s_3(\tau_3) s_0(\bar{\tau}_0) + (1 - a_3^2) \tau_1 \bar{\tau}_2, \\
 \bar{\tau}_0 s_0 s_3(\tau_3) &= a_0^{-2} s_3(\tau_3) s_0(\bar{\tau}_0) + (1 - a_0^{-2}) \tau_1 \bar{\tau}_2.
 \end{aligned}
 \tag{3.11}$$

Proof. Eliminating f_0 from (3.9) with $i = 0$, we obtain

$$1 - a_0^2 \frac{\bar{\tau}_0 s_0(\tau_0)}{\tau_0 s_0(\bar{\tau}_0)} = (1 - a_0^2) \frac{\bar{\tau}_3 \tau_1}{\tau_0 s_0(\bar{\tau}_0)}.
 \tag{3.12}$$

From (3.2) and (3.9), we get

$$\frac{\tau_1 s_0 s_1(\bar{\tau}_1)}{s_0(\bar{\tau}_0) \tau_2} = 1 + a_0^2 \left(\frac{\tau_1 s_1(\bar{\tau}_1)}{\bar{\tau}_0 \tau_2} - 1 \right) \frac{\bar{\tau}_0 s_0(\tau_0)}{\tau_0 s_0(\bar{\tau}_0)},
 \tag{3.13}$$

which leads to the first equation of (3.11) by using (3.12). The other equations are derived by the similar way. □

Let us define the translation operators T_i ($i = 0, 1, 2, 3$) by

$$T_1 = \pi s_3 s_2 s_1, \quad T_2 = s_1 \pi s_3 s_2, \quad T_3 = s_2 s_1 \pi s_3, \quad T_0 = s_3 s_2 s_1 \pi,
 \tag{3.14}$$

which commute with each other and satisfy $T_1 T_2 T_3 T_0 = 1$. These operators act on parameters a_i as

$$T_i(a_{i-1}) = q^{-1} a_{i-1}, \quad T_i(a_i) = q a_i, \quad T_i(a_j) = a_j \quad (j \neq i - 1, i).
 \tag{3.15}$$

In terms of T_i , τ -functions in (3.11) are expressed as

$$\begin{aligned}
 \tau_1 &= T_1(\tau_0), & \tau_2 &= T_1 T_2(\tau_0), & \tau_3 &= T_0^{-1}(\tau_0), \\
 s_0(\tau_0) &= T_0^{-1} T_1(\tau_0), & s_1(\tau_1) &= T_2(\tau_0), & s_2(\tau_2) &= T_1 T_3(\tau_0), \\
 s_3(\tau_3) &= T_3^{-1}(\tau_0), & s_0 s_1(\tau_1) &= T_1 T_2 T_0^{-1}(\tau_0), & s_1 s_0(\tau_0) &= T_2 T_0^{-1}(\tau_0), \\
 s_1 s_2(\tau_2) &= T_2 T_3(\tau_0), & s_2 s_1(\tau_1) &= T_3(\tau_0), & s_2 s_3(\tau_3) &= T_2^{-1}(\tau_0), \\
 s_3 s_2(\tau_2) &= T_1 T_0(\tau_0), & s_3 s_0(\tau_0) &= T_1 T_3^{-1}(\tau_0), & s_0 s_3(\tau_3) &= T_1 T_3^{-1} T_0^{-1}(\tau_0).
 \end{aligned}
 \tag{3.16}$$

Introducing a notation,

$$(3.17) \quad \tau_{l,m,n} = T_2^l T_3^m T_0^n(\tau_0), \quad \bar{\tau}_{l,m,n} = T_2^l T_3^m T_0^n(\bar{\tau}_0), \quad l, m, n \in \mathbb{Z},$$

we can express the bilinear relations (3.11) as

$$(3.18) \quad \begin{aligned} \tau_{l,m,n} \bar{\tau}_{l,m-1,n-2} &= a_0^2 q^{2n} \tau_{l-1,m-1,n-2} \bar{\tau}_{l+1,m,n} \\ &\quad + (1 - a_0^2 q^{2n}) \tau_{l,m-1,n-1} \bar{\tau}_{l,m,n-1}, \\ \bar{\tau}_{l-1,m-1,n-1} \tau_{l+1,m,n-1} &= a_1^{-2} q^{2l} \tau_{l-1,m-1,n-2} \bar{\tau}_{l+1,m,n} \\ &\quad + (1 - a_1^{-2} q^{2l}) \tau_{l,m-1,n-1} \bar{\tau}_{l,m,n-1}, \\ \tau_{l-1,m-1,n-1} \bar{\tau}_{l+1,m+1,n} &= a_1^2 q^{-2l} \tau_{l+1,m,n} \bar{\tau}_{l-1,m,n-1} \\ &\quad + (1 - a_1^2 q^{-2l}) \tau_{l,m,n-1} \bar{\tau}_{l,m,n}, \\ \bar{\tau}_{l,m-1,n-1} \tau_{l,m+1,n} &= a_2^{-2} q^{2(-l+m)} \tau_{l+1,m,n} \bar{\tau}_{l-1,m,n-1} \\ &\quad + (1 - a_2^{-2} q^{2(-l+m)}) \tau_{l,m,n-1} \bar{\tau}_{l,m,n}, \\ \tau_{l,m-1,n-1} \bar{\tau}_{l-1,m,n} &= a_2^2 q^{2(l-m)} \tau_{l-1,m,n-1} \bar{\tau}_{l,m-1,n} \\ &\quad + (1 - a_2^2 q^{2(l-m)}) \tau_{l,m,n} \bar{\tau}_{l-1,m-1,n-1}, \\ \bar{\tau}_{l,m,n-1} \tau_{l-1,m-1,n} &= a_3^{-2} q^{2(-m+n)} \tau_{l-1,m,n-1} \bar{\tau}_{l,m-1,n} \\ &\quad + (1 - a_3^{-2} q^{2(-m+n)}) \tau_{l,m,n} \bar{\tau}_{l-1,m-1,n-1}, \\ \tau_{l,m,n-1} \bar{\tau}_{l-1,m-2,n-1} &= a_3^2 q^{2(m-n)} \tau_{l,m-1,n} \bar{\tau}_{l-1,m-1,n-2} \\ &\quad + (1 - a_3^2 q^{2(m-n)}) \tau_{l-1,m-1,n-1} \bar{\tau}_{l,m-1,n-1}, \\ \bar{\tau}_{l,m,n} \tau_{l-1,m-2,n-2} &= a_0^{-2} q^{-2n} \tau_{l,m-1,n} \bar{\tau}_{l-1,m-1,n-2} \\ &\quad + (1 - a_0^{-2} q^{-2n}) \tau_{l-1,m-1,n-1} \bar{\tau}_{l,m-1,n-1}. \end{aligned}$$

§4. Construction of rational solutions

In this section, we construct a family of rational solutions of q -P_V. Similarly to the continuous case, we consider the fixed points of Dynkin diagram automorphism π^2 to get a seed solution. It is clear that the symmetric form of q -P_V (1.1) has a particular solution,

$$(4.1) \quad \begin{aligned} (a_0, a_1, a_2, a_3) &= (q^{-\frac{1}{2}} a, a^{-1}, q^{-\frac{1}{2}} a, a^{-1}), \\ (f_0, f_1, f_2, f_3) &= (x^{-1}, x^{-1}, x^{-1}, x^{-1}), \\ x^2 &= c, \end{aligned}$$

Applying Bäcklund transformations to the seed solution (4.1), we obtain a family of rational solutions of q -P_V.

Calculating $\tau_{l,m,n}$ and $\bar{\tau}_{l,m,n}$ from (3.2), (3.9) and (3.14), and putting as (4.1) and

$$(4.2) \quad \tau_i = \bar{\tau}_i = 1,$$

we obtain the τ -functions for the rational solutions of q -P_V. For small l, m, n , we observe that $\tau_{l,m,n}$ and $\bar{\tau}_{l,m,n}$ are factorized as the form

$$(4.3) \quad \tau_{l,m,n} = c_k U_{l,m,n}, \quad \bar{\tau}_{l,m,n} = \tilde{c}_k \bar{U}_{l,m,n}, \quad k = m - n - l.$$

It is possible to guess that $U_{l,m,n} = U_{l,m,n}(x, a)$ are some polynomials in x^{-1} , $a^{\pm 1}$ and $q^{\pm \frac{1}{2}}$, and that the factors c_k and \tilde{c}_k are determined by the recurrence relations

$$(4.4) \quad c_{k+1} \tilde{c}_{k-1} = (1 + q^{\frac{k}{2}} a^{-1} x^{-1}) c_k \tilde{c}_k, \quad \tilde{c}_{k+1} c_{k-1} = (1 + q^{-\frac{k}{2}} a x^{-1}) c_k \tilde{c}_k,$$

with the initial conditions

$$(4.5) \quad c_0 = c_1 = 1, \quad \tilde{c}_0 = \tilde{c}_1 = 1.$$

Some examples of them are listed in Appendix A.

Notice that we have

$$(4.6) \quad T_2^l T_0^l(a_0, a_1, a_2, a_3) = (q^{-\frac{1}{2}} \tilde{a}, \tilde{a}^{-1}, q^{-\frac{1}{2}} \tilde{a}, \tilde{a}^{-1}), \quad \tilde{a} = q^l a, \quad l \in \mathbb{Z},$$

under the specialization of (4.1). Comparing (4.6) with (4.1), we see that the effect of T_2 is absorbed by that of T_0^{-1} and rescaling of the parameter a . Then, we do not need to consider the translation T_2 for constructing the family of rational solutions of q -P_V, and it is possible to put

$$(4.7) \quad U_{l,m,n}(x, a) = U_{0,m,n-l}(x, q^l a).$$

Now, by (4.1), (4.3) and (4.7), we can rewrite the bilinear relations

(3.18) in terms of $U_{m,n} = U_{0,m,n}$. We have

$$\begin{aligned}
 U_{m,n}\bar{U}_{m-1,n-2} &= a^2q^{2n-1}\mu_{m-n}U_{m-1,n-1}^--\bar{U}_{m,n-1}^{++} \\
 &\quad + (1 - a^2q^{2n-1})U_{m-1,n-1}\bar{U}_{m,n-1}, \\
 \bar{U}_{m-1,n}^{--}U_{m,n-2}^{++} &= a^2\mu_{m-n}U_{m-1,n-1}^--\bar{U}_{m,n-1}^{++} \\
 &\quad + (1 - a^2)U_{m-1,n-1}\bar{U}_{m,n-1}, \\
 U_{m-1,n}^--\bar{U}_{m+1,n-1}^{++} &= a^{-2}\nu_{m-n}U_{m,n-1}^{++}\bar{U}_{m,n}^-- \\
 &\quad + (1 - a^{-2})U_{m,n-1}\bar{U}_{m,n}, \\
 \bar{U}_{m-1,n-1}U_{m+1,n} &= a^{-2}q^{2m+1}\nu_{m-n}U_{m,n-1}^{++}\bar{U}_{m,n}^-- \\
 &\quad + (1 - a^{-2}q^{2m+1})U_{m,n-1}\bar{U}_{m,n}, \\
 (4.8) \quad U_{m-1,n-1}\bar{U}_{m,n+1}^-- &= a^2q^{-2m-1}\mu_{m-n}U_{m,n}^--\bar{U}_{m-1,n} \\
 &\quad + (1 - a^2q^{-2m-1})U_{m,n}\bar{U}_{m-1,n}^--, \\
 \bar{U}_{m,n-1}U_{m-1,n+1}^-- &= a^2q^{-2(m-n)}\mu_{m-n}U_{m,n}^--\bar{U}_{m-1,n} \\
 &\quad + (1 - a^2q^{-2(m-n)})U_{m,n}\bar{U}_{m-1,n}^--, \\
 U_{m,n-1}\bar{U}_{m-2,n}^-- &= a^{-2}q^{2(m-n)}\nu_{m-n}U_{m-1,n}\bar{U}_{m-1,n-1}^-- \\
 &\quad + (1 - a^{-2}q^{2(m-n)})U_{m-1,n}\bar{U}_{m-1,n-1}^--, \\
 \bar{U}_{m,n}U_{m-2,n-1}^-- &= a^{-2}q^{-2n+1}\nu_{m-n}U_{m-1,n}\bar{U}_{m-1,n-1}^-- \\
 &\quad + (1 - a^{-2}q^{-2n+1})U_{m-1,n}\bar{U}_{m-1,n-1}^--,
 \end{aligned}$$

with the initial conditions

$$(4.9) \quad U_{-1,-1} = U_{-1,0} = U_{0,-1} = U_{0,0} = 1,$$

where we denote as

$$\begin{aligned}
 (4.10) \quad \mu_k &= (1 + q^{\frac{1}{2}(k+1)}a^{-1}x^{-1})(1 + q^{\frac{k}{2}}a^{-1}x^{-1}), \\
 \nu_k &= (1 + q^{-\frac{1}{2}(k-1)}ax^{-1})(1 + q^{-\frac{k}{2}}ax^{-1}),
 \end{aligned}$$

and

$$(4.11) \quad U_{m,n}^{\pm\pm} = U_{m,n}(x, q^{\pm 1}a).$$

Conversely, by solving the bilinear relations (4.8) with (4.9), we can construct the family of rational solutions of q -PV. Applying $T_3^m T_0^n$ to (3.10) and denoting as $T_3^m T_0^n(f_i) = f_i(x, a)$, we have the following proposition.

PROPOSITION 4.1. Let $U_{m,n} = U_{m,n}(x, a)$ ($m, n \in \mathbb{Z}$) be polynomials in x^{-1} , $a^{\pm 1}$ and $q^{\pm \frac{1}{2}}$ which satisfy the bilinear equations (4.8) with the initial conditions (4.9). Then, $f_i(x, a)$ given by

$$\begin{aligned}
 (4.12) \quad & 1 + q^{\frac{1}{2}(2n-1)} a f_0(x, a) \\
 &= (1 + q^{-\frac{1}{2}(m-n+1)} a x^{-1}) \frac{U_{m,n}(x, a) U_{m-1,n-1}(q^{\frac{1}{2}} x, q^{-1} a)}{U_{m,n-1}(q^{\frac{1}{2}} x, a) U_{m-1,n}(x, q^{-1} a)}, \\
 1 + a^{-1} f_1(x, a) &= (1 + q^{\frac{1}{2}(m-n)} a^{-1} x^{-1}) \frac{U_{m-1,n}(x, q^{-1} a) U_{m,n-1}(q^{\frac{1}{2}} x, qa)}{U_{m,n}(q^{\frac{1}{2}} x, a) U_{m-1,n-1}(x, a)}, \\
 1 + q^{-\frac{1}{2}(2m+1)} a f_2(x, a) &= (1 + q^{-\frac{1}{2}(m-n+1)} a x^{-1}) \frac{U_{m-1,n-1}(x, a) U_{m,n}(q^{\frac{1}{2}} x, q^{-1} a)}{U_{m-1,n}(q^{\frac{1}{2}} x, q^{-1} a) U_{m,n-1}(x, a)}, \\
 1 + q^{m-n} a^{-1} f_3(x, a) &= (1 + q^{\frac{1}{2}(m-n)} a^{-1} x^{-1}) \frac{U_{m,n-1}(x, a) U_{m-1,n}(q^{\frac{1}{2}} x, a)}{U_{m-1,n-1}(q^{\frac{1}{2}} x, a) U_{m,n}(x, a)},
 \end{aligned}$$

solve the q -PV (1.1) for the parameters

$$(4.13) \quad (a_0, a_1, a_2, a_3) = (q^{n-\frac{1}{2}} a, a^{-1}, q^{-m-\frac{1}{2}} a, q^{m-n} a^{-1}).$$

Moreover, the above solutions admit the other expressions as

$$\begin{aligned}
 (4.14) \quad & 1 + q^{-\frac{1}{2}(2n-1)} a^{-1} f_0(x, a) \\
 &= (1 + q^{\frac{1}{2}(m-n+1)} a^{-1} x^{-1}) \frac{U_{m,n}(q^{\frac{1}{2}} x, a) U_{m-1,n-1}(x, q^{-1} a)}{U_{m,n-1}(q^{\frac{1}{2}} x, a) U_{m-1,n}(x, q^{-1} a)}, \\
 1 + a f_1(x, a) &= (1 + q^{-\frac{1}{2}(m-n)} a x^{-1}) \frac{U_{m-1,n}(q^{\frac{1}{2}} x, q^{-1} a) U_{m,n-1}(x, qa)}{U_{m,n}(q^{\frac{1}{2}} x, a) U_{m-1,n-1}(x, a)}, \\
 1 + q^{\frac{1}{2}(2m+1)} a^{-1} f_2(x, a) &= (1 + q^{\frac{1}{2}(m-n+1)} a^{-1} x^{-1}) \frac{U_{m-1,n-1}(q^{\frac{1}{2}} x, a) U_{m,n}(x, q^{-1} a)}{U_{m-1,n}(q^{\frac{1}{2}} x, q^{-1} a) U_{m,n-1}(x, a)}, \\
 1 + q^{-m+n} a f_3(x, a) &= (1 + q^{-\frac{1}{2}(m-n)} a x^{-1}) \frac{U_{m,n-1}(q^{\frac{1}{2}} x, a) U_{m-1,n}(x, a)}{U_{m-1,n-1}(q^{\frac{1}{2}} x, a) U_{m,n}(x, a)}.
 \end{aligned}$$

§5. Proof of Theorem 2.3

In this section, we give the proof for Theorem 2.3.

PROPOSITION 5.1. *We have*

$$(5.1) \quad U_{m,n} = (-1)^{\binom{n+1}{2}} \kappa_m \kappa_n S_{m,n},$$

where $S_{m,n} = S_{m,n}(x, a)$ is defined in Theorem 2.3 and κ_n is the factor determined by

$$(5.2) \quad \kappa_{n+1} \bar{\kappa}_{n-1} = q^{-\frac{1}{4}(2n+1)} x^{-1} (1 - q^{2n+1}) \kappa_n \bar{\kappa}_n, \quad \kappa_0 = \kappa_{-1} = 1.$$

We notice that κ_n for $n \geq 0$ is expressed as

$$(5.3) \quad \kappa_n = q^{-\binom{n+1}{3} - \frac{1}{4} \binom{n+1}{2}} x^{-\binom{n+1}{2}} \prod_{k=1}^n \prod_{j=1}^k (1 - q^{2j-1}).$$

By substituting (5.1) into (4.12), we find that Theorem 2.3 is a direct consequence of Propositions 4.1 and 5.1. Taking (5.1) and (2.8) into account, we obtain the bilinear relations for $R_{m,n}^{(b)}$.

PROPOSITION 5.2. *The following bilinear relations hold:*

$$(5.4) \quad \begin{aligned} & -q^{\frac{1}{4}(2m-2n-1)} x(1 - q^{2n+1}) R_{m,n+1}^+ \bar{R}_{m-1,n-1} \\ & \quad = b^2 q^{m+n+1} \mu^+ R_{m-1,n}^- \bar{R}_{m,n}^{++} + x^2 (1 - b^2 q^{m+n+1}) R_{m-1,n}^+ \bar{R}_{m,n}, \\ & -q^{\frac{1}{4}(2m-6n-3)} x(1 - q^{2n+1}) \bar{R}_{m-1,n+1} R_{m,n-1}^+ \\ & \quad = b^2 q^{m-n} \mu^+ R_{m-1,n}^- \bar{R}_{m,n}^{++} + x^2 (1 - b^2 q^{m-n}) R_{m-1,n}^+ \bar{R}_{m,n}, \\ & q^{\frac{1}{4}(-6m+2n-3)} x(1 - q^{2m+1}) R_{m-1,n}^- \bar{R}_{m+1,n-1} \\ & \quad = b^{-2} q^{-m+n} \nu R_{m,n-1}^+ \bar{R}_{m,n}^{--} + x^2 (1 - b^{-2} q^{-m+n}) R_{m,n-1}^- \bar{R}_{m,n}, \\ & q^{\frac{1}{4}(-2m+2n-1)} x(1 - q^{2m+1}) \bar{R}_{m-1,n-1} R_{m+1,n}^- \\ & \quad = b^{-2} q^{m+n+1} \nu R_{m,n-1}^+ \bar{R}_{m,n}^{--} + x^2 (1 - b^{-2} q^{m+n+1}) R_{m,n-1}^- \bar{R}_{m,n}, \\ & -q^{-\frac{1}{4}(2m+6n+3)} x(1 - q^{2n+1}) R_{m-1,n-1} \bar{R}_{m,n+1}^- \\ & \quad = b^2 q^{-m-n-1} \mu R_{m,n}^{--} \bar{R}_{m-1,n}^+ + x^2 (1 - b^2 q^{-m-n-1}) R_{m,n} \bar{R}_{m-1,n}^-, \\ & -q^{-\frac{1}{4}(2m+2n+1)} x(1 - q^{2n+1}) \bar{R}_{m,n-1}^- R_{m-1,n+1} \end{aligned}$$

$$\begin{aligned}
 &= b^2 q^{-m+n} \mu R_{m,n}^- \bar{R}_{m-1,n}^+ + x^2 (1 - b^2 q^{-m+n}) R_{m,n} \bar{R}_{m-1,n}^-, \\
 q^{-\frac{1}{4}(2m+2n+1)} x (1 - q^{2m+1}) R_{m+1,n-1}^- \bar{R}_{m-1,n}^- \\
 &= b^{-2} q^{m-n+2} \nu^- R_{m,n} \bar{R}_{m,n-1}^- + x^2 (1 - b^{-2} q^{m-n+2}) R_{m,n}^- \bar{R}_{m,n-1}^-, \\
 q^{-\frac{1}{4}(6m+2n+3)} x (1 - q^{2m+1}) \bar{R}_{m+1,n}^- R_{m-1,n-1}^- \\
 &= b^{-2} q^{-m-n+1} \nu^- R_{m,n} \bar{R}_{m,n-1}^- + x^2 (1 - b^{-2} q^{-m-n+1}) R_{m,n}^- \bar{R}_{m,n-1}^-,
 \end{aligned}$$

with

$$(5.5) \quad \mu = (x + b^{-1})(x + q^{\frac{1}{2}} b^{-1}), \quad \nu = (x + b)(x + q^{-\frac{1}{2}} b),$$

where we denote as

$$(5.6) \quad X^{\overbrace{\pm \cdots \pm}^j} = X^{\overbrace{\pm \cdots \pm}^j}(b) = X(q^{\pm \frac{j}{2}} b).$$

From the above discussion, now the proof of Theorem 2.3 is reduced to that of Proposition 5.2.

It is possible to reduce the number of bilinear relations to be proved in (5.4) by the following symmetry of $R_{m,n}^{(b)}(y|q)$.

LEMMA 5.3. *We have the relations for $m, n \in \mathbb{Z}_{\geq 0}$*

$$(5.7) \quad \begin{aligned} R_{n,m}^{(b^{-1})}(y|q^{-1}) &= R_{m,n}^{(b)}(y|q), \\ R_{n,m}^{(b^{-1})} &= (-1)^{m(m+1)/2+n(n+1)/2} R_{m,n}^{(b)}. \end{aligned}$$

Proof. From (2.7), it is easy to see that

$$(5.8) \quad q_k^{(b)}(y|q) = p_k^{(b^{-1})}(y|q^{-1}),$$

which leads to the first relation of Lemma 5.3. To verify the second relation, we introduce polynomials $\tilde{q}_k^{(b)} = \tilde{q}_k^{(b)}(y|q)$ by

$$(5.9) \quad \sum_{k=0}^{\infty} \tilde{q}_k^{(b)} \lambda^k = \frac{(-q^{\frac{1}{4}} b^{-1} \lambda, -q^{\frac{3}{4}} b^{-1} \lambda; q)_{\infty}}{(q^{\frac{1}{4}} x \lambda, q^{\frac{3}{4}} x^{-1} \lambda; q)_{\infty}}, \quad \tilde{q}_k^{(b)} = 0 \text{ for } k < 0.$$

Comparing the generating functions, we see that each \tilde{q}_k is a linear combination of q_j , $j = k, k - 2, k - 4, \dots$. Therefore we can express $R_{m,n}^{(b)}$ for

$m, n \in \mathbb{Z}_{\geq 0}$ in terms of p_k and \tilde{q}_k as

$$(5.10) \quad R_{m,n}^{(b)} = \begin{vmatrix} \tilde{q}_1^{(b)} & \tilde{q}_0^{(b)} & \cdots & \tilde{q}_{-m+2}^{(b)} & \tilde{q}_{-m+1}^{(b)} & \cdots & \tilde{q}_{-m-n+3}^{(b)} & \tilde{q}_{-m-n+2}^{(b)} \\ \tilde{q}_3^{(b)} & \tilde{q}_2^{(b)} & \cdots & \tilde{q}_{-m+4}^{(b)} & \tilde{q}_{-m+3}^{(b)} & \cdots & \tilde{q}_{-m-n+5}^{(b)} & \tilde{q}_{-m-n+4}^{(b)} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \tilde{q}_{2m-1}^{(b)} & \tilde{q}_{2m-2}^{(b)} & \cdots & \tilde{q}_m^{(b)} & \tilde{q}_{m-1}^{(b)} & \cdots & \tilde{q}_{m-n+1}^{(b)} & \tilde{q}_{m-n}^{(b)} \\ p_{n-m}^{(b)} & p_{n-m+1}^{(b)} & \cdots & p_{n-1}^{(b)} & p_n^{(b)} & \cdots & p_{2n-2}^{(b)} & p_{2n-1}^{(b)} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots \\ p_{-n-m+4}^{(b)} & p_{-n-m+5}^{(b)} & \cdots & p_{-n+3}^{(b)} & p_{-n+4}^{(b)} & \cdots & p_2^{(b)} & p_3^{(b)} \\ p_{-n-m+2}^{(b)} & p_{-n-m+3}^{(b)} & \cdots & p_{-n+1}^{(b)} & p_{-n+2}^{(b)} & \cdots & p_0^{(b)} & p_1^{(b)} \end{vmatrix}.$$

Noticing that \tilde{q}_k and p_k are related as

$$(5.11) \quad \tilde{q}_k^{(b)}(y|q) = (-1)^k p_k^{(b^{-1})}(y|q),$$

we obtain the second relation of Lemma 5.3. □

From the symmetries of $R_{m,n}^{(b)}(y|q)$ described by (2.4) and Lemma 5.3, it is sufficient to prove the first two relations in (5.4) for $m, n \in \mathbb{Z}_{\geq 0}$, which are equivalent to

$$(5.12) \quad -q^{\frac{1}{4}(2m-2n-1)} R_{m,n+1}^+ \bar{R}_{m-1,n-1} + q^{\frac{1}{4}(2m+2n+1)} \bar{R}_{m-1,n+1} R_{m,n-1}^+ = x R_{m-1,n}^+ \bar{R}_{m,n},$$

$$(5.13) \quad -q^{\frac{1}{4}(2m-6n-3)} x(1 - q^{2n+1}) \bar{R}_{m-1,n+1} R_{m,n-1}^+ = b^2 q^{m-n} \mu^+ R_{m-1,n}^- \bar{R}_{m,n}^{++} + x^2(1 - b^2 q^{m-n}) R_{m-1,n}^+ \bar{R}_{m,n},$$

In the following, we show that these bilinear relations are reduced to Jacobi’s identity of determinants. Let D be an $(m + n + 1) \times (m + n + 1)$ determinant and $D \begin{bmatrix} i_1 & i_2 & \cdots & i_k \\ j_1 & j_2 & \cdots & j_k \end{bmatrix}$ the minor which are obtained by deleting the rows with indices i_1, \dots, i_k and the columns with indices j_1, \dots, j_k . Then we have Jacobi’s identity

$$(5.14) \quad D \cdot D \begin{bmatrix} m & m + 1 \\ 1 & m + n + 1 \end{bmatrix} = D \begin{bmatrix} m \\ 1 \end{bmatrix} D \begin{bmatrix} m + 1 \\ m + n + 1 \end{bmatrix} - D \begin{bmatrix} m + 1 \\ 1 \end{bmatrix} D \begin{bmatrix} m \\ m + n + 1 \end{bmatrix}.$$

We first choose proper determinants as D (D itself should be expressed in terms of $R_{m,n}^{(b)}$). Secondly, we construct such formulas that express the minor

determinants by $R_{m,n}^{(b)}$. Then, Jacobi's identity yields bilinear equations for $R_{m,n}^{(b)}$ which are nothing but (5.12) and (5.13).

We have the following lemmas.

LEMMA 5.4. *We set*

$$(5.15) \quad D \equiv \begin{vmatrix} q^{-\frac{m+n-2}{2}} q^{\frac{1}{4}} x^{-1} q_1^+ & \bar{q}_1 & \cdots & \bar{q}_{-m-n+3} & \bar{q}_{-m-n+2} \\ q^{-\frac{m+n-4}{2}} q^{\frac{1}{4}} x^{-1} q_3^+ & \bar{q}_3 & \cdots & \bar{q}_{-m-n+5} & \bar{q}_{-m-n+4} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ q^{-\frac{n-m}{2}} q^{\frac{1}{4}} x^{-1} q_{2m-1}^+ & \bar{q}_{2m-1} & \cdots & \bar{q}_{m-n+1} & \bar{q}_{m-n} \\ q^{-n} p_{n-m+1}^+ & \bar{p}_{n-m+2} & \cdots & \bar{p}_{2n} & \bar{p}_{2n+1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ q^{-1} p_{-n-m+3}^+ & \bar{p}_{-n-m+4} & \cdots & \bar{p}_2 & \bar{p}_3 \\ p_{-n-m+1}^+ & \bar{p}_{-n-m+2} & \cdots & \bar{p}_0 & \bar{p}_1 \end{vmatrix}.$$

Then, we have

$$(5.16) \quad \begin{aligned} D &= q^{\frac{1}{4}m^2 - \frac{1}{4}n(n+5)} x^{-m} R_{m,n+1}^+, \\ D \begin{bmatrix} cm \\ 1 \end{bmatrix} &= \bar{R}_{m-1,n+1}, \quad D \begin{bmatrix} cm+1 \\ 1 \end{bmatrix} = \bar{R}_{m,n}, \\ D \begin{bmatrix} m \\ m+n+1 \end{bmatrix} &= q^{\frac{1}{4}(m-1)^2 - \frac{1}{4}(n-1)(n+4) - 1} x^{-m+1} R_{m-1,n}^+, \\ D \begin{bmatrix} m+1 \\ m+n+1 \end{bmatrix} &= q^{\frac{1}{4}m^2 - \frac{1}{4}(n-2)(n+3) - 1} x^{-m} R_{m,n-1}^+, \\ D \begin{bmatrix} ccm & m+1 \\ 1 & m+n+1 \end{bmatrix} &= \bar{R}_{m-1,n-1}. \end{aligned}$$

LEMMA 5.5. *Define $P_{j,k}^{[-m-n+j]}$ and $Q_{j,k}^{[-m-n+j]}$ by*

$$(5.17) \quad \begin{aligned} P_{j,k}^{[-m-n+j]} &= \prod_{i=1}^{m+n-j} \left(q^{-\frac{1}{4} - \frac{i}{2}} x \right) q^{\frac{1}{2}(m+n-j)(k - \frac{1}{2}(m+n-j) + \frac{1}{2})} p_k^{[-m-n+j]}, \\ Q_{j,k}^{[-m-n+j]} &= q^{-\frac{1}{2}(m+n-j)k} q_k^{[-m-n+j]}, \end{aligned}$$

where we denote

$$(5.18) \quad X^{[j]} = X^{[j]}(x, b) = X\left(q^{\frac{j}{2}} x, q^{\frac{j}{2}} b\right).$$

Then, setting

$$(5.19) \quad D \equiv \begin{vmatrix} \tilde{Q}_{0,1}^{[-m-n]++} & Q_{1,1}^{[-m-n+1]} & \dots & Q_{m+n-1,-m-n+3}^{[-1]} & Q_{m+n,-m-n+2}^{[0]} \\ \tilde{Q}_{0,3}^{[-m-n]++} & Q_{1,3}^{[-m-n+1]} & \dots & Q_{m+n-1,-m-n+5}^{[-1]} & Q_{m+n,-m-n+4}^{[0]} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \tilde{Q}_{0,2m-1}^{[-m-n]++} & Q_{1,2m-1}^{[-m-n+1]} & \dots & Q_{m+n-1,m-n+1}^{[-1]} & Q_{m+n,m-n}^{[0]} \\ \hat{P}_{0,2n}^{[-m-n]++} & P_{1,2n+1}^{[-m-n+1]} & \dots & P_{m+n-1,2n+1}^{[-1]} & P_{m+n,2n+1}^{[0]} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \hat{P}_{0,2}^{[-m-n]++} & P_{1,3}^{[-m-n+1]} & \dots & P_{m+n-1,3}^{[-1]} & P_{m+n,3}^{[0]} \\ \hat{P}_{0,0}^{[-m-n]++} & P_{1,1}^{[-m-n+1]} & \dots & P_{m+n-1,1}^{[-1]} & P_{m+n,1}^{[0]} \end{vmatrix},$$

where

$$(5.20) \quad \begin{aligned} \hat{P}_{0,2k}^{[-m-n]++} &= -\frac{P_{0,2k}^{[-m-n]++}}{1 - q^{2k+1}}, \\ \tilde{Q}_{0,2k-1}^{[-m-n]++} &= \frac{Q_{0,2k-1}^{[-m-n]++}}{q^{m+n+1-2k}(1 - q^{-m-n+2k}b^2)}, \end{aligned}$$

we have

$$(5.21) \quad \begin{aligned} D &= (-1)^{n+1} \frac{\prod_{j=1}^{m+n} \mu^{[-m-n+j]}}{(q^{-\frac{1}{4}}b^{-2}x)^{m+n} \prod_{k=0}^n (1-q^{2k+1}) \prod_{i=1}^m q^{m+n+1-2i}(1-q^{-m-n+2i}b^2)} R_{m,n}^{++}, \\ D \begin{bmatrix} m \\ 1 \end{bmatrix} &= R_{m-1,n+1}, \quad D \begin{bmatrix} m+1 \\ 1 \end{bmatrix} = R_{m,n}, \\ D \begin{bmatrix} m \\ m+n+1 \end{bmatrix} &= (-1)^{n+1} x^{n+1} \\ &\quad \times \frac{q^{-\frac{1}{4}m^2 + \frac{1}{4}n^2 + \frac{1}{4}m - \frac{1}{2}n - \frac{3}{4}} \prod_{j=1}^{m+n-1} \mu^{[-m-n+j]}}{(q^{-\frac{1}{4}}b^{-2}x)^{m+n-1} \prod_{k=0}^n (1-q^{2k+1}) \prod_{i=1}^{m-1} q^{m+n+1-2i}(1-q^{-m-n+2i}b^2)} R_{m-1,n}^+, \\ D \begin{bmatrix} m+1 \\ m+n+1 \end{bmatrix} &= (-1)^n x^n \\ &\quad \times \frac{q^{-\frac{1}{4}m^2 + \frac{1}{4}n^2 - \frac{1}{4}m - n} \prod_{j=1}^{m+n-1} \mu^{[-m-n+j]}}{(q^{-\frac{1}{4}}b^{-2}x)^{m+n-1} \prod_{k=0}^{n-1} (1-q^{2k+1}) \prod_{i=1}^m q^{m+n+1-2i}(1-q^{-m-n+2i}b^2)} R_{m,n-1}^+, \end{aligned}$$

$$D \begin{bmatrix} m & m + 1 \\ 1 & m + n + 1 \end{bmatrix} = q^{-\frac{1}{4}m^2 + \frac{1}{4}n^2 + \frac{1}{4}m - \frac{1}{2}n} x^n \underline{R}_{m-1, n}^-.$$

It is easy to see that the bilinear relations (5.12) and (5.13) follow immediately from Jacobi’s identity (5.14) by using Lemmas 5.4 and 5.5, respectively. We give the proof of Lemmas 5.4 and 5.5 in Appendix B. This completes the proof of our main result Theorem 2.3.

§6. Remarks

The q -P_V (1.1) admits the ultra-discrete limit [14]. The limiting procedure is the same as the case of q -P_{IV} [4] and preserves the symmetry of the extended affine Weyl group of type $A_3^{(1)}$. Moreover, it is observed that $U_{m,n} = U_{m,n}(x, a)$ are polynomials in $x^{-1}, a^{\pm 1}$ and $q^{\pm \frac{1}{2}}$ with positive coefficients. Then, the rational solutions of q -P_V (1.1) in Theorem 2.3 are thought to survive after taking the ultra-discrete limit.

It is known that the special polynomials associated with the rational solutions of the Painlevé equations possess the mysterious combinatorial properties [16], [9], [15]. It is interesting problem to investigate whether the polynomials $U_{m,n}$ admit such properties.

In [4], it has been shown that the q -P_{IV} coincides with Sakai’s Mul.6 system [13]. As mentioned in Section 1, the q -P_V (1.1) has $\widetilde{W}(A_1^{(1)} \times A_3^{(1)})$ symmetry by the original construction. On the other hand, Sakai’s Mul.5 system [13], which should be also regarded as a q -analogue of the Painlevé V equation, admits the symmetry of $\widetilde{W}(A_4^{(1)})$. It might be an important problem to study the relationship between the equation (1.1) and Sakai’s Mul.5 system.

Acknowledgment. The author would like to thank Prof. M. Noumi, Prof. Y. Yamada and Prof. K. Kajiwara for fruitful discussions.

A. Table of c_k, \tilde{c}_k and $U_{m,n}$

The polynomials $U_{m,n}(x, a)$.

$$\begin{aligned} U_{0,0} &= 1, \\ U_{1,0} &= 1 + q^{\frac{1}{2}} x^{-2} + a^{-1} q^{\frac{1}{2}} (1 + q^{\frac{1}{2}}) x^{-1}, \\ U_{2,0} &= 1 + q^{\frac{3}{2}} x^{-6} + a^{-1} (1 + q^{\frac{1}{2}}) (1 + q + q^2) (x^{-1} + qx^{-5}) \\ &\quad + q^{-\frac{1}{2}} (1 + q + q^2) \left[1 + a^{-2} q^{\frac{3}{2}} (1 + q^{\frac{1}{2}})^2 \right] (x^{-2} + q^{\frac{1}{2}} x^{-4}) \end{aligned}$$

$$+ a^{-1}q^{\frac{1}{2}}(1 + q^{\frac{3}{2}})\left[2(1 + q^{\frac{1}{2}} + q) + a^{-2}q^2(1 + q^{\frac{1}{2}})^2\right]x^{-3},$$

$$U_{0,1} = 1 + aq^{\frac{1}{2}}(1 + q^{\frac{1}{2}})x^{-1} + q^{\frac{1}{2}}x^{-2},$$

$$U_{0,2} = 1 + q^{\frac{3}{2}}x^{-6} + a(1 + q^{\frac{1}{2}})(1 + q + q^2)(x^{-1} + qx^{-5}) \\ + q^{-\frac{1}{2}}(1 + q + q^2)\left[1 + a^2q^{\frac{3}{2}}(1 + q^{\frac{1}{2}})^2\right](x^{-2} + q^{\frac{1}{2}}x^{-4}) \\ + aq^{\frac{1}{2}}(1 + q^{\frac{3}{2}})\left[2(1 + q^{\frac{1}{2}} + q) + a^2q^2(1 + q^{\frac{1}{2}})^2\right]x^{-3},$$

$$U_{1,1} = 1 + qx^{-4} + (1 + q^{\frac{1}{2}})(a + a^{-1})(x^{-1} + q^{\frac{1}{2}}x^{-3}) \\ + q^{-\frac{1}{2}}(1 + q)(1 + q^{\frac{1}{2}} + q)x^{-2},$$

$$U_{1,2} = 1 + q^2x^{-8} + a^{-1}q^{-\frac{1}{2}}(1 + q^{\frac{1}{2}})\left[1 + a^2(1 + q + q^2)\right](x^{-1} + q^{\frac{3}{2}}x^{-7}) \\ + q^{-\frac{3}{2}}\left[a^2q^{\frac{3}{2}}(1 + q^{\frac{1}{2}})^2(1 + q + q^2) \\ + (1 + q^{\frac{1}{2}} + q + q^2)(1 + q + q^{\frac{3}{2}} + q^2)\right](x^{-2} + qx^{-6}) \\ + a^{-1}q^{-1}(1 + q^{\frac{3}{2}})\left[a^4q^2(1 + q^{\frac{1}{2}})^2 + a^2(1 + q + q^2)(2 + 3q^{\frac{1}{2}} + 2q) \\ + (1 + q^{\frac{1}{2}} + q)\right](x^{-3} + q^{\frac{1}{2}}x^{-5}) \\ + q^{-1}(1 + q^{\frac{1}{2}} + q + q^{\frac{3}{2}} + q^2) \\ \times \left[a^2q^{\frac{1}{2}}(1 + q^{\frac{1}{2}})(1 + q)(1 + q^{\frac{3}{2}}) + 2(1 + q + q^2)\right]x^{-4},$$

$$U_{2,1} = 1 + q^2x^{-8} + a^{-1}q^{-\frac{1}{2}}(1 + q^{\frac{1}{2}})\left[a^2 + (1 + q + q^2)\right](x^{-1} + q^{\frac{3}{2}}x^{-7}) \\ + a^{-2}q^{-\frac{3}{2}}\left[a^2(1 + q^{\frac{1}{2}} + q + q^2)(1 + q + q^{\frac{3}{2}} + q^2) \\ + q^{\frac{3}{2}}(1 + q^{\frac{1}{2}})^2(1 + q + q^2)\right](x^{-2} + qx^{-6}) \\ + a^{-3}q^{-1}(1 + q^{\frac{3}{2}})\left[a^4(1 + q^{\frac{1}{2}} + q) + a^2(1 + q + q^2)(2 + 3q^{\frac{1}{2}} + 2q) \\ + q^2(1 + q^{\frac{1}{2}})^2\right](x^{-3} + q^{\frac{1}{2}}x^{-5}) \\ + a^{-2}q^{-1}(1 + q^{\frac{1}{2}} + q + q^{\frac{3}{2}} + q^2) \\ \times \left[2a^2(1 + q + q^2) + q^{\frac{1}{2}}(1 + q^{\frac{1}{2}})(1 + q)(1 + q^{\frac{3}{2}})\right]x^{-4}.$$

The factor c_k and \tilde{c}_k .

$$\begin{aligned}
 c_0 &= 1, & c_1 &= 1, \\
 c_2 &= 1 + q^{\frac{1}{2}} a^{-1} x^{-1}, \\
 c_3 &= (1 + q^{\frac{1}{2}} a^{-1} x^{-1})(1 + qa^{-1}x^{-1})(1 + q^{-\frac{1}{2}} ax^{-1}), \\
 c_4 &= (1 + q^{\frac{1}{2}} a^{-1} x^{-1})^2(1 + qa^{-1}x^{-1})(1 + q^{\frac{3}{2}} a^{-1} x^{-1})(1 + q^{-\frac{1}{2}} ax^{-1}) \\
 &\quad \times (1 + q^{-1} ax^{-1}), \\
 c_5 &= (1 + q^{\frac{1}{2}} a^{-1} x^{-1})^2(1 + qa^{-1}x^{-1})^2(1 + q^{\frac{3}{2}} a^{-1} x^{-1})(1 + q^2 a^{-1} x^{-1}) \\
 &\quad \times (1 + q^{-\frac{1}{2}} ax^{-1})^2(1 + q^{-1} ax^{-1})(1 + q^{-\frac{3}{2}} ax^{-1}), \\
 \\
 \tilde{c}_0 &= 1, & \tilde{c}_1 &= 1, \\
 \tilde{c}_2 &= 1 + q^{-\frac{1}{2}} ax^{-1}, \\
 \tilde{c}_3 &= (1 + q^{-\frac{1}{2}} ax^{-1})(1 + q^{-1} ax^{-1})(1 + q^{\frac{1}{2}} a^{-1} x^{-1}), \\
 \tilde{c}_4 &= (1 + q^{-\frac{1}{2}} ax^{-1})^2(1 + q^{-1} ax^{-1})(1 + q^{-\frac{3}{2}} ax^{-1})(1 + q^{\frac{1}{2}} a^{-1} x^{-1}) \\
 &\quad \times (1 + qa^{-1}x^{-1}), \\
 \tilde{c}_5 &= (1 + q^{-\frac{1}{2}} ax^{-1})^2(1 + q^{-1} ax^{-1})^2(1 + q^{-\frac{3}{2}} ax^{-1})(1 + q^{-1} ax^{-2}) \\
 &\quad \times (1 + q^{\frac{1}{2}} a^{-1} x^{-1})^2(1 + qa^{-1}x^{-1})(1 + q^{\frac{3}{2}} a^{-1} x^{-1}).
 \end{aligned}$$

B. Proof of Lemmas 5.4 and 5.5

We first note that the following contiguity relations hold,

$$(B.1) \quad p_k^+ - q^{\frac{k}{2}} \bar{p}_k = -q^{\frac{1}{4}} x p_{k-1}^+, \quad q_k^+ - q^{-\frac{k}{2}} \bar{q}_k = -q^{-\frac{1}{4}} x^{-1} q_{k-1}^+,$$

$$(B.2) \quad p_k - q^{\frac{k}{2}} \underline{p}_k^- = -q^{\frac{3}{4}} x^{-1} p_{k-1}, \quad q_k - q^{-\frac{k}{2}} \underline{q}_k^- = -q^{-\frac{3}{4}} x q_{k-1},$$

and

$$\begin{aligned}
 (B.3) \quad &(1 - q^{k+1})p_{k+1} = q^{\frac{k}{2} + \frac{1}{4}} bx(b - b^{-1})\underline{p}_k^+ - q^{\frac{1}{4}} b^2 x^{-1} \mu p_k^{++}, \\
 &(1 - q^{k+1} b^2)q_{k+1} = q^{\frac{1}{2}(k+1)} b(b^{-1} - b)\underline{q}_{k+1}^+ + q^{k + \frac{1}{4}} b^2 x^{-1} \mu q_k^{++},
 \end{aligned}$$

which are easily derived from (2.1).

Let us prove Lemma 5.4. Noticing that $p_1 = 1$ and $p_k = 0$ for $k < 0$, we see that $R_{m,n}$ can be rewritten as

$$(B.4) \quad R_{m,n} = \begin{vmatrix} q_1 & q_0 & \cdots & q_{-m-n+3} & q_{-m-n+2} & q_{-m-n+1} \\ q_3 & q_2 & \cdots & q_{-m-n+5} & q_{-m-n+4} & q_{-m-n+3} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ q_{2m-1} & q_{2m-2} & \cdots & q_{m-n+1} & q_{m-n} & q_{m-n-1} \\ p_{n-m} & p_{n-m+1} & \cdots & p_{2n-2} & p_{2n-1} & p_{2n} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ p_{-n-m+4} & p_{-n-m+5} & \cdots & p_2 & p_3 & p_4 \\ p_{-n-m+2} & p_{-n-m+3} & \cdots & p_0 & p_1 & p_2 \\ p_{-n-m} & p_{-n-m+1} & \cdots & p_{-2} & p_{-1} & p_0 \end{vmatrix}.$$

Adding the $(j-1)$ -th column multiplied by $q^{\frac{1}{4}}x$ to the j -th column of $R_{m,n}^+$ for $j = m+n, m+n-1, \dots, 2$ and using (B.1), we get

$$(B.5) \quad R_{m,n}^+ = q^{-\frac{1}{4}m^2 + \frac{1}{4}(n-1)(n+4)} x^m \times \begin{vmatrix} q^{-\frac{m+n-3}{2}} q^{\frac{1}{4}} x^{-1} q_1^+ & \bar{q}_1 & \cdots & \bar{q}_{-m-n+4} & \bar{q}_{-m-n+3} \\ q^{-\frac{m+n-5}{2}} q^{\frac{1}{4}} x^{-1} q_3^+ & \bar{q}_3 & \cdots & \bar{q}_{-m-n+6} & \bar{q}_{-m-n+5} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ q^{-\frac{n-m-1}{2}} q^{\frac{1}{4}} x^{-1} q_{2m-1}^+ & \bar{q}_{2m-1} & \cdots & \bar{q}_{-n+2} & \bar{q}_{m-n+1} \\ q^{-n+1} p_{n-m}^+ & \bar{p}_{n-m+1} & \cdots & \bar{p}_{2n-2} & \bar{p}_{2n-1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ q^{-1} p_{-n-m+4}^+ & \bar{p}_{-n-m+5} & \cdots & \bar{p}_2 & \bar{p}_3 \\ p_{-n-m+2}^+ & \bar{p}_{-n-m+3} & \cdots & \bar{p}_0 & \bar{p}_1 \end{vmatrix}.$$

From (B.4) and (B.5), we obtain Lemma 5.4.

We next prove Lemma 5.5. Adding the $(i+1)$ -th column multiplied by $q^{\frac{1}{4}-(m+n-j)}x$ to the i -th column of $R_{m,n}$ for $i = 1, 2, \dots, j, j = m+n-1, m+n-2, \dots, 1$ and using (B.2), we get

$$(B.6) \quad R_{m,n} = \begin{vmatrix} Q_{1,1}^{[-m-n+1]} & Q_{2,0}^{[-m-n+2]} & \cdots & Q_{m+n-1,-m-n+3}^{[-1]} & Q_{m+n,-m-n+2}^{[0]} \\ Q_{1,3}^{[-m-n+1]} & Q_{2,2}^{[-m-n+2]} & \cdots & Q_{m+n-1,-m-n+5}^{[-1]} & Q_{m+n,-m-n+4}^{[0]} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ Q_{1,2m-1}^{[-m-n+1]} & Q_{2,2m-2}^{[-m-n+2]} & \cdots & Q_{m+n-1,m-n+1}^{[-1]} & Q_{m+n,m-n}^{[0]} \\ P_{1,2n-1}^{[-m-n+1]} & P_{2,2n-1}^{[-m-n+2]} & \cdots & P_{m+n-1,2n-1}^{[-1]} & P_{m+n,2n-1}^{[0]} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ P_{1,3}^{[-m-n+1]} & P_{2,3}^{[-m-n+2]} & \cdots & P_{m+n-1,3}^{[-1]} & P_{m+n,3}^{[0]} \\ P_{1,1}^{[-m-n+1]} & P_{2,1}^{[-m-n+2]} & \cdots & P_{m+n-1,1}^{[-1]} & P_{m+n,1}^{[0]} \end{vmatrix}.$$

Noticing that $p_1 = 1$ and $p_k = 0$ for $k < 0$, we see that $R_{m,n}$ can be

rewritten as

$$(B.7) \quad R_{m,n} = \begin{pmatrix} Q_{0,1}^{[-m-n]} & Q_{1,0}^{[-m-n+1]} & \dots & Q_{m+n-1,-m-n+2}^{[-1]} & Q_{m+n,-m-n+1}^{[0]} \\ Q_{0,3}^{[-m-n]} & Q_{1,2}^{[-m-n+1]} & \dots & Q_{m+n-1,-m-n+4}^{[-1]} & Q_{n+n,-m-n+3}^{[0]} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ Q_{0,2m-1}^{[-m-n]} & Q_{1,2m-2}^{[-m-n+1]} & \dots & Q_{m+n-1,m-n}^{[-1]} & Q_{m+n,m-n-1}^{[0]} \\ P_{0,2n}^{[-m-n]} & P_{1,2n}^{[-m-n+1]} & \dots & P_{m+n-1,2n}^{[-1]} & P_{m+n,2n}^{[0]} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ P_{0,2}^{[-m-n]} & P_{1,2}^{[-m-n+1]} & \dots & P_{m+n-1,2}^{[-1]} & P_{m+n,2}^{[0]} \\ P_{0,0}^{[-m-n]} & P_{1,0}^{[-m-n+1]} & \dots & P_{m+n-1,0}^{[-1]} & P_{m+n,0}^{[0]} \end{pmatrix}.$$

Then, adding the j -th column multiplied $\frac{q^{\frac{3}{4}+m+n-j}b^{-2}x(1-q^{-m-n+j}b^2)}{\mu^{[-m-n+j]}}$ to $(j+1)$ -th column of $R_{m,n}^{++}$ for $j = m+n, m+n-1, \dots, 1$ and using (B.3), we obtain

$$(B.8) \quad R_{m,n}^{++} = (-1)^{n+1} \frac{(q^{-\frac{1}{4}}b^{-2}x)^{m+n} \prod_{k=0}^n (1-q^{2k+1}) \prod_{i=1}^m q^{m+n+1-2i}(1-q^{-m-n+2i}b^2)}{\prod_{j=1}^{m+n} \mu^{[-m-n+j]}} \times \begin{pmatrix} \tilde{Q}_{0,1}^{[-m-n]++} & Q_{1,1}^{[-m-n+1]} & \dots & Q_{m+n-1,-m-n+3}^{[-1]} & Q_{m+n,-m-n+2}^{[0]} \\ \tilde{Q}_{0,3}^{[-m-n]++} & Q_{1,3}^{[-m-n+1]} & \dots & Q_{m+n-1,-m-n+5}^{[-1]} & Q_{n+n,-m-n+4}^{[0]} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \tilde{Q}_{0,2m-1}^{[-m-n]++} & Q_{1,2m-1}^{[-m-n+1]} & \dots & Q_{m+n-1,m-n+1}^{[-1]} & Q_{m+n,m-n}^{[0]} \\ \tilde{P}_{0,2n}^{[-m-n]++} & P_{1,2n+1}^{[-m-n+1]} & \dots & P_{m+n-1,2n+1}^{[-1]} & P_{m+n,2n+1}^{[0]} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \tilde{P}_{0,2}^{[-m-n]++} & P_{1,3}^{[-m-n+1]} & \dots & P_{m+n-1,3}^{[-1]} & P_{m+n,3}^{[0]} \\ \tilde{P}_{0,0}^{[-m-n]++} & P_{1,1}^{[-m-n+1]} & \dots & P_{m+n-1,1}^{[-1]} & P_{m+n,1}^{[0]} \end{pmatrix},$$

where we use the relations

$$(B.9) \quad \begin{aligned} \underline{P}_{j,k}^{[-m-n+j+1]-} &= q^{-\frac{k}{2} + \frac{1}{4} + \frac{1}{2}(m+n-j)} x^{-1} P_{j,k}^{[-m-n+j]}, \\ \underline{Q}_{j,k}^{[-m-n+j+1]-} &= q^{\frac{k}{2}} Q_{j,k}^{[-m-n+j]}. \end{aligned}$$

Lemma 5.5 follows from (B.6), (B.7) and (B.8).

C. Continuum limit to the rational solutions of P_V

We consider the continuum limit of the rational solutions of q - P_V to those of the Painlevé V equation. In the previous paper [8], we have presented a determinant formula for the rational solutions of P_V .

PROPOSITION C.1. Let $\hat{p}_k^{(r)} = \hat{p}_k^{(r)}(z)$ and $\hat{q}_k^{(r)} = \hat{q}_k^{(r)}(z)$ be polynomials defined by

$$(C.1) \quad \hat{p}_k^{(r)}(z) = L_k^{(r-1)}(z), \quad \hat{q}_k^{(r)}(z) = L_k^{(r-1)}(-z),$$

respectively, where $L_k^{(r)}(z)$ are the Laguerre polynomials. For $m, n \in \mathbb{Z}_{\geq 0}$, we define a family of polynomials $\widehat{R}_{m,n}^{(r)} = \widehat{R}_{m,n}^{(r)}(z)$ by

$$(C.2) \quad \widehat{R}_{m,n}^{(r)}(z) = \begin{vmatrix} \hat{q}_1^{(r)} & \hat{q}_0^{(r)} & \cdots & \hat{q}_{-m+2}^{(r)} & \hat{q}_{-m+1}^{(r)} & \cdots & \hat{q}_{-m-n+3}^{(r)} & \hat{q}_{-m-n+2}^{(r)} \\ \hat{q}_3^{(r)} & \hat{q}_2^{(r)} & \cdots & \hat{q}_{-m+4}^{(r)} & \hat{q}_{-m+3}^{(r)} & \cdots & \hat{q}_{-m-n+5}^{(r)} & \hat{q}_{-m-n+4}^{(r)} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \hat{q}_{2m-1}^{(r)} & \hat{q}_{2m-2}^{(r)} & \cdots & \hat{q}_m^{(r)} & \hat{q}_{m-1}^{(r)} & \cdots & \hat{q}_{m-n+1}^{(r)} & \hat{q}_{m-n}^{(r)} \\ \hat{p}_{n-m}^{(r)} & \hat{p}_{n-m+1}^{(r)} & \cdots & \hat{p}_{n-1}^{(r)} & \hat{p}_n^{(r)} & \cdots & \hat{p}_{2n-2}^{(r)} & \hat{p}_{2n-1}^{(r)} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \hat{p}_{-n-m+4}^{(r)} & \hat{p}_{-n-m+5}^{(r)} & \cdots & \hat{p}_{-n+3}^{(r)} & \hat{p}_{-n+4}^{(r)} & \cdots & \hat{p}_2^{(r)} & \hat{p}_3^{(r)} \\ \hat{p}_{-n-m+2}^{(r)} & \hat{p}_{-n-m+3}^{(r)} & \cdots & \hat{p}_{-n+1}^{(r)} & \hat{p}_{-n+2}^{(r)} & \cdots & \hat{p}_0^{(r)} & \hat{p}_1^{(r)} \end{vmatrix}.$$

For $m, n \in \mathbb{Z}_{< 0}$, we define $\widehat{R}_{m,n}^{(r)}$ by

$$(C.3) \quad \widehat{R}_{m,n}^{(r)} = (-1)^{m(m+1)/2} \widehat{R}_{-m-1,n}^{(r)}, \quad \widehat{R}_{m,n}^{(r)} = (-1)^{n(n+1)/2} \widehat{R}_{m,-n-1}^{(r)}.$$

Moreover, we introduce $S_{m,n} = S_{m,n}(t, s)$ as

$$(C.4) \quad \widehat{R}_{m,n}^{(r)}(z) = \widehat{S}_{m,n}(t, s), \quad z = \frac{t}{2}, \quad r = 2s - m + n.$$

Then, $\varphi_i = \varphi_i(t, s)$ given by

$$(C.5) \quad \begin{aligned} \varphi_0 &= \frac{\sqrt{t}}{2} \frac{\widehat{S}_{m,n}(t, s) \widehat{S}_{m-1,n-1}(t, s-1)}{\widehat{S}_{m,n-1}(t, s) \widehat{S}_{m-1,n}(t, s-1)}, \\ \varphi_1 &= \frac{\sqrt{t}}{2} \frac{\widehat{S}_{m-1,n}(t, s-1) \widehat{S}_{m,n-1}(t, s+1)}{\widehat{S}_{m,n}(t, s) \widehat{S}_{m-1,n-1}(t, s)}, \\ \varphi_2 &= \frac{\sqrt{t}}{2} \frac{\widehat{S}_{m-1,n-1}(t, s) \widehat{S}_{m,n}(t, s-1)}{\widehat{S}_{m-1,n}(t, s-1) \widehat{S}_{m,n-1}(t, s)}, \\ \varphi_3 &= \frac{\sqrt{t}}{2} \frac{\widehat{S}_{m,n-1}(t, s) \widehat{S}_{m-1,n}(t, s)}{\widehat{S}_{m-1,n-1}(t, s) \widehat{S}_{m,n}(t, s)}, \end{aligned}$$

solve the symmetric form of P_V (1.10) for the parameters

$$(C.6) \quad (\alpha_0, \alpha_1, \alpha_2, \alpha_3) = \left(\frac{1}{2} - s - n, s, \frac{1}{2} - s + m, s - m + n \right).$$

Let us consider the continuum limit of the result in Theorem 2.3. First, we note that the continuous q -Laguerre polynomials $P_k^{(\alpha)}(y|q)$ are expressed as

$$(C.7) \quad P_k^{(\alpha)}(y|q) = \frac{(q^{\alpha+1}; q)_k}{(q; q)_k} {}_3\phi_2 \left(\begin{matrix} q^{-k}, q^{\frac{1}{2}\alpha + \frac{1}{4}} e^{i\theta}, q^{\frac{1}{2}\alpha + \frac{1}{4}} e^{-i\theta} \\ q^{\alpha+1}, 0 \end{matrix} \middle| q; q \right),$$

$y = \cos \theta$ for $k \geq 0$,

in terms of the basic hypergeometric functions [6]. Then, from (2.2), (2.6) and (2.7), the polynomials $p_k^{(b)}(y|q)$ are written as

$$(C.8) \quad p_k^{(b)}(y|q) = (q^{\frac{3}{4}} b^{-1})^k \frac{(b^2; q)_k}{(q; q)_k} {}_3\phi_2 \left(\begin{matrix} q^{-k}, -q^{-\frac{1}{2}} bx, -bx^{-1} \\ b^2, 0 \end{matrix} \middle| q; q \right) \quad \text{for } k \geq 0.$$

Setting

$$(C.9) \quad b = q^{\frac{r}{2}}, \quad q = e^{\frac{\varepsilon^2}{2}}, \quad x = -e^{\frac{\varepsilon}{2}} \sqrt{t},$$

and taking the limit as $\varepsilon \rightarrow 0$, we obtain

$$(C.10) \quad \lim_{\varepsilon \rightarrow 0} p_k^{(b)}(y|q) = \frac{(r)_k}{k!} {}_1F_1 \left(\begin{matrix} -k \\ r \end{matrix} \middle| \frac{t}{2} \right) = L_k^{(r-1)}(t/2) \quad \text{for } k \geq 0.$$

Similarly, we see that $q_k^{(b)}$ reduce to $L_k^{(r-1)}(-t/2)$. Thus, we get

$$(C.11) \quad \lim_{\varepsilon \rightarrow 0} R_{m,n}^{(b)}(y|q) = \widehat{R}_{m,n}^{(r)}(z),$$

and

$$(C.12) \quad \lim_{\varepsilon \rightarrow 0} S_{m,n}(x, a) = \widehat{S}_{m,n}(t, s),$$

with $a = q^s$. Finally, setting as

$$(C.13) \quad f_i = -e^{-\varepsilon \varphi_i},$$

we find that (2.10) and (2.9) reduce to (C.5) and (C.6), respectively. It is shown that q - P_V (1.1) reduce to the symmetric form of P_V (1.10) by the above limiting procedure in Section 1. Therefore, the rational solutions of q - P_V stated in Theorem 2.3 reduce to those of P_V .

REFERENCES

- [1] B. Grammaticos and A. Ramani, *Discrete Painlevé equations: coalescences, limits and degeneracies*, *Physica*, **A228** (1996), 160–171.
- [2] B. Grammaticos, A. Ramani and V. G. Papageorgiou, *Do integrable mappings have the Painlevé property*, *Phys. Rev. Lett.*, **67** (1991), 1825–1828.
- [3] K. Kajiwara, M. Noumi and Y. Yamada, *A Study on the fourth q -Painlevé Equation*, *J. Phys. A: Math. Gen.*, **34** (2001), 8563–8581.
- [4] K. Kajiwara, M. Noumi and Y. Yamada, *Discrete dynamical systems with $W(A_{m-1}^{(1)} \times A_{n-1}^{(1)})$ symmetry*, *Lett. Math. Phys.*, **60** (2002), 211–219.
- [5] K. Kajiwara, M. Noumi and Y. Yamada, private communication.
- [6] R. Koekoek and R. F. Swarttouw, *The Askey-scheme of hypergeometric orthogonal polynomials and its q -analogue*, Delft University of Technology, Department of Technical Mathematics and Informatics Report no. 98-17 (1998).
- [7] K. Koike, *On the decomposition of tensor products of the representations of the classical groups: by means of the universal characters*, *Adv. Math.*, **74** (1989), 57–86.
- [8] T. Masuda, Y. Ohta and K. Kajiwara, *A determinant formula for a class of rational solutions of Painlevé V equation*, *Nagoya Math. J.*, **168** (2002), 1–25.
- [9] M. Noumi, S. Okada, K. Okamoto, and H. Umemura, *Special polynomials associated with the Painlevé equations II*, *Proceedings of the Taniguchi Symposium, 1997, Integrable Systems and Algebraic Geometry* (M. H. Saito, Y. Shimizu, K. Ueno, eds.), Singapore: World Scientific (1998), pp. 349–372.
- [10] M. Noumi and Y. Yamada, *Higher order Painlevé equations of type $A_l^{(1)}$* , *Funkcial. Ekvac.*, **41** (1998), 483–503.
- [11] K. Okamoto, *Studies on the Painlevé equations III, second and fourth Painlevé equations, P_{II} and P_{IV}* , *Math. Ann.*, **275** (1986), 222–254.
- [12] A. Ramani, B. Grammaticos and J. Hietarinta, *Discrete versions of the Painlevé equations*, *Phys. Rev. Lett.*, **67** (1991), 1829–1832.
- [13] H. Sakai, *Rational surfaces associated with affine root systems and geometry of the Painlevé equations*, *Commun. Math. Phys.*, **220** (2001), no. 1, 165–229.
- [14] T. Tokihiro, D. Takahashi, J. Matsukidaira and J. Satsuma, *From soliton equations to integrable cellular automata through a limiting procedure*, *Phys. Rev. Lett.*, **76** (1996), 3247–3250.
- [15] M. Taneda, *Polynomials associated with an algebraic solution of the sixth Painlevé equation*, to appear in *Jap. J. Math.*, **27** (2001).
- [16] H. Umemura, *Special polynomials associated with the Painlevé equations I*, preprint.

Department of Mathematics
 Kobe University
 Rokko
 Kobe, 657-8501
 Japan
 masuda@math.kobe-u.ac.jp