

## Heritability of Hwabyung Symptoms in South Korean Adolescent and Young Adult Twins

Yoon-Mi Hur,<sup>1</sup> Eun-ji Choi,<sup>2</sup> Jong-Woo Kim,<sup>2</sup> Hee-Jeong Jin,<sup>3</sup> and Siwoo Lee<sup>3</sup>

<sup>1</sup>Research Institute for the Welfare Society, Mokpo National University, Jeonnam, South Korea

<sup>2</sup>Department of Neuropsychiatry, Kyung Hee University, Seoul, South Korea

<sup>3</sup>Mibyeong Research Center, Korea Institute of Oriental Medicine, Daejeon, South Korea

The present study aimed to estimate heritability of Hwabyung (HB) symptoms in adolescent and young adult twins in South Korea. The sample included 1,601 twins consisting of 143 pairs of monozygotic male (MZM), 67 pairs of dizygotic male (DZM), 295 pairs of monozygotic female (MZF), 114 pairs of dizygotic female (DZF), and 117 pairs of opposite-sex dizygotic (OSDZ) twins and 129 twins with non-participating co-twins (mean age = 19.1 ± 3.1 years; range: 12–29 years). An HB symptom questionnaire was given to twins via a telephone interview. Consistent with the literature of HB, the mean level of HB was significantly higher in females than in males. Maximum likelihood twin correlations for HB were 0.31 (95% CI [0.16, 0.45]) for MZM, 0.19 (95% CI [-0.05, 0.41]) for DZM, 0.50 (95% CI [0.41, 0.58]) for MZF, 0.28 (95% CI [0.11, 0.44]) for DZF, and 0.23 (95% CI [0.05, 0.40]) for OSDZ twins. These patterns of twin correlations suggested the presence of additive genetic influences on HB. Model-fitting analysis showed that additive genetic and individual-specific environmental influences on HB were 44% (95% CI [37, 51]) and 56% (95% CI [49, 63]), respectively. Shared environmental influences were not significant. These parameter estimates were not significantly different between two sexes, and did not change significantly with age in the present sample, suggesting that genetic and environmental influences on HB in both sexes are stable across adolescence and young adulthood.

■ **Keywords:** Hwabyung, anger, heritability, adolescents, twin, structural equation model

Hwabyung (HB; known as an anger syndrome) is characterized by chronic suppression of anger, continued feelings of unfairness and resentment, subjective anger, external anger, somatic complaints, heat sensation, chest tightness, sighing, fatigue, insomnia, dysphoric affect, and indigestion (American Psychiatric Association, 1994; Min, 2013). HB was listed as culture-bound syndrome in the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* (American Psychiatric Association, 1994). However, it was deleted from the category of cultural syndromes in the *DSM-V* (American Psychiatric Association, 2013). The prevalence of HB syndrome has been estimated to be between 4.2% and 13.3% in the Korean general population and Korean immigrants in Western countries, and the prevalence is known to be higher in females than in males and is lower in higher social classes (Kim & Park, 2004; Lee & Lee, 2008; Min et al., 1990; Rhi, 2004).

Prior studies have shown that HB is frequently comorbid with many psychiatric disorders, including major depres-

sive disorder, generalized anxiety disorder, panic disorder, somatization disorder, dysthymia, and post-traumatic embitterment disorder (PTED), a subtype of post-traumatic stress disorder (PTSD; Joe et al., 2017; Kim et al., 2010; Lin et al., 1992; Min & Suh, 2010). As with many other psychiatric illnesses, the onset of HB has been associated with exposures to stressful life events. For example, Chung et al. (2013) found in a sample of South Koreans that major stressful life events that preceded the onset of HB were spousal infidelity, serious financial problems, and family conflicts. Chung et al. (2013) further showed that the risk for developing HB increased with lower social

RECEIVED 30 January 2018; ACCEPTED 1 May 2018. First published online 22 June 2018.

ADDRESS FOR CORRESPONDENCE: Siwoo Lee, Mibyeong Research Center, Korea Institute of Oriental Medicine, Daejeon, South Korea. E-mail: [bfree@kiom.re.kr](mailto:bfree@kiom.re.kr)

support, higher spousal criticisms, and higher interpersonal relationship problems.

Several researchers suggested that the collectivism cultures in East Asian countries play a strong etiologic role in HB. In collectivistic cultures, people are socialized to internalize stress and repress negative emotions such as anger, hostility, and psychological distress to maintain harmonious social relationships (Chiao, 2015). If aversive emotion is not relieved properly and accumulated over a long period, then the negative emotion reaches a threshold limit and often explodes in rage, which can lead to the development of HB (Pang, 1990). After a review of large medical records of the patients with HB, Min (2008) concluded that one of the most common precipitating factors of HB in South Korean women was 'unfair treatment'. Many South Korean female patients with HB felt that they were victims of unfair treatment, especially the chronic verbal and physical violence of husbands and in-law families. The patients typically felt that they were supposed to suppress their anger to conform to the Korean traditional patriarchal system and collectivism cultures where housewives are obliged to obey their husbands and in-law families. Other precipitating factors of HB reported by Min (2008) included poverty resulting from unfair social deprivation, experiences of social injustice, discrimination, and unfair juridical decisions.

In a functional magnetic resonance imaging (fMRI) study on neural responses to neutral, sad, and angry facial stimuli, Lee et al. (2009) found that as compared to healthy controls, HB patients showed increased activations in the lingual gyrus and fusiform gyrus and reduced activation in the thalamus in responses to all three types of facial stimuli. HB patients also showed lower activities in the right anterior cingulate cortex (ACC) in response to the neutral condition than did healthy controls. These findings indicated that the suppression of emotion may lead to aberrant function of the brain regions of the visual pathway and functional impairment in the ACC. However, as the sample size of the Lee et al.'s study is small ( $n = 24$ ), the findings need to be replicated in a larger sample to draw a firm conclusion.

While several psychosocial and neurological risk factors contributing to the development of HB have been suggested, the genetic etiology of HB remains poorly understood. In an early study, Min et al. (1990) found that the rates of HB and psychiatric illnesses such as depression, anxiety disorders, insomnia, personality disorders, violence, and suicide were higher in the first-degree relatives of the HB patients than those of healthy controls, suggesting that HB may have a genetic basis. The major goal of the present study, therefore, was to explore genetic and environmental influences on the HB symptoms in South Korean adolescent and young adult twins. We also examined age and sex differences in genetic and environmental influences on HB during adolescence and young adulthood.

## Methods

### Sample

The sample included 1,601 twins consisting of 143 pairs of monozygotic male (MZM), 67 pairs of dizygotic male (DZM), 295 pairs of monozygotic female (MZF), 114 pairs of dizygotic female (DZF), 117 pairs of opposite-sex dizygotic (OSDZ) twins, and 129 twins with non-participating co-twins drawn from the South Korean Twin Registry (Hur et al., 2013). The mean age of the twins was  $19.1 \pm 3.1$  years (range 12–29 years). Twins under 20 years of age were recruited mostly from schools throughout South Korea, while those of 20 years of age or older were from Facebook, twin clubs on the internet, and colleges throughout South Korea. The present sample has an over-representation of females (62%), which was due in part to the fact that some of the male twins were in the military service at the time of the recruitment of the subjects because it is mandatory for young adult males in South Korea to complete army service. In terms of parental educational achievement, 49% of fathers and 41% of mothers of twins completed university education or more.

Zygosity of the twins was assessed using a three-item zygosity questionnaire. As compared to DNA analysis, this questionnaire method has been shown to achieve over 90% accuracy (Ooki et al., 1993). The number of monozygotic (MZ) twins was much greater than that of dizygotic (DZ) twins in the present sample, which likely reflected the low DZ twin birth rates in the South Korean population for the birth cohorts in the present study rather than sampling bias (Hur & Kwon, 2005).

### Measures

To measure HB, we used the HB symptom scale (Kwon et al., 2008), which consists of 15 self-report items concerning typical symptoms of HB. Psychometric properties of the HB symptom scale have been intensively studied, and reliabilities and validity of the scale were found to be acceptable (Kwon et al., 2008). A telephone interview was conducted to assess twins. In the interview, twins were instructed to rate themselves on a 5-point Likert scale from *not true* (0) to *certainly true* (4) for each of the 15 items. The ratings were summed to obtain a total score so that higher scores represent more severe symptoms. As with many other psychopathology measures, the total score of the HB symptom scale was somewhat positively skewed, with a skewness index of 0.65. We performed square-root transformation of the scores of the HB symptoms scale, which resulted in a distribution of skewness of -0.14. Thus, twin correlation and model-fitting analyses below were based on the transformed data. Cronbach's alpha of the 15 items was 0.92 in the present sample, which was close to that in the normative sample (Kwon et al., 2008).

**TABLE 1**  
Means, Standard Deviations, and Maximum Likelihood Twin Correlations and Their 95% CIs for Hwabyung Syndrome by Zygosity

	MZM	DZM	MZF	DZF	OSDZ
N (individual)	318 (143 pairs + 32)	151 (67 pairs + 17)	628 (295 pairs + 38)	247 (114 pairs + 19)	257 (117 pairs + 23)
Mean (SD)	14.6 (10.5)	14.4 (10.3)	17.1 (11.1)	16.2 (10.6)	16.2 (10.9)
Twin correlation	0.31 [0.16, 0.45]	0.19 [-0.05, 0.41]	0.50 [0.41, 0.58]	0.28 [0.11, 0.44]	0.23 [0.05, 0.40]

Note: 95% CIs are in parentheses. MZM = monozygotic male twins; DZM = dizygotic male twins; MZF = monozygotic female twins; DZF = dizygotic female twins; OSDZ = opposite-sex dizygotic twins. Twin correlations were based on the square-root transformed data.

## Statistical Analysis

To estimate genetic and environmental influences on HB, we conducted maximum likelihood twin correlation and structural equation model-fitting analyses. A general sex-limitation model (Neale & Cardon, 1992) was applied to the raw data for HB. The full model included additive genetic factors (A) (i.e., the sum of the average effect of all alleles that influence a trait), shared environmental factors (C) (i.e., those environmental factors shared between the two members of a twin pair), and individual-specific environmental factors (E) (i.e., those environmental factors unique to each member of a twin pair and measurement error). On the basis of the degree of genetic relatedness, the correlations for A were set at 1.0 for MZ, and 0.50 for the same-sex DZ twins for both sexes. The correlation for C was set at 1.0 for both types of twins for both sexes. The correlation for E was set at 0.0 because by definition, this factor was uncorrelated between the two members of a twin pair. To determine sex differences in genetic and environmental influences on HB, we allowed the magnitudes of A, C, and E to differ across sexes in the full model. In addition, to test sex-specific genetic effects, the genetic correlation for opposite-sex DZ twins was set to be different from 0.5. Age was treated as a covariate in the model.

We used the maximum likelihood raw data option in Mx (Neale et al., 2003), which calculates twice the negative log-likelihood (-2LL) of the data. To determine the best-fitting, most parsimonious model, parameters in the full model are constrained sequentially, and the resulting changes in -2LL were evaluated. As the difference in -2LL is chi-square distributed, when models were nested to each other, the likelihood ratio test (LRT) was used to compare alternative models. A significant change in chi-square would indicate that the reduction of the parameters in the nested model causes a significant decrease in model fit, whereas a non-significant change would suggest that constraining parameters in the nested model is acceptable. When alternative models were not nested to each other, Akaike's information criterion ( $AIC = -2LL - 2df$ ) for alternative models were compared to evaluate superiority among competing models. Models having lower AIC are considered more parsimonious and are thus preferred (Akaike, 1987).

## Results

### Descriptive Statistics and Twin Correlations

Table 1 presents means, standard deviations, and maximum likelihood twin correlations for HB by zygosity. Although there was no significant sex difference in the variance of HB, the mean was significantly higher in females than in males ( $t = 4.7, p < .001$ ). The correlation between age and HB was significant but small ( $r = 0.07$ ).

As a preliminary step, we computed maximum likelihood twin correlations across zygosity groups to predict the results from general sex-limitation model-fitting analysis. As can be seen in Table 1, MZ twin correlations were greater than DZ twin correlations in both sexes, suggesting the presence of genetic effects on HB. The sizes of DZ twin correlations were only slightly greater than half the MZ correlations in both sexes, suggesting that shared environmental influences may be very small. The size of OSDZ twin correlation was not significantly lower than same-sex DZ twins, which suggested that genes for HB may not be sex-specific. MZ correlations were much less than 1.0 in both sexes, indicating that individual-specific environmental factors may be important in HB. These observations of the twin correlations were formally tested using general sex-limitation model-fitting analysis.

### Model-Fitting Analysis

Table 2 presents the results of model-fitting analysis. When we set the genetic correlation to be the same across OSDZ and same-sex twins, the resulting chi-square change was negligible (Model 2). When we removed age covariate from the full model (Model 3), the resulting chi-square change was significant, suggesting that the age covariate should be retained in the model. We next equated the magnitudes of additive genetic, shared, and individual-specific environmental effects across two sexes, which yielded no significant change in chi-square (Model 4). Taken together, these result suggested that sex-specific genes might not be present in HB, and that the magnitudes of genetic and shared and individual-specific environmental influences on HB might be the same across sexes.

Models 5 and 6 removed shared environmental effects and additive genetic effects, respectively, from Model 2. The resulting chi-square change was significant in Model 6 but

**TABLE 2**  
Model-Fitting Results for the Hwabyung Symptoms Scale

Model	Description	Goodness-of-fit statistics					Parameter estimates					
		-2LL	AIC	df	$\Delta$ -2LL	$\Delta$ df	Male			Female		
							A	C	E	A	C	E
1	Full model	5,074.1	2,108.1	1,483			0.16 [0.00, 0.43]	0.14 [0.00, 0.39]	0.70 [0.57, 0.84]	0.42 [0.08, 0.58]	0.09 [0.00, 0.39]	0.50 [0.42, 0.59]
2	$r_g$ for OSDZ twins = 0.5	5,074.1	2,106.1	1,484	0.0	1	0.99	0.16 [0.00, 0.43]	0.14 [0.00, 0.39]	0.70 [0.57, 0.83]	0.42 [0.08, 0.58]	0.50 [0.42, 0.58]
3	$r_g$ for OSDZ twins = 0.5; Age covariate = 0	5,080.6	2,110.6	1,485	6.6	2	0.04	0.15 [0.00, 0.42]	0.15 [0.00, 0.38]	0.71 [0.57, 0.84]	0.39 [0.07, 0.58]	0.49 [0.42, 0.58]
4	$r_g$ for OSDZ twins = 0.5; $A_m = A_r, C_m = C_r, E_m = E_r$	5,080.6	2,106.6	1,487	6.5	4	0.16	0.36 [0.11, 0.50]	0.08 [0.00, 0.29]	0.57 [0.50, 0.64]	0.36 [0.11, 0.50]	0.57 [0.50, 0.64]
5	$r_g$ for OSDZ twins = 0.5; $A_m = A_r, E_m = E_r, C_m = 0, C_r = 0$	5,081.1	2,105.1	1,488	7.0	5	0.22	<b>0.44 [0.37, 0.51]</b>	-	<b>0.56 [0.49, 0.63]</b>	<b>0.44 [0.37, 0.51]</b>	<b>0.56 [0.49, 0.63]</b>
6	$r_g$ for OSDZ twins = 0.5; $C_m = C_r, E_m = E_r, A_m = 0, A_r = 0$	5,088.7	2,112.7	1,488	14.6	5	0.01	-	0.37 [0.30, 0.43]	0.63 [0.57, 0.70]	-	0.63 [0.57, 0.70]

Note: Model-fitting analyses were based on the square-root transformed data. LL = log likelihood. The best-fitting model is indicated in bold type. A = additive genetic variance, C = shared environmental variance, E = individual-specific environmental variance plus measurement error.  $r_g$  = genetic correlation. OSDZ = opposite-sex dizygotic twin. Subscripts m and f represent male and female, respectively.

not in Model 5, indicating that shared environmental effects are negligible, while genetic influences are significant. From these model comparisons, we concluded that Model 5 was the best fit, where additive genetic and individual-specific environmental effects including measurement error was 44% (95% CI [37%, 51%]) and 56% (95% CI [49%, 63%]), respectively.

## Discussion

The present study was the first to report heritability of HB. Model-fitting analysis revealed that 44% of the variation in HB symptoms was due to genetic effects, with the remaining variance being attributable to individual-specific environmental influences and measurement error. The genetic contribution to HB found in the present study points to a need to conduct molecular genetic studies in the future for a better understanding of the pathophysiology of HB. In an association study of the oxytocin receptor gene (*OXT* rs53576), Kim et al. (2011) found that as compared to those who carry the A allele, those with the G allele exhibited more culturally congruent patterns of emotion regulation. While suppression of emotion is a normative behavior in Korean collectivistic cultures, expression of emotion is a normative behavior in Western individualistic cultures. Consistent with cultural norms, the American G allele carriers showed less habitual use of emotional suppression than did the American A allele carriers, whereas the Korean G allele carriers showed greater habitual use of emotional suppression than did the Korean A allele carriers. These results suggest a possibility that expression of genes related to HB may be modified by cultural norms. Although emotion regulation is likely to be an endophenotype of HB, the Kim et al. study employed a candidate gene approach that has been criticized for the high rate of false positive findings (Marian, 2012). Thus, it is necessary in the future to carry out genome-wide association studies of HB to identify genes involved in HB symptoms.

The magnitude of genetic influence found in HB in the present sample was broadly similar to those found in affective disorders and stress-related psychopathology such as depression, anxiety disorders, somatization, and PTSD (Duncan et al., 2017; Shimada-Sugimoto et al., 2015). Given the frequent comorbidity of HB with these disorders, multivariate genetic studies may need to be performed in the future to elucidate shared genetic basis of HB and related disorders.

In the present sample, we found no significant sex differences in genetic and environmental influences on HB. Similar results were reported in twin studies of anxiety disorders (Shimada-Sugimoto et al., 2015) but not in those of depression or PTSD (Bangasser & Valentino, 2014; Perry et al., 2017). Given the higher incidence of stress-related psychiatric disorders in females (Marcus et al., 2005) and sex differences in biological pathways to psychiatric disorders

reported in the literature (e.g., Perry et al., 2017), it is possible that our sample size was not sufficiently large to detect sex differences in genetic and environmental influences on HB. Future twin studies should, therefore, increase sample size to test sex differences.

Our twin model assumes that there is no gene environment (GE) correlation or gene by environment interaction ( $G \times E$ ). It has been argued that omission of GE correlation and  $G \times E$  in a twin model may lead to an overestimation of genetic and/or environmental variance components (Eaves et al., 2003). Genetically vulnerable individuals may create a stressful environment that can precipitate anger, which can contribute to the development of HB (GE correlation). Genetically influenced risk factors for HB include hasty, impatient, and convulsive personality characteristics (Moon et al., 1988), poor social skills and stress coping strategies (Kim & Kim, 2013), and low self-esteem (Park & Chae, 2001). Longitudinal twin studies may be necessary to delineate the process of GE correlation in the development of HB.

Genetic factors in HB may be expressed differently across different environmental settings ( $G \times E$ ). Parental education is an example of shared environment. Given the findings that lower parental education is associated with a higher risk factor for HB (Kim & Park, 2004; Lee & Lee, 2008; Min et al., 1990; Rhi, 2004), we conducted a  $G \times E$  model-fitting analysis including the parental educational level as a moderator. However, genetic and environmental influences on HB in our sample did not differ significantly across the levels of parental education (data available upon request), suggesting that our heritability estimate may not be inflated due to the effect of gene by parental education interaction.

Family conflicts have been consistently suggested to be risk factors for the development of HB (Kim & Park, 2004). However, our study showed that shared environmental effects were negligible and that environmental effects on HB were mainly individual-specific, suggesting that family conflicts relevant to HB may be specific to individuals within the family. If family conflicts are subsumed under shared environmental influences in the development of HB, then our heritability estimate may encompass the gene by family conflict interaction component. If, however, family conflicts are subsumed under individual-specific environment, then our estimate of individual-specific environment may include the gene by family conflict interaction component.

Unfortunately, we did not measure family conflicts in our study, and we were not able to examine how family conflicts act and interact with genetic effects on HB.

There were a few limitations in this study that deserve mention. First, we collected the HB data through telephone interviews. Although some studies suggest that telephone interview is comparable to the face-to-face interviews in assessing psychiatric phenotypes (e.g., Sobin et al., 1993), others suggest that responses elicited by telephone are sus-

ceptible to social desirability bias (e.g., Sibbald et al., 1994). Therefore, it is necessary to replicate our findings with the face-to-face interview method. Second, our sample only included adolescents and young adults, and therefore heritability of HB found in our sample may not be generalized in other age groups. Third, HB is known to be more prevalent in middle-aged adults than in adolescent populations (Min, 2008). Thus, one could argue that patients with HB may not be present and only mild cases may have been available in our sample. However, on the basis of the diagnostic criteria of the HB symptom scale (Kwon et al., 2008), approximately 10% of the present sample could be considered at risk for the development of HB, which is within the range of the prevalence rate of HB estimated for the general population in South Korea (Kim & Park, 2004; Lee & Lee, 2008; Min et al., 1990; Rhi, 2004). Finally, we employed a symptom scale to measure the phenotype of HB. However, heritability estimate may be different if a clinical diagnosis of HB is employed.

## Acknowledgments

This work was supported by the ‘Development of Health Prediction Technology based on Big Data’ (K18092) funded by the Ministry of Science and ICT (MSIT) of South Korea, given to the Korea Institute of Oriental Medicine (KIOM).

## References

- Akaike, H. (1987). Factor analysis and AIC. *Psychometrika*, 52, 317–332.
- American Psychiatric Association. (1994). *Diagnostic and statistical manual of mental disorders* (4th ed.). Washington, DC: Author.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Washington, DC: Author.
- Bangasser, D. A., & Valentino, R. J. (2014). Sex differences in stress-related psychiatric disorders: Neurobiological perspectives. *Frontiers in Neuroendocrinology*, 35, 303–319.
- Chiao, J. Y. (2015). Current emotion research in cultural neuroscience. *Emotion Review*, 7, 280–293.
- Chung, S.-Y., Song, S.-Y., & Kim, J.-W. (2013). Clinical guidelines for Hwabyung II (research on the status of Hwabyung in Korea). *Journal of Oriental Neuropsychiatry*, 24, 15–22.
- Duncan, L. E., Ratanatharathorn, A., Aiello, A. E., Almli, L. M., Amstadter, A. B., Ashley-Koch, A. E., ... Koenen, K. C. (2017). Largest GWAS of PTSD (N = 20070) yields genetic overlap with schizophrenia and sex differences in heritability. *Molecular Psychiatry*, 23, 666–673.
- Eaves, L., Silberg, J., & Erkanli, A. (2003). Resolving multiple epigenetic pathways to adolescent depression. *Journal of Child Psychology and Psychiatry*, 44, 1006–1014.
- Hur, Y.-M., & Kwon, J. S. (2005). Changes in twinning rates in South Korea; 1981–2002. *Twin Research and Human Genetics*, 8, 76–79.

- Hur, Y.-M., Jeong, H.-U., Chung, K. W., Shin, J. S., & Song, T.-B. (2013). The South Korean Twin Registry: An update. *Twin Research and Human Genetics*, *16*, 237–240.
- Joe, S., Lee, J. S., Kim, S. Y., Won, S. H., Lim, J. S., & Ha, K. S. (2017). Posttraumatic Embitterment disorder and Hwabyung in the general Korean population. *Psychiatry Investigation*, *14*, 392–399.
- Kim, H.-K., & Park, J.-Y. (2004). Prevalence and related factors of Hwabyung for the aged woman in rural community. *Journal of Korea Community Health Nursing Academic Society*, *18*, 234–242.
- Kim, H. S., Sherman, D. K., Mojaverian, T., Sasaki, J. Y., Park, J., Suh, E. M., & Taylor, S. E. (2011). Gene-culture interaction: Oxytocin receptor polymorphism (OXTR) and emotion regulation. *Social Psychological and Personality Science*, *2*, 665–672.
- Kim, J.-W., Chung, S.-Y., Suh, H.-U., Jung, I.-C., Lee, S.-G., Kim, B.-K., & Kim, S.-H. (2010). The characteristics of Hwa-byung patients based on Hwa-byung epidemiologic data. *Journal of Oriental Neuropsychiatry*, *21*, 157–169.
- Kim, J.-W., & Kim, S. Y. (2013). Clinical guidelines for Hwabyung I. (Overview). *Journal of Oriental Neuropsychiatry*, *21*, 157–169.
- Kwon, J. H., Kim, J. W., Park, D. G., Lee, M. S., Min, S. G., & Kwon, H. I. (2008). Development and validation of the Hwa-Byung Scale. *Korean Journal of Clinical Psychology*, *27*, 237–252.
- Lee, B. T., Paik, J. W., Kang, R. H., Chung, S. Y., Kwon, H. I., Khang, H. S., ... Ham, B. J. (2009). The neural substrates of affective face recognition in patients with Hwa-byung and healthy individuals in Korea. *The World Journal of Biological Psychiatry*, *10*, 552–559.
- Lee, J.-K., & Lee, J.-H. (2008). Study on the prevalence of Hwa-byung diagnosed by HBDIS in general population in Kangwon Province. *Journal of Oriental Neuropsychiatry*, *19*, 133–139.
- Lin, K. M., Lau, J. K., Yamamoto, J., Zheng, Y. P., Kim, H. S., Cho, K. H., & Nakasaki, G. (1992). Hwa-byung. A community study of Korean-Americans. *The Journal of Nervous and Mental Disease*, *180*, 386–391.
- Marcus, S. M., Young, E. A., Kerber, K. B., Kornstein, S., Farabaugh, A. H., Mitchell, J., ... Rush, A. J. (2005). Gender differences in depression: Findings from the STAR\*D study. *Journal of Affective Disorders*, *87*, 141–150.
- Marian, A. J. (2012). Molecular genetic studies of complex phenotypes. *Translational Research*, *159*, 64–79.
- Min, S. K. (2008). Clinical correlates of Hwa-byung and a proposal for a new anger disorder. *Psychiatry Investigation*, *5*, 125–141.
- Min, S. K. (2013). Culture and somatic symptoms: Hwa-byung, a culture-related anger syndrome. In K. Koh (Eds.), *Somatization and psychosomatic symptoms* (pp. 51–60). New York, NY: Springer.
- Min, S. K., Namkoong, K., & Lee, H. (1990). An epidemiological study on Hwabyung. *Journal of the Korean Neuropsychiatric Association*, *29*, 867–874.
- Min, S. K., & Suh, S. Y. (2010). The anger syndrome Hwabyung and its comorbidity. *Journal of Affective Disorders*, *124*, 211–214.
- Moon, C. M., Kim, J. H., & Whang, W. W. (1988). The bibliographical study on stress and Hwa. *Journal of Korean Oriental Internal Medicine*, *9*, 153–159.
- Neale, M. C., & Cardon, L. R. (1992). *Methodology for genetic studies of twins and families*. London: Kluwer.
- Neale, M., Boker, S. M., Xie, G., & Maes, H. (2003). *Mx: Statistical modeling*. Richmond, VA: Department of Psychiatry, Virginia Institute for Psychiatric and Behavioral Genetics, Virginia Commonwealth University.
- Ooki, S., Yamada, K., & Asaka, A. (1993). Zygosity diagnosis of twins by questionnaire for twins' mothers. *Acta Geneticae Medicae et Gemellologicae (Roma)*, *42*, 17–22.
- Pang, K. Y. (1990). Hwabyung: The construction of a Korean popular illness among Korean elderly immigrant women in the United States. *Culture, Medicine, and Psychiatry*, *14*, 495–512.
- Park, Y. S., & Chae, E. O. (2001). A study on Hwabyung. *Community Nursing*, *12*, 705–715.
- Perry, L. M., Goldstein-Piekarski, A. N., & Williams, L. M. (2017). Sex differences modulating serotonergic polymorphisms implicated in the mechanistic pathways of risk for depression and related disorders. *Journal of Neuroscience Research*, *95*, 737–762.
- Rhi, B. Y. (2004). Hwabyung – An overview. *Psychiatry Investigation*, *1*, 21–24.
- Shimada-Sugimoto, M., Otowa, T., & Hettema, J. M. (2015). Genetics of anxiety disorders: Genetic epidemiological and molecular studies in humans. *Psychiatry and Clinical Neurosciences*, *69*, 388–401.
- Sibbald, B., Addington-Hall, J., Brennehan, D., & Freeling, P. (1994). Telephone versus postal surveys of general practitioners: Methodological considerations. *British Journal of General Practice*, *44*, 297–300.
- Sobin, C., Weissman, M. M., Goldstein, R. B., Adams, P., Wickramaratne, P., Warner, V., & Lish, J. (1993). Diagnostic interviewing for family studies: Comparing telephone and face-to-face interview methods for the diagnosis of lifetime psychiatric disorders. *Psychiatric Genetics*, *3*, 227–233.