

If you're seeking referral to a Gender Identity Clinic to receive hormone treatment or surgery, this will only lengthen the process. Psychiatric assessments are not required by Gender Identity Clinics, and your GP is qualified to make the referral'.

Given the high proportion of students who will have families of origin outside Oxfordshire and the high level of awareness within Oxford University that a psychiatric assessment is unnecessary, it appears likely that a significant number are bypassing the service.

Another factor not considered by the authors is the almost exponential rise in people presenting to gender identity services who are under the age of 17. Provision is via a centralised national service for young people. At 17, their care is transferred to an appropriate gender identity clinic. Transfers of care now comprise an increasing proportion of referrals to the Northern Region Gender Dysphoria Service. As this is the case nationally, such patients will also bypass Dr Bass and will not feature in the statistics presented.

Michael Shaw, Consultant in Gender Dysphoria, Northern Region Gender Dysphoria Service; email: NRGDS@ntw.nhs.uk

doi:10.1192/bjb.2018.109



© The Author 2019. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

Contrast with referrals to Tavistock and Portman Gender Identity Disorder Service

This article shows a modest increase in referrals year on year, approximately 18%, with a majority of referrals received being for those assigned male at birth (AMAB). This is an interesting contrast with data from the Tavistock and Portman Gender Identity Disorder Service (GIDS) for children and adolescents.

Referrals to GIDS have increased from 97 in 2009/10 to 2016 in 2016/17. From 2014/15 to 2015/16, referrals increased by over 100% and from 2015/16 to 2016/17 they increased by 41%. Ages at referral seen by the service ranged from a very few at 3 to 17 years old.

Also in contrast to this much larger increase in referral rates is a marked change in the proportion of those assigned female at birth (AFAB). Up until 2011 there were more referrals of those AMAB. Since then the number of those AFAB referred has grown steadily, and in 2016/17 more than twice as many referrals were made for those AFAB as those AMAB (data available on GIDS website).

We need to be looking at these striking differences, and more research is required to determine the reasons for them. It may be that reducing stigma has led to higher referral rates, particularly among teenage girls but it could also be that the characteristics of those being referred are changing. This links with the finding that there seems to be a higher prevalence of autism spectrum conditions (ASC) in

clinically referred, gender dysphoric adolescents than in the general adolescent population. Holt, Skagerberg and Dunsford (2014) found that 13.3% of referrals to the GIDS service in 2012 mentioned comorbid ASC (although this is likely to be an underestimate).

In this context, it is alarming that referral rates are increasing at a rate that services and research cannot keep up with. Both the American Academy of Pediatrics and the Australian Standards of Care and Treatment for Transgender and Gender Diverse Children and Adolescents appear to support both medical and surgical transition in adolescents. And yet long-term outcomes in this group are not known. We know that adults who have gender dysphoria and who transition report the dysphoria beginning in early childhood. We do not know yet know that those experiencing dysphoria in childhood will go onto experience dysphoria in adulthood. Indeed, we know that 80% of individuals referred to GIDS do not proceed to transition.

In the UK, we are fortunate to have a national service for children that follows the best available evidence, but there is an urgent need for both research and discussion. This is not always easy in a highly emotionally charged atmosphere.

Katherine Rae Clyde, Consultant in Older Persons' Mental Health, Royal College of Psychiatrists; email: katie.clyde@nhs.net

doi:10.1192/bjb.2018.110



© The Author 2019. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

Author reply: Our paper aimed to highlight the marked increase in referrals seen locally within recent years. We accept that, as Dr Shaw notes, some people, particularly those from the student population who also have families of origin outside Oxfordshire, may have been referred directly to gender identity clinics and so were not captured by the data. This would mean that our findings likely represent an underestimate of the total increase in numbers of people seeking assistance from specialist clinics, which makes the increase we did find even more striking.

We also appreciate Dr Shaw's points regarding the referral pathway in Oxfordshire and the need to minimise the barriers transgender people experience to accessing services. The clinical pathway has indeed been reviewed and altered since the period described in the study, with service users now being referred directly by general practitioners, generally to the specialist clinic in Northamptonshire.

The clinic did not assess people under the age of 17, so we did not include data on this age group in our study, but it is certainly notable, as both Dr Shaw and Dr Clyde's letters highlight, that referrals to gender identity disorder services (GIDS) for children and adolescents have risen dramatically over the past five years. Dr Clyde in her letter draws attention to the high rates of referral to GIDS for children and adolescents since 2010, and in particular the increase in rates of

referral of those assigned female at birth. This has also been our experience, although the increase was less marked in our adult population than in the child and adolescent population Dr Clyde describes, and it is interesting to consider possible reasons for this. In our data, we found a marked trend towards more natal females being referred over the six-year period from 2011 to 2016, with a ratio of approximately 1:2 compared with natal males requesting transition. In our previous audit published in 2011 and covering the period 2006–2011, this ratio was 1:3 (Saunders and Bass, 2011). However this apparent trend did not reach statistical significance in our data. We also identified 8.5% of people presenting with non-binary gender identities, which were not noted in our previous audit. The finding of increased rates of autism spectrum disorder is also of considerable interest, and our finding of 7.8% is almost certainly an underestimate. We agree that clinics are being

overwhelmed, and that there is an urgent need for both more research and discussion in this rapidly changing field, in order to best meet the needs of transgender young people and adults.

Josephine Fielding and **Christopher Bass**, email: j.fielding2@nhs.net

doi:10.1192/bjb.2018.111



© The Authors 2019. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.