

## EPV1172

**The Delicate Balance: Aptitude of Physicians with Psychiatric diseases**

N. Khaterchi<sup>1</sup>, G. Bahri<sup>2</sup>, I. Youssef<sup>2</sup>, M. Mersni<sup>2</sup>, H. Ben Said<sup>2</sup>, D. Brahim<sup>2</sup>, N. Mechergui<sup>2</sup>, M. Methni<sup>3\*</sup>, C. Ben Said<sup>3</sup>, N. Bram<sup>3</sup> and N. Ladhari<sup>2</sup>

<sup>1</sup>occupational medicine; <sup>2</sup>Charles Nicolle Hospital and <sup>3</sup>Forensic Psychiatry departement, Razi Hospital, La Manouba, Tunis, Tunisia

\*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1721

**Introduction:** The delicate balance between the need to ensure quality patient care and the reality of physicians dealing with psychiatric diseases poses a major challenge within the medical field. This issue raises fundamental ethical, legal, and medical questions, highlighting the complexity of decision-making regarding professional aptitude for practitioners affected by mental disorders.

**Objectives:** To examine the impact of psychiatric diseases on the medical aptitude of physicians.

**Methods:** This was a retrospective descriptive study that focused on physicians with psychiatric diseases referred to the occupational pathology clinic at Charles Nicolle Hospital in Tunis for medical evaluations of their work aptitude between January 1, 2021, and September 15, 2023.

**Results:** During the study period, we collected data from 20 patients. The mean age was  $38 \pm 11$  years, with a sex-ratio (F/M) of 4.5. Five examined physicians had family histories of psychiatric disorders. Medical specialties were the most represented (N=17), including three general practitioners, two family medicine practitioners, and two anesthesiologists. The study population included 10 residents, eight hospital assistants, and two medical interns. The most common psychiatric diagnosis was depression (N=7), followed by bipolar disorder (N=5). The medical treatment prescribed was combinations of antidepressants and anxiolytics in seven cases, antipsychotics in five cases, and antidepressants in two cases. Medication adherence was noted in 10 physicians. Fourteen physicians had taken long-term sick leave, with an average duration of 203 days. Five physicians were declared fit to continue their regular professional activities, seven physicians were declared fit with restrictions on night work, and one physician was declared fit with workplace accommodations.

**Conclusions:** This study highlights the challenges surrounding the medical aptitude of physicians with psychiatric diseases. However, it is imperative to promote mental health awareness and to implement support measures to ensure both compassion for physicians and patient safety.

**Disclosure of Interest:** None Declared

## EPV1169

**Assessment of the Capability of Artificial Intelligence for Psychiatric Diagnosis**

N. Laherrán, R. Palacios\* and A. Vázquez

Psychiatry, Jerez de la Frontera University Hospital, Jerez De La Frontera, Spain

\*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1722

**Introduction:** Psychopathological exploration (PPE) involves an assessment of the mental state of patients, where psychological signs and symptoms are analyzed, which collectively form a syndrome. To conduct this assessment, the clinician must utilize their expertise to identify the presence and authenticity of a series of symptoms that, once recognized, allow for a diagnosis (1). The presence of this subjective component could explain why, despite the continuous growth of artificial intelligence (AI), its application in clinical psychiatry practice remains limited. However, the combination of the clinician's work with AI could enhance diagnostic accuracy and our understanding of diseases (2).

**Objectives:** The objective of this study is to investigate whether AI can make accurate diagnoses through an initial psychopathological evaluation.

**Methods:** A random sample was selected from our medical records of all patients admitted to our acute mental health inpatient unit through the hospital's emergency services in the year 2022. An anonymized database was created, including sociodemographic information, the results of the psychopathological assessment in the emergency department, and the diagnosis at the time of discharge. The psychopathological assessment conducted in the emergency department was provided to the AI chatbot ChatGPT, with a request to establish a diagnosis according to the DSM-5. Diagnoses such as brief psychotic disorder, schizophreniform disorder, and schizophrenia were considered, given their acute symptom similarities, as well as major depressive disorder (unipolar) and bipolar disorder. The level of agreement between both diagnoses was evaluated using the kappa coefficient.

**Results:** The sample consisted of 15 patients, of whom 60% were male, with a mean age of 45 years (standard deviation = 15.6). 73.3% of the patients had prior mental health follow-up, and 66.7% had been previously hospitalized. Diagnoses included psychotic disorder in 33.3% of cases, bipolar disorder with manic episode in 26.7%, depressive disorder in 13.3%, delusional disorder in 13.3%, schizoaffective disorder in 6.7%, and borderline personality disorder in 6.7%. A kappa value of 0.561 was obtained, indicating a moderate degree of agreement between the diagnoses.

**Conclusions:** Despite the inherent subjectivity in psychopathological assessment, this study suggests that AI, in the form of natural language processing chatbots like ChatGPT, can be a useful tool to assist mental health professionals in the diagnostic process. While AI shows promising potential, it should not entirely replace the experience and clinical judgment of mental health professionals. Instead, the importance of potential collaboration between AI and clinicians for achieving more precise and comprehensive diagnoses is highlighted.

**Disclosure of Interest:** None Declared

## EPV1172

**Dermatitis artefacta in a 60 year old man: a case report**

P. Setién Preciados\*, C. Díaz Mayoral and E. Arroyo Sánchez

Servicio de Psiquiatría, Hospital Universitario Príncipe de Asturias, Alcalá de Henares, Spain

\*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1723

**Introduction:** Dermatitis artefacta (DA), also known as factitious dermatitis, is a condition among factitious disorders, whereby self-