

HIPPARCOS EXTRAGALACTIC LINK

Preliminary Bonn, Potsdam and Kiev solutions

P. BROSCHE¹, M. GEFFERT¹, S. HIRTE², N. KHARCHENKO³,
V. KISLYUK³, M. ODENKIRCHEN¹, S. RYBKA³, E. SCHILBACH²,
R.-D. SCHOLZ², H.-J. TUCHOLKE¹ AND A. YATSENKO³

¹ *Sternwarte der Universität Bonn, Germany*

² *WIP Astronomie, Universität Potsdam, Germany*

³ *Main Astronomical Observatory, Kiev, Ukraine*

Hipparcos proper motions contain an unknown angular velocity ω relative to a non-rotating system. The basic equations for its derivation are:

$$\begin{aligned} \Delta\mu_\alpha \cos \delta &= -\omega_1 \cos \alpha \sin \delta - \omega_2 \sin \alpha \sin \delta + \omega_3 \cos \delta \\ \Delta\mu_\delta &= +\omega_1 \sin \alpha - \omega_2 \cos \alpha \end{aligned} \quad (1)$$

where $\Delta\mu_\alpha$ and $\Delta\mu_\delta$ are absolute minus Hipparcos proper motions.

	Bonn	Potsdam	Kiev
photographic plates	astrograph	Schmidt	astrograph
<i>m</i> link fields	8	10	183
<i>n</i> link stars	33	104	1015
galaxies per field	1 to 5	300 to 2000	3 to 5
base line [years]	70 to 90	20 to 40	20 to 40
random p.m. error per star [mas/yr]	0.5 to 1.5	3 to 5	5 to 12
syst. abs. p.m. error per field [mas/yr]	1.0 to 1.5	~ 2	~ 4
<i>rms</i> of solution of (1) [mas/yr]	5	8	14
$\omega_1 \pm \sigma(\omega_1)$ [mas/yr]	+1.2 ± 1.0	+0.8 ± 1.0	-1.5 ± 0.7
$\omega_2 \pm \sigma(\omega_2)$ [mas/yr]	+3.2 ± 0.7	-0.7 ± 1.0	-2.0 ± 0.5
$\omega_3 \pm \sigma(\omega_3)$ [mas/yr]	+0.0 ± 1.1	+0.5 ± 1.0	+1.2 ± 0.5

The Table describes three different absolute proper motion programmes and shows preliminary link results with H 30 data. The number of Bonn and Potsdam link fields will be increased (to 15 and 50, respectively) so that the influence of possible systematic effects - not represented by the formal errors $\sigma(\omega_i)$ - can be further reduced. We expect to provide an accuracy of the final link of the Hipparcos proper motions of better than 1 mas/yr, competitive with other link programmes (Lick/Yale, VLBI, HST).