

RELATIONS ON TOPOLOGICAL SPACES: URYSOHN'S LEMMA

Y.-F. LIN¹

(Received 7 December 1965, revised 19 August 1966)

1. Introduction and Preliminaries

Let X be a topological space equipped with a binary relation R ; that is, R is a subset of the Cartesian square $X \times X$. Following Wallace [5], we write

$$\begin{aligned}xR &= \{y : (x, y) \in R\}, & Rx &= \{y : (y, x) \in R\}, \\ RA &= \cup \{Rx : x \in A\} \text{ and } AR = \cup \{xR : x \in R\}.\end{aligned}$$

Deviating from [7], we shall follow Wallace [4] to call the relation R *continuous* if $RA^* \subset (RA)^*$ for each $A \subset X$, where $*$ designates the topological closure. Borrowing the language from the Ordered System, though our relation R need not be any kind of order relation, we say that a subset S of X is *R -decreasing* (*R -increasing*) if $RS \subset S(SR \subset S)$, and that S is *R -monotone* if S is either R -decreasing or R -increasing. Two R -monotone subsets are of the *same type* if they are either both R -decreasing or both R -increasing.

DEFINITION 1. A topological space X equipped with a relation R is said to be *R -normal* if, and only if, A_1 and A_2 are two disjoint closed subsets of X such that either A_1 is R -decreasing or A_2 is R -increasing then there exist two disjoint R -monotone open subsets U_1 and U_2 of X such that $A_1 \subset U_1$, $A_2 \subset U_2$ and that U_1 is R -decreasing and U_2 is R -increasing.

Our definition of R -normality agrees with that of strong normality of Ward's [6].

It should be noted that the *trivial* relation, $\Delta = \{(x, x) : x \in X\}$, is continuous and that a normal space is a particular R -normal space in which R is trivial. Recall that a family of subsets of X is *point-finite* (*locally finite*) if every point of X belongs to (has a neighborhood that meets) at most a finite number of sets in the family; in particular, all finite, all star finite [2], and all locally finite families are point-finite.

¹ Presented to the 71st Summer Meeting of the American Mathematical Society, September 1, 1966. This work was supported by the Research Council of the University of South Florida. The author is indebted to the referee for pointing out some errors in the earlier version of this paper.

The purpose of this note is to prove the following generalization of the famous Urysohn's Lemma.

THEOREM. *Let X be a topological space equipped with a continuous relation R such that X is R -normal, F an R -increasing (R -decreasing) closed subset of X , and $\{U_\alpha : \alpha \in A\}$ a locally finite family of R -increasing (R -decreasing) open sets in X such that $\cup \{U_\alpha : \alpha \in A\} \supset F$. Then there exists a family $\{f_\alpha : \alpha \in A\}$ of continuous functions on X with values in $[0, 1]$ such that*

- (i) *if $(x, y) \in R$, then $\sum_{\alpha \in A} f_\alpha(x) \leq \sum_{\alpha \in A} f_\alpha(y)$;*
- (ii) *$\sum_{\alpha \in A} f_\alpha(x) = 1$ ($\sum_{\alpha \in A} f_\alpha(x) = 0$) for all $x \in F$; and*
- (iii) *for each $\alpha \in A$, $f_\alpha(x) = 0$ ($f_\alpha(x) = 1$) for all $x \in X \setminus U_\alpha$.*

We are inspired by [3], [4], [5], and [6].

2. Proof of the main theorem

Let us record a number of readily established facts and some useful lemmas. *In all that follows, let X be always a topological space equipped with a relation R .*

2.1 *A subset A of X is R -decreasing (R -increasing) if, and only if, $X \setminus A$ is R -increasing (R -decreasing) [6].*

2.2 *If A is an R -decreasing (R -increasing) subset of X and if R is continuous, then A^* and A^0 are R -decreasing (R -increasing), where 0 denotes interior.*

2.3 *If $\{A_\alpha\}$ is a family of R -decreasing (R -increasing) subsets of X , then*

$$\cup_\alpha A_\alpha \text{ and } \cap_\alpha A_\alpha$$

are R -decreasing (R -increasing) [6].

2.4 *The empty subset \square and the whole space X are both R -decreasing as well as R -increasing.*

LEMMA 1. (Urysohn-Nachbin-Ward). *The space X is R -normal if, and only if, A and B are two disjoint closed sets in X such that either A is R -decreasing or B is R -increasing, then there exists a continuous function h on X with values in $[0, 1]$ such that $h(x) \leq h(y)$ whenever $(x, y) \in R$ and that $h(x) = 0$ for all $x \in A$ and $h(x) = 1$ for all $x \in B$.*

PROOF. To prove the sufficiency, let $U = \{x \in X | 0 \leq h(x) < \frac{1}{2}\}$ and $V = \{x \in X | \frac{1}{2} < h(x) \leq 1\}$, then $U \cap V = \square$, $A \subset U$ and $B \subset V$. By the continuity of h , one sees that U and V are open. Finally, since h satisfies

the property that $(x, y) \in R$ implies $h(x) \leq h(y)$, U is R -decreasing and V is R -increasing. Thus, X is R -normal.

The necessity part of the proof may be found in Ward [6, page 363].

LEMMA 2. *Let R be a continuous relation on X , then X is R -normal if, and only if, for any R -decreasing (R -increasing) closed set A contained in any open set $U \subset X$, there exists an R -decreasing (R -increasing) open set $U_0 \subset X$ such that $A \subset U_0 \subset U_0^* \subset U$.*

PROOF. If X is R -normal, A an R -decreasing (R -increasing) closed set contained in an open set U , then A and $X \setminus U$ are two disjoint closed sets in X such that A is R -decreasing (R -increasing). The R -normality of X implies that there exists an R -increasing (R -decreasing) open set V containing $X \setminus V$ and missing an R -decreasing (R -increasing) neighborhood of A . Taking $U_0 = X \setminus V^*$, one shows by (2.1) and (2.2) that U_0 fulfills the requirements.

Conversely, if A_1 and A_2 are two disjoint closed subsets of X such that A_1 is R -decreasing; the case where A_2 is R -increasing may be argued similarly. Then, since A_1 is contained in the open set $X \setminus A_2$, there exists an R -decreasing open set U_0 such that $A_1 \subset U_0 \subset U_0^* \subset X \setminus A_2$. Taking $U_1 = U_0$ and $U_2 = X \setminus U_0^*$, then U_1 and U_2 are two disjoint open sets such that $A_1 \subset U_1$, $A_2 \subset U_2$ and that U_1 is R -decreasing and U_2 is R -increasing, by (2.1) and (2.2). Thus, X is R -normal.

We say that a family of subsets of X is R -increasing (R -decreasing, R -monotone) if each of its member is an R -increasing (R -decreasing, R -monotone) subset of X ; thus, an R -increasing open cover of X is a cover of X by R -increasing open sets.

LEMMA 3. *Let X be an R -normal space such that R is continuous, $\mathcal{U} = \{U_\alpha : \alpha \in A\}$ a point-finite, R -increasing (R -decreasing) open cover of X , then there exists an R -increasing (R -decreasing) open cover $\mathcal{V} = \{V_\alpha : \alpha \in A\}$ ² of X such that $V_\alpha^* \subset U_\alpha$, for all $\alpha \in A$.*

PROOF. We shall prove only the case where \mathcal{U} is R -increasing, the case where \mathcal{U} is R -decreasing may be proved similarly.

By the well-ordering principle, let the indexing set A be well-ordered by a certain well-ordering denoted by \leq . Let $\mathcal{V}^{(o)} = \mathcal{U}$ where o denotes a symbol not in A ; define $o < \alpha$ for all $\alpha \in A$. For each β in A , we shall construct inductively an R -increasing open cover $\mathcal{V}^{(\beta)} = \{V_{\alpha,\beta} : \alpha \in A\}$ satisfying

² It should be remarked that we allow some of the V_α 's to be empty [cf. (2.4)]. In fact, some of the V_α 's may have to be empty even when all U_α are nonempty. The referee has contributed the following example: let $X = \{x_1, x_2\}$ with open sets: X , $\{x_2\}$ and the empty set \square . Let X be equipped with the trivial relation $R = \Delta$, then X is R -normal. Consider $\mathcal{U} = \{U_1, U_2\}$ where $U_1 = \{x_2\}$ and $U_2 = X$; in this case, the only possible choice of $\mathcal{V} = \{V_1, V_2\}$ is $V_1 = \square$ and $V_2 = X$.

(i) $V_{\alpha,\beta}^* \subset U_\alpha$ for all $\alpha \leq \beta$, and (ii) $V_{\alpha,\beta} = U_\alpha$ for all $\alpha > \beta$. Suppose for some $\gamma \in A$ that all $\mathcal{V}^{(\beta)}$ with $\beta < \gamma$ have been constructed, then we shall define $\mathcal{V}^{(\gamma)}$ as follows: define

$$V_{\alpha,\gamma} = \begin{cases} V_{\alpha,\alpha} & \text{for all } \alpha < \gamma, \\ U_\alpha & \text{for all } \alpha > \gamma. \end{cases}$$

To define $V_{\gamma,\gamma}$, we first observe that

$$U_\gamma \cup [\cup \{V_{\alpha,\gamma} : \alpha \in A \ \& \ \alpha \neq \gamma\}] = X$$

and hence $X \setminus U_\gamma$ and $X \setminus \{V_{\alpha,\gamma} : \alpha \in A \ \& \ \alpha \neq \gamma\}$ are two disjoint (R -monotone) closed sets. By R -normality of X , there exist two disjoint open sets S and T such that $X \setminus U_\gamma \subset S$ and $T \supset X \setminus \cup \{V_{\alpha,\gamma} : \alpha \in A \ \& \ \alpha \neq \gamma\}$. Since $X \setminus U_\gamma$ is R -decreasing, by Lemma 2, there exists an R -decreasing open set Q such that $X \setminus U_\gamma \subset Q \subset Q^* \subset S$. Now let $V_{\gamma,\gamma} = X \setminus Q^*$, then by (2.1) and (2.2) $V_{\gamma,\gamma}$ is R -increasing. We have $V_{\gamma,\gamma}^* \subset X \setminus Q \subset U_\gamma$. Furthermore, since Q^* and $X \setminus \cup \{V_{\alpha,\gamma} : \alpha \in A \ \& \ \alpha \neq \gamma\}$ are disjoint, we have

$$\begin{aligned} \cup \{V_{\alpha,\alpha} : \alpha \in A\} &= V_{\gamma,\gamma} \cup [\cup \{V_{\alpha,\gamma} : \alpha \in A \ \& \ \alpha \neq \gamma\}] \\ &= (X \setminus Q^*) \cup [\cup \{V_{\alpha,\gamma} : \alpha \in A \ \& \ \alpha \neq \gamma\}] = X. \end{aligned}$$

Thus, by the transfinite induction, $\mathcal{V}^{(\gamma)}$ is well-defined for every $\gamma \in A$. It is to be noted that the γ^{th} cover $\mathcal{V}^{(\gamma)}$ contains an initial segment of every cover $\mathcal{V}^{(\beta)}$ for $\beta < \gamma$. Let $\mathcal{V} = \{V_\alpha : \alpha \in A\}$ be defined by $V_\alpha = V_{\alpha,\alpha}$ for all $\alpha \in A$. Then, it needs only to be proved that \mathcal{V} covers X . To this end, let x be an arbitrary element in X , then by the point-finiteness of \mathcal{U} there exist at least one and at most finitely many $U_{\alpha_1}, \dots, U_{\alpha_n}$ containing x . Let $\alpha_0 = \max \{\alpha_1, \dots, \alpha_n\}$, under the well-ordering we have introduced at the very beginning. Consequently, $x \notin U_\gamma$ for all $\gamma > \alpha_0$; but since $\mathcal{V}^{(\alpha_0)}$ covers X , there exists an $\beta \leq \alpha_0$ such that $x \in V_{\beta,\alpha_0} = V_{\beta,\beta} = V_\beta \in \mathcal{V}$.

PROOF OF THEOREM. Since F is an R -increasing (R -decreasing) closed subset of the R -normal space X , the set F equipped with the relative topology and the relation $R_F = R \cap (F \times F)$ forms an R_F -normal space, and the family $\{U_\alpha \cap F; \alpha \in A\}$ is a point-finite R_F -increasing (R_F -decreasing) open (in F) cover of F . By Lemma 3, there exists an R_F -increasing (R_F -decreasing) open (in F) cover $\{V_\alpha : \alpha \in A\}$ of F such that $V_\alpha^* \subset U_\alpha \cap F \subset U_\alpha$ for all $\alpha \in A$. Since each V_α is an R_F -increasing (R_F -decreasing) subset of the R -increasing (R -decreasing) subset F of X , each V_α is R -increasing (R -decreasing) in X and hence, by (2.2), each V_α^* is R -increasing (R -decreasing) in X . Now, for each $\alpha \in A$, applying Lemma 1 to two disjoint R -monotone closed sets V_α^* and $X \setminus U_\alpha$ ³ in X , we have a continuous function

³ If $V_\alpha^* = \square$, regardless of $X \setminus U_\alpha = \square$ or otherwise, one may choose, for instance, the constant function $h_\alpha(x) = 0$ (resp. $h_\alpha(x) = 1$) for all $x \in X$; if $V_\alpha^* \neq \square$ and $X \setminus U_\alpha = \square$, then one chooses, for instance, the constant function $h_\alpha(x) = 1$ (resp. $h_\alpha(x) = 0$) for all $x \in X$.

$h_\alpha : X \rightarrow [0, 1]$ such that

$$\begin{aligned} h_\alpha(x) &\leq h_\alpha(y) \text{ whenever } (x, y) \in R, \\ h_\alpha(x) &= 1 \text{ (} h_\alpha(x) = 0 \text{) for all } x \in V_\alpha^*, \text{ and} \\ h_\alpha(x) &= 0 \text{ (} h_\alpha(x) = 1 \text{) for all } x \in X \setminus U_\alpha. \end{aligned}$$

For each $\alpha \in A$, define a function $f_\alpha : X \rightarrow [0, 1]$ by

$$f_\alpha(x) = h_\alpha(x) / \max \left\{ \sum_{\alpha \in A} h_\alpha(x), 1 \right\}$$

for all x in X . To show that each f_α is continuous, it suffices to show that the function $\sum_{\alpha \in A} h_\alpha : X \rightarrow [0, \infty)$ is continuous: for each x in X , by the local finiteness of $\{U_\alpha : \alpha \in A\}$, there exists an open neighborhood N_x of x such that $N_x \cap U_\alpha \neq \emptyset$ for at least one and at most finitely many α 's, say $\alpha_1, \alpha_2, \dots, \alpha_m$; therefore, for any open set G in $[0, \infty)$,

$$\left(\sum_{\alpha \in A} h_\alpha \right)^{-1}(G) \cap N_x = (h_{\alpha_1} + \dots + h_{\alpha_m})^{-1}(G) \cap N_x$$

which is open, because $h_{\alpha_1} + \dots + h_{\alpha_m} : X \rightarrow [0, \infty)$, as a sum of finitely many continuous functions, is continuous; and hence,

$$\left(\sum_{\alpha \in A} h_\alpha \right)^{-1}(G) = \bigcup_{x \in X} \left[\left(\sum_{\alpha \in A} h_\alpha \right)^{-1}(G) \cap N_x \right],$$

as a union of open sets, is open. Finally, by a routine verification, one verifies that the family $\{f_\alpha : \alpha \in A\}$ of continuous functions satisfies the conditions (i), (ii) and (iii) stated in the theorem.

Since every normal space is R -normal, where R is the (continuous) trivial relation Δ , and every finite family of open sets is locally finite, therefore we have the following Dieudonné's partition of unity.

COROLLARY (Dieudonné [1]). *Let X be a normal space, F a closed subset of X , and U_1, U_2, \dots, U_n open sets such that $\bigcup_{k=1}^n U_k \supset F$. Then there exist continuous functions h_1, \dots, h_n on X with values in $[0, 1]$ such that*

- (i) $\sum_{k=1}^n h_k(x) = 1$ for all $x \in F$;
- (ii) $h_k(x) = 0$ for all $x \in X \setminus U_k$ and for $k = 1, 2, \dots, n$.

I am grateful to Dr. Frank L. Cleaver for his encouragement.

References

- [1] J. Dieudonné, 'Sur les fonctions continues numériques définies dans un produit de deux espaces compacts', *C.R. Acad. Sci. Paris* 205 (1937), 593–595.
- [2] J. L. Kelley, *General topology*, (New York, 1955).
- [3] L. Nachbin, *Topologia e ordem*, (Chicago, 1950).

- [4] A. D. Wallace, *Relations on topological spaces* (Proc. Symp. on General Topology and its Relations to Modern Analysis and Algebra (Prague 1961), 356–360.)
- [5] A. D. Wallace, *Relation-theory*, Lecture Notes, (Univ. of Fla. 1963–1964).
- [6] L. E. Ward, Jr., 'Binary relations in topological spaces', *Anais Acad. Basil. Ci.* 36 (1954), 357–373.
- [7] L. E. Ward, Jr., 'Partially ordered topological spaces', *Proc. Amer. Math. Soc.* 5 (1954), 144–161.

The University of South Florida
Tampa, Florida