

Adolescents' diet quality in relation to their relatives' and peers' diet engagement and encouragement: the Healthy Lifestyle in Europe by Nutrition in Adolescence (HELENA) study

Jérémy Vanhelst^{1,*}, Laurent Béghin¹, Elodie Drumez², Alain Duhamel², Stefaan De Henauw³, Jonatan R Ruiz⁴, Anthony Kafatos⁵, Yannis Manios⁶, Kurt Widhalm⁷, Béatrice Mauro⁸, Michael Sjöström⁹, Mathilde Kersting¹⁰ and Frédéric Gottrand¹

¹Univ. Lille, Inserm, CHU Lille, U995 – LIRIC – Lille Inflammation Research International Center, CIC 1403 – Centre d'investigation clinique, Lille F-59000, France: ²Univ. Lille, CHU Lille, EA 2694 – Santé publique: épidémiologie et qualité des soins, Lille, France: ³Department of Movement and Sport Sciences, Ghent University, Ghent, Belgium: ⁴PROmoting FITness and Health through physical activity research group (PROFITH), Department of Physical Education and Sports, Faculty of Sport Sciences, University of Granada, Granada, Spain: ⁵Department of Social Medicine, Preventive Medicine and Nutrition, University of Crete School of Medicine, Heraklion, Crete, Greece: ⁶Department of Nutrition and Dietetics, Harokopio University, Athens, Greece: ⁷Department of Pediatrics, Private Medical University Salzburg, Salzburg, Austria: ⁸Agricultural Research Council – Research Center on Food and Nutrition – CRANUT (formerly INRAN), Rome, Italy: ⁹Unit for Preventive Nutrition, Department of Biosciences and Nutrition, Karolinska Institutet, Huddinge, Sweden: ¹⁰Forschungsinstitut fuer Kinderernaehrung, Rheinischen Friedrich-Wilhelms-Universitaet Bonn, Dortmund, Germany

Submitted 11 October 2017: Final revision received 30 May 2018: Accepted 14 June 2018: First published online 10 August 2018

Abstract

Objective: To examine the associations between adolescents' diet quality and their perceived relatives' and peers' diet engagement and encouragement.

Design: Cross-sectional study performed in European countries. Diet quality was scored using the Diet Quality Index for Adolescents (DQI-A) based on four components: quality, diversity, balance and meal frequency. Perceived diet quality engagement and perceived encouragement of the relatives/peers were assessed using the questions 'How healthy is each of the following persons' diet?' and 'How often does each of the following persons encourage you to eat a healthy diet?'

Setting: Vienna, Ghent, Lille, Athens, Heraklion, Pecs, Rome, Dortmund, Zaragoza and Stockholm.

Subjects: Healthy adolescents (*n* 2943).

Results: The perceived engagement level of the mother, father and sister was each positively associated with the DQI-A ($P < 0.05$). A positive association was found for the perceived engagement level of siblings, father and mother with all specific components ($P < 0.05$). DQI-A was negatively associated with the perceived encouragement level from a best friend and positively associated with the encouragement level of the mother and father ($P < 0.05$). Diversity, balance and quality components were positively associated with the perceived encouragement level from the mother and father ($P < 0.05$), whereas the best friend's perceived encouragement was negatively associated with the meal frequency component ($P < 0.01$).

Conclusions: These findings highlight the role of social engagement and encouragement of relatives and peers in adolescents' diet quality. Intervention or promotion programmes aimed at enhancing diet quality in adolescents should target both family and peers.

Keywords
Youth
Assessment
Nutrition
Family
Epidemiological study

Adolescence is an important period in life that includes multiple physiological and psychological changes that have a considerable effect on dietary habits^(1,2). Unhealthy food consumption patterns during childhood and adolescence are linked with both the occurrence of obesity in

youth and the later risk of developing diseases such as cancer, obesity and CVD in adulthood⁽³⁾.

Dietary habits are influenced by individual, social and environmental factors, including food choice decisions, food choice motivations, religious adherence, food

cravings, taste, hunger, time and effort required for food preparation and consumption, cost, body image and socio-economic status^(4–8). Dietary habits are also influenced by cultural traditions, which differ between countries^(9,10). Family and peers are considered important sociocultural influences that have a strong impact on dietary habits during adolescence^(11–17). Previous studies have consistently demonstrated the importance of parents to healthy eating habits during adolescence, specifically vegetable and fruit consumption^(13,16,17). Previous studies of both encouragement and engagement have also found that friends influence the intake of healthy foods such as vegetables, energy drinks, snacks, desserts, fruits, whole grains and biscuits^(13–15). However, previous studies have not included the influence of siblings' encouragement and engagement in terms of the diet. Moreover, the aim of previous research was to compare the social influence of parents and friends on eating attitudes of adolescents by focusing specifically on particular food groups^(11–17). To our knowledge, no previous studies have examined the influence of family and peers on diet quality taking account into quality, diversity, equilibrium and meal frequency in adolescents.

Therefore, the aim of the present study was to examine the associations between adolescents' diet quality and their perceived relatives' and peers' (father, mother, brother, sister and best friend) diet engagement and encouragement.

Methods

Study design

The present study is a secondary data analysis of the Healthy Lifestyle in Europe by Nutrition in Adolescence (HELENA) Study (www.helenastudy.com) performed in European adolescents (2006–2007). The aim of the HELENA Study was to obtain a broad range of standardized, reliable and comparable nutrition- and health-related data from a random sample of European adolescents aged 12.5–17.5 years.

The random selection of schools and classes was performed centrally. The first step of the recruitment strategy consisted of telephone contact with the director/principal of the school. During the call, a meeting with the director/principal and main/principal teachers of selected classes was organized to present the study aims and procedures and obtain consent to participate. The second step consisted of a meeting with adolescents from selected classes and their main/principal teacher. During this meeting, the study aims, procedures and tests were explained. Information and consent forms were then distributed, and the adolescents were asked to return the written/signed consent form (including the signatures of the adolescent and both parents) within a maximum of two weeks after the meeting. Table 1 presents an overview of the participation

rate of the different sampling units for the whole study and for each centre individually. In total, 3528 adolescents were included in the HELENA Study, 83% of whom completed the dietary habits questionnaire and were therefore included in the present study. The participants' characteristics are presented in Table 2. No significant differences were observed between the included and excluded adolescents' characteristics.

The local ethics committee for each country approved the HELENA Study, and all procedures were performed in accordance with the ethical standards of the Helsinki Declaration of 1975, as revised in 2008, and the European Good Clinical Practices.

Measurements

Assessment of relatives' and peers' diet engagement and encouragement

A self-administered questionnaire was used to assess healthy diet determinants. A paper version of the questionnaire was administered in a classroom under the supervision of a HELENA fieldworker. Two questions on engagement and encouragement were extracted for the present study⁽¹⁸⁾. The adolescents were asked about the perceived diet quality engagement of their relatives and peers (father, mother, brother(s), sister(s) and best friend(s)) using the following question: 'How healthy is each of the following persons' diet: father, mother, sister(s), brother(s), best friend(s)?' The adolescents' perceived engagement of their relatives and peers was classified as low if the answer to the question was 'very unhealthy' or 'not very healthy', medium if 'average' or 'quite healthy', and high if 'very healthy'. The adolescents were also asked about the perceived diet encouragement provided by their relatives and peers using the following question: 'How often does each of the following persons encourage you to eat a healthy diet: father, mother, sister(s), brother(s), best friend(s)?' The answers were classified as low if the answer to the question was 'not at all' or 'not much', medium if 'sometimes' or 'often', and high if 'very often'. These questions regarding perceived relatives' and peers' diet engagement and encouragement were extracted from a healthy diet determinants questionnaire that has been found to be valid and reliable⁽¹⁸⁾.

Dietary habits

Dietary intake was assessed by two non-consecutive 24 h recalls performed on two convenient weekdays one week apart. The 24 h recalls were recorded using the self-administered, computer-based HELENA Dietary Intake Assessment Tool (HELENA-DIAT), which has been validated in European adolescents⁽¹⁹⁾. The HELENA-DIAT tool is based on intake assessments at six meal occasions (breakfast, morning snack, lunch, afternoon snack, evening meal and evening snack) on the previous day. Trained dietitians assisted the adolescents to complete the

Table 1 Number of approached/participating classes and adolescents in the Healthy Lifestyle in Europe by Nutrition in Adolescence (HELENA) Study†

	Centre									
	Athens	Dortmund	Ghent	Heraklion	Lille	Pecs	Rome	Stockholm	Vienna	Zaragoza
No. of eligible schools in the city	82	55	43	22	40	12	290	25	347	83
No. of schools approached/participating	17/10	14/11	11/9	11/10	13/12	8/7	18/10	14/10	23/13	16/12
No. of classes approached/participating	14/14	23/23	20/19	22/20	19/18	24/14	24/22	25/23	35/19	26/23
No. of adolescents approached in all approached classes	458	603	429	429	538	720	420	645	870	597
No. of adolescents approached in all participating classes/no. of adolescents participating	458/370	603/515	413/347	400/340	508/308	420/401	430/339	535/377	536/427	537/441
Adolescents included in HELENA Study										
<i>n</i>	321	476	336	284	287	394	304	341	403	382
%	70%‡	79%‡	78%‡	66%‡	53%‡	55%‡	65%‡	53%‡	63%‡	64%‡

†Data were collected from 2006 to 2007.

‡Percentage calculated to reflect the ratio of selected adolescents for statistical analysis to adolescents approached in all approached classes.

24 h recalls when needed. To calculate energy and nutrient intakes, data from HELENA-DIAT were linked to the German Food Code and Nutrient Database (Bundeslebensmittelschlüssel, version II.3.1)⁽²⁰⁾. The Multiple Source Method was used to estimate the usual energy, nutrient and food intakes.

The Diet Quality Index for Adolescents (DQI-A) is composed of four components – quality, diversity, equilibrium and meal frequency – that were previously validated in the HELENA population^(21–24). Daily diet was divided into nine recommended food groups: (i) water; (ii) bread and cereal; (iii) potatoes and grains; (iv) vegetables; (v) fruits; (vi) milk products; (vii) cheese; (viii) meat, fish and substitutes; and (ix) fats and oils. Dietary quality indicated whether an adolescent made optimal food quality choices within a food group and was represented by a ‘preference group’ (i.e. the healthiest foods: cereal/brown bread, fresh fruit and fish), an ‘intermediate group’ (e.g. white bread, minced meat) and a ‘low-nutrient, energy-dense group’ (i.e. the unhealthiest foods: soft drinks, sweet snacks and chicken nuggets) using predefined criteria. The dietary quality score was then calculated by multiplying the amount of the food consumed (in grams) by a weighting factor (+1 for the preference group, 0 for the intermediate group and –1 for the low-nutrient, energy-dense group) and dividing by the total amount of food (in grams). The diet quality score was expressed as a percentage, meaning that it could vary between –100 and 100%. The diversity component corresponds to the degree of variation in the diet. The score was obtained by assigning 1 point for each food group that had at least one serving at the preference level, dividing by 9 (which represents the maximum score), and then expressed as a percentage between 0 and 100%. Dietary equilibrium was calculated as the difference between the adequacy component (the percentage of food groups with intake above the minimum recommended value) and the excess component (the percentage of food groups exceeding the upper level of the recommended intake) and ranged between 0 and 100%. Meal frequency was scored as 0 when no food was consumed and 1 when some food was consumed at each of the three main meal occasions. The scores for the three occasions were summed and expressed as a percentage; the possible scores were thus 0% (no consumption at any of the main meals), 33% (consumption at only one main meal), 66% (consumption at two main meals) and 100% (consumption at all three main meals).

The four DQI-A components are presented as percentages. The quality component ranged from –100 to 100%, whereas the diversity, equilibrium and meal frequency components ranged from 0 to 100%. The DQI-A was computed as the arithmetic mean of these four components; hence, the DQI-A ranged from –25 to 100%, with higher scores reflecting a higher-quality diet. The score was calculated for each day and the mean daily score was taken as the individual’s overall index.

Table 2 Characteristics of the study population of adolescents (*n* 2943) aged 12.5–17.5 years from ten European cities, Healthy Lifestyle in Europe by Nutrition in Adolescence (HELENA) Study

	Before imputation		After imputation
	Without missing DQI-A data	With missing DQI-A data	
<i>n</i>	2943	585	3528
Sex (% male)	47.2	50.3	47.7
Age (years)			
Mean	14.8	14.5*	14.7
SD	1.2	1.2	1.2
Height (cm)			
Mean	166.2	163.8*	165.8
SD	9.2	8.7	9.1
Body mass (kg)			
Mean	59.0	59.9	59.1
SD	12.7	12.9	12.7
BMI Z-score			
Mean	0.32	0.62*	0.37
SD	0.9	0.9	0.9
Nutritional status (% UW/% NW/% OW/% O)†	6.7/72.0/16.4/4.9	3.1/64.6/22.9/9.4*	6.1/70.8/17.5/5.6
Father's education level (% I/% II/% III)‡	37.4/27.0/35.6	39.9/32.0/28.1*	38.7/27.5/33.8
Mother's education level (% I/% II/% III)‡	34.0/30.9/35.1	40.2/33.9/25.9*	35.5/31.0/33.5

DQI-A, Diet Quality Index for Adolescents.

**P* < 0.05 for comparison between the two samples, without and with missing data on DQI-A.

†Nutritional status: underweight (UW), normal weight (NW), overweight (OW), obese (O).

‡Education level: lower education (I); higher secondary education (II); higher education or university degree (III).

Participants' characteristics

Body weight was measured with the participant wearing light clothes and without shoes to the nearest 0.1 kg using an electronic scale (model 871; SECA, Hamburg, Germany). Height was measured without shoes to the nearest 0.1 cm using a telescopic height-measuring instrument (model 225; SECA). BMI was calculated as [weight (kg)]/[height (m)]². The nutritional status was assessed using the International Obesity Task Force scale⁽²⁵⁾. An extended and detailed manual of operations was designed for and thoroughly read by every researcher involved in fieldwork before the data collection started⁽²⁶⁾. In addition, a workshop training week was carried out before the study began to standardize and harmonize the data collection methods. The instructions given to the participants for every measurement were standardized for all cities and translated into the local language.

Parental educational level was classified into one of four categories using a specific questionnaire adapted from the International Standard Classification of Education (ISCED; <http://www.uis.unesco.org/Library/Documents/isced97-en.pdf>). Parental educational level was scored as 1 for primary and lower education (levels 0, 1 and 2 in the ISCED classification); 2 for higher secondary (levels 3 and 4 in the ISCED classification); and 3 for tertiary (levels 5 and 6 in the ISCED classification).

Statistical analyses

The data are presented as percentages for qualitative variables and as means and standard deviations for quantitative variables. Normality of distribution was checked graphically and by using the Shapiro–Wilk test.

To assess the potential bias related to missing or incomplete data for the DQI-A, the main adolescent characteristics were compared between adolescents with and without DQI-A data using Student's *t* test for quantitative variables, the χ^2 test for categorical variables and the Mantel–Haenszel trend test for ordered categorical variables (Table 1).

We examined the associations between the DQI-A (overall index and each component) and each perceived relative's and peer's diet engagement and encouragement levels using linear mixed models adjusted for prespecified confounding factors, including age, sex and parental educational level as fixed effects, and city, city × school and city × school × class as random effects^(21,27,28). The adjusted means for the DQI-A and their standard errors were calculated using the least-square means. Because the perceived relatives' and peers' diet engagement or encouragement levels were classified into three ordered levels, we used linear contrasts to perform the trend test. Comparisons of overall DQI-A between the ten perceived relatives' and peers' diet engagement or encouragement levels (main objective) were adjusted for multiple comparisons using the false discovery rate controlling method⁽²⁹⁾.

To avoid case deletion in the analyses, missing data were imputed by multiple imputations using the regression-switching approach (chained equations with *m* = 20 imputations obtained using R statistical software version 3.03)⁽³⁰⁾. The imputation procedure was performed under the missing-at-random assumption using all adolescents' characteristics, relatives' and peers' diet engagement and encouragement levels and DQI-A components with the predictive mean-matching method for

quantitative variables, logistic regression for binary variables and ordinal logistic regression for ordered categorical variables. Rubin's rules were used to combine the estimates derived from multiple imputed data sets⁽³¹⁾. We performed a key subgroup analysis according to sex for the associations of overall DQI-A and meal frequency component with each perceived relative's and peer's diet engagement and encouragement levels. Inclusion of the corresponding interaction term into the multivariable linear mixed model was used to assess heterogeneity.

All statistical tests were done at the two-tailed α level of $P < 0.05$. Data were analysed using the statistical software package SAS version 9.3.

Results

Physical characteristics of the adolescents are presented in Table 2.

The adolescents' overall DQI-A score was positively and significantly associated with their perceived mother's, brother's and sister's diet engagement (Table 3). Having a high level of perceived mother's, brother's and sister's engagement resulted in 6, 5 and 4% higher diet quality scores, respectively, compared with the low level (Table 3). We found also significant positive associations between perceived sister's diet engagement and the adolescents' quality component (39.5 (SE 2.8) *v.* 43.2 (SE 1.8) *v.* 48.1 (SE 2.6) for low, medium and high, respectively; $P = 0.004$; +21.7%; Fig. 1). Similarly, a significant positive association was observed between perceived brother's diet engagement and adolescents' diversity component (72.4 (SE 1.1) *v.* 73.9 (SE 0.9) *v.* 76.2 (SE 1.3) for low, medium and high, respectively; $P = 0.003$; +5.2%); between perceived father's diet engagement and adolescents' balance (40.1 (SE 0.6) *v.* 41.0 (SE 0.3) *v.* 41.5 (SE 0.5)

for low, medium and high, respectively; $P = 0.047$; +3.5%) and diversity components (72.7 (SE 1.1) *v.* 73.6 (SE 0.9) *v.* 75.1 (SE 1.1) for low, medium and high, respectively; $P = 0.003$; +3.3%); and between perceived mother's diet engagement and adolescents' balance component (39.3 (SE 0.9) *v.* 40.8 (SE 0.3) *v.* 41.9 (SE 0.4) for low, medium and high, respectively; $P = 0.005$; +6.6%; Fig. 1). Similar findings were observed for the adolescents' diversity (71.2 (SE 1.4) *v.* 73.4 (SE 0.9) *v.* 75.3 (SE 0.9) for low, medium and high, respectively; $P = 0.005$; +5.7%) and meal frequency components (89.8 (SE 1.0) *v.* 91.7 (SE 0.7) *v.* 92.6 (SE 0.8) for low, medium and high, respectively; $P = 0.023$; +3.1%; Fig. 1) with perceived mother's diet engagement.

The overall DQI-A score was significantly negatively associated with the perceived best friend's encouragement and positively associated with the perceived father's and mother's encouragement (Table 4). Having a high level of perceived best friend's, father's and mother's encouragement resulted in 4.0% lower, and 4.4 and 4.4% higher, diet quality scores, respectively, compared with the low level (Table 4). The perceived father's encouragement level was positively and significantly associated with the quality, diversity and balance components (Fig. 2). The difference observed in the diet quality score between the low and high levels of perceived father's encouragement was 13.4, 3.9 and 3.2% for the quality, diversity and balance components, respectively (Fig. 2). Positive associations were also found between the perceived mother's encouragement and the quality, balance and diversity components. The difference observed in the diet quality score between the low and high levels of perceived mother's encouragement was 13.0, 4.0 and 5.5% for the quality, balance and diversity components, respectively (Fig. 2). Another positive association was found between the perceived sister's diet encouragement and the adolescents' balance component, with a difference of 4.6% in diet quality score

Table 3 Overall diet quality scores according to their relatives' and peers' diet engagement among adolescents (n 2943) aged 12.5–17.5 years from ten European cities, Healthy Lifestyle in Europe by Nutrition in Adolescence (HELENA) Study†

Relative/peer	Diet engagement	<i>n</i>	Mean DQI-A	SE	<i>P</i> ‡
Father	Low	475	62.07	0.98	0.077
	Medium	2447	62.21	0.68	
	High	606	63.92	0.84	
Mother	Low	203	60.13	1.27	0.008
	Medium	2413	62.21	0.67	
	High	912	63.74	0.77	
Brother	Low	757	61.84	0.81	0.008
	Medium	2443	62.36	0.68	
	High	328	64.93	1.06	
Sister	Low	458	61.03	1.00	0.032
	Medium	2622	62.53	0.67	
	High	448	63.45	0.99	
Best friend	Low	612	62.01	0.86	0.36
	Medium	2651	62.75	0.67	
	High	265	60.98	1.16	

DQI-A, Diet Quality Index for Adolescents.

†Number, adjusted mean, SE and *P* value for trend across relatives'/peers' diet engagement were calculated using linear mixed models including age, sex and parental educational level as fixed effects and city, city \times school and city \times school \times class as random effects after handling missing data by multiple imputation.

‡Controlled for multiple comparisons using the false discovery rate method.




Fig. 1 Diet quality components, scored using the Diet Quality Index for Adolescents (DQI-A), according to relatives' and peers' diet engagement (■, low; □, medium; ▒, high), among adolescents (*n* 2943) aged 12.5–17.5 years from ten European cities, Healthy Lifestyle in Europe by Nutrition in Adolescence (HELENA) Study. Values are means (with their standard errors represented by vertical bars) of each component, calculated using linear mixed models including age, sex and parental educational level as fixed effects and city, city × school and city × school × class as random effects after handling missing data by multiple imputation. *Adjusted *P* for trend < 0.05 across the relatives' engagement

Table 4 Overall diet quality scores according to their relatives' and peers' diet encouragement among adolescents (*n* 2943) aged 12.5–17.5 years from ten European cities, Healthy Lifestyle in Europe by Nutrition in Adolescence (HELENA) Study†

Relative/peer	Diet encouragement	<i>n</i>	Mean DQI-A	SE	<i>P</i> ‡
Father	Low	1097	61.52	0.73	< 0.001
	Medium	1791	62.46	0.68	
	High	640	64.23	0.87	
Mother	Low	612	60.77	0.82	< 0.001
	Medium	1865	62.50	0.67	
	High	1051	63.46	0.75	
Brother	Low	2412	62.46	0.68	0.23
	Medium	893	62.18	0.79	
	High	223	64.10	1.34	
Sister	Low	2194	62.10	0.67	0.32
	Medium	1030	63.10	0.79	
	High	304	63.36	1.26	
Best friend	Low	2235	62.71	0.69	0.040
	Medium	1069	62.51	0.78	
	High	224	60.20	1.25	

DQI-A, Diet Quality Index for Adolescents.

†Number, adjusted mean, SE and *P* value for trend across relatives'/peers' diet encouragement were calculated using linear mixed models including age, sex and parental educational level as fixed effects and city, city × school and city × school × class as random effects after handling missing data by multiple imputation.

‡Controlled for multiple comparisons using the false discovery rate method.

between the low and high perceived encouragement levels (Fig. 2). The perceived best friend's encouragement was negatively associated with the meal frequency

component, with a difference of 3.7% in the diet quality score between the low and high perceived encouragement levels (Fig. 2).




Fig. 2 Diet quality components, scored using the Diet Quality Index for Adolescents (DQI-A), according to relatives' and peers' diet encouragement (■, low; ▒, medium; □, high), among adolescents (n 2943) aged 12–17.5 years from ten European cities, Healthy Lifestyle in Europe by Nutrition in Adolescence (HELENA) Study. Values are means (with their standard errors represented by vertical bars) of each component, calculated using linear mixed models including age, sex and parental educational level as fixed effects and city, city \times school and city \times school \times class as random effects after handling missing data by multiple imputation. *Adjusted P for trend < 0.05 across the relatives' encouragement

The association between adolescents' overall DQI-A and perceived mother's diet encouragement was stronger in boys than in girls, although the heterogeneity test did not reach the level of significance (P for heterogeneity = 0.089). In boys, the adjusted mean DQI-A was 58.8 (SE 1.1) *v.* 61.0 (SE 0.9) *v.* 62.2 (SE 1.1) for the low, medium and high perceived mother's diet encouragement levels, respectively (P for trend = 0.002). By contrast, in girls, the adjusted mean DQI-A was 63.1 (SE 1.1) *v.* 64.1 (SE 0.8) *v.* 64.7 (SE 0.8) for the low, medium and high perceived mother's diet encouragement levels, respectively (P for trend = 0.12). We found no other significant heterogeneity based on the adolescents' sex (data not shown).

Discussion

Our study aimed to investigate the associations between adolescents' diet and their perceived relatives' and peers' (father, mother, brothers, sisters and best friends) diet engagement and encouragement. Since our study directly addressed adolescents, we only have information about perceived engagement and encouragement; relatives' and

peers' engagement and encouragement were not directly assessed. Although we acknowledge that this could have influenced our results, we believe that adolescents' perceptions influenced their own diet quality more than relatives' or peers' engagement and encouragement.

The main finding of our study is that both perceived relatives' diet engagement and encouragement were associated with the diet quality of the adolescents studied. However, the magnitude of the associations with the adolescents' DQI-A component scores varied according to the perceived parent's, sibling's or peer's diet engagement and encouragement. A strong positive association between the perceived mother's diet engagement and the adolescents' diet quality was found. This shows that mothers play a key role in family food choices, including adolescents' choices^(32,33), and is consistent with previous studies showing the importance of mothers to adolescents' meals⁽³⁴⁾. This finding also confirms that the perceived mother's engagement is associated with adolescents' diet quality. Moreover, it concurs with previous studies showing that mother-adolescent communication is more effective than father-adolescent communication in changing adolescents'

nutritional behaviour^(35,36). Our finding is also consistent with the results of the Healthy Eating Questionnaire, which showed that the mother is the family member most likely to promote healthy dietary habits⁽³⁷⁾. However, we also found an association between perceived fathers' encouragement and adolescents' diet quality. No previous studies have assessed the influence of brothers and sisters, and our data show for the first time a positive relationship between their perceived diet engagement or encouragement and adolescents' diet quality, balance and diversity components. This outcome shows the importance of siblings on the diet quality of the adolescent. Therefore, this suggests that intervention programmes that aim to enhance diet quality in adolescent populations might be more successful if parents and siblings are also included in the intervention. Our results concur with previous published studies showing that youth diet behaviours, particularly in obese paediatric patients, may be improved when parents attend and are directly involved with services and are provided with training in the skills required to support lifestyle modifications in accordance with expert guidelines^(38–41). In this context and from a practical point of view, primary care may play a major role in the improvement of parenting behaviours linked to child health^(42,43). Indeed, children and adolescents, most of the time accompanied by their parents, regularly access primary care where specialists or generalist physicians are present. Even if health-care providers report having inadequate time and a lack of expertise and resources to effectively work with parents and provide key messages regarding a healthy lifestyle, attempting to implement a specific time for discussion with parents and siblings during primary care visits remains important⁽⁴⁴⁾. This point of view is supported by several committees' recommendations regarding the prevention and treatment of youth overweight and obesity and the promotion of family-centred interventions in primary care^(45,46). Caregivers' policies should be discussed and new ways to address child and family care should be created for care providers such as paediatricians, family physicians, nurse practitioners and physicians' assistants. Concerning the roles of siblings, few existing preventive interventions target sibling relationships⁽⁴⁷⁾. Therefore, clinicians should also consider offering specific sessions for siblings that focus on healthy eating habits and instruction regarding how to promote and reinforce these habits among their siblings.

Most of the previous studies of the influence of relatives and peers have focused on the dietary behaviours of girls, and few studies have also included boys^(48,49). A significant difference between girls and boys was found only for the association between adolescents' DQI-A score and perceived mother's diet encouragement. In contrast to the results of previous studies of dietary behaviours, we found that girls' diet quality did not correlate with the perceived mother's diet encouragement⁽⁴⁸⁾. Indeed, it has been

shown that weight-control behaviours among young girls are modelled partially on their mothers' behaviours^(48,50,51). In the present study, we focused on diet quality components, but not directly on weight control.

During the transition from childhood to adolescence, children decrease the time spent with parents and spend more time alone and/or with friends⁽⁴⁹⁾. One unexpected finding of our study is the negative association between perceived encouragement of peers to eat healthily and adolescent unhealthy food consumption. This also contrasts with a recent study showing that friends' unhealthy food consumption was associated with an individual's unhealthy food consumption, although that study examined consumption rather than encouragement⁽⁵²⁾. One possible explanation is that those adolescents with unhealthy food consumption are encouraged by their peers to eat more healthily, independently of their peers' food consumption habits. Differences in our study in the associations between the adolescents' DQI-A scores and the perceived relatives' (positive association) and perceived peers' (negative association) diet engagement might reflect a better awareness of healthy lifestyle in adults than in adolescents. However, the influence of the relatives may also have adverse effects in the medium term⁽⁵³⁾. Indeed, if perceived relatives' engagement or encouragement is too important, it could lead to eating disorders and have a negative impact on future health. Several studies have highlighted concerns about the effectiveness of their role in dieting and the potential for increasing the risk of unintentional weight gain, disordered eating and eating disorders^(53–57).

The strengths of the present study are the large sample size of adolescents with sex-specific information in ten European cities, the use of standardized procedures, the inclusion of many confounding factors in the analyses and the strong methodology for assessing dietary habits⁽⁵⁸⁾. The limitations of the study include the cross-sectional and observational design to examine the associations, which cannot be interpreted to reflect causal relationships. The proxy report of the parent's, sibling's and peer's diet engagement and encouragement is another limitation that could lead to misclassification. Moreover, we cannot rule out bias because of the estimated values for missing data, as the multiple-imputation procedure to replace missing values with a set of plausible values was done under a missing-at-random assumption. Finally, in the present study, we found that the mean differences between variables was low, which raises the question of their clinical significance.

Conclusion

In conclusion, our findings highlight the role of social encouragement and engagement in adolescents' diet quality. Implementing intervention or promotion programmes

that aim to encourage a healthy diet in adolescents might be more successful if the family and peers are also targeted. Indeed, interventions aimed at improving diet quality in young people might be more successful when family members are also encouraged to engage in healthy diet quality and support adolescents' diet quality. Another important point is the fact that adolescents' perceptions of their peers'/families' engagement/encouragement may also play a major role in their dietary quality and should be addressed in intervention programmes focusing on adolescents.

Acknowledgements

Acknowledgements: The authors thank the participants for taking part in the study. *Financial support:* The HELENA Study is made possible by the financial support of the European Community Sixth RTD Framework Programme (Contract FOOD-CT-20056007034) and the Spanish Ministry of Science and Innovation (RYC-2010-05957 and RYC-2011-09011). The funders had no role in the design, analysis or writing of this article. The content of this paper reflects only the authors' views, and the European Community is not liable for any use that may be made of the information contained therein. *Conflict of interest:* The authors declared no conflict of interest. *Authorship:* Each author of this article significantly contributed to the study. F.G., S.D.H., A.K., K.W., M.S., M.K. and J.R.R. designed the research; J.V., L.B., J.R.R., M.S., A.K., M.K. and K.W. conducted the research; E.D. analysed data; A.D. analysed data and performed statistical analyses; J.V., L.B. and F.G. wrote the paper; F.G. had primary responsibility for the final content. All authors read and approved the final manuscript. *Ethics of human subject participation:* This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects were approved by the local ethics committee for each country. Written informed consent was obtained from all subjects and their parents.

References

1. Von Post-Skagegård M, Samuelson G, Karlström B *et al.* (2002) Changes in food habits in healthy Swedish adolescents during the transition from adolescence to adulthood. *Eur J Clin Nutr* **56**, 532–538.
2. Lake AA, Mathers JC, Rugg-Gunn AJ *et al.* (2006) Longitudinal change in food habits between adolescence (11–12 years) and adulthood (32–33 years): the ASH30 Study. *J Public Health* **28**, 10–16.
3. Niemeier HM, Raynor HA, Lloyd-Richardson EE *et al.* (2006) Fast food consumption and breakfast skipping: predictors of weight gain from adolescence to adulthood in a nationally representative sample. *J Adolesc Health* **39**, 842–849.
4. Furst T, Connors M, Bisogni CA *et al.* (1996) Food choice: a conceptual model of the process. *Appetite* **26**, 247–265.
5. Fitzgerald A, Heary C, Nixon E *et al.* (2010) Factors influencing the food choices of Irish children and adolescents: a qualitative investigation. *Health Promot Int* **25**, 289–298.
6. Kant AK & Graubard BI (2007) Secular trends in the association of socio-economic position with self-reported dietary attributes and biomarkers in the US population: National Health and Nutrition Examination Survey (NHANES) 1971–1975 to NHANES 1999–2002. *Public Health Nutr* **10**, 158–167.
7. Lallukka T, Laaksonen M, Rahkonen O *et al.* (2007) Multiple socio-economic circumstances and healthy food habits. *Eur J Clin Nutr* **61**, 701–710.
8. Hulshof KF, Brussaard JH, Kruizinga AG *et al.* (2003) Socioeconomic status, dietary intake and 10 y trends: the Dutch National Food Consumption Survey. *Eur J Clin Nutr* **57**, 128–137.
9. Slimani N, Fahey M, Welch AA *et al.* (2002) Diversity of dietary patterns observed in the European Prospective Investigation into Cancer and Nutrition (EPIC) project. *Public Health Nutr* **5**, 1311–1328.
10. Agudo A, Slimani N, Ocké MC *et al.* (2002) Consumption of vegetables, fruit and other plant foods in the European Prospective Investigation into Cancer and Nutrition (EPIC) cohorts from 10 European countries. *Public Health Nutr* **5**, 1179–1196.
11. Pedersen S, Grønhøj A & Thøgersen J (2015) Following family or friends. Social norms in adolescent healthy eating. *Appetite* **86**, 54–60.
12. Wouters EJ, Larsen JK, Kremers S *et al.* (2010) Peer influence on snacking behavior in adolescence. *Appetite* **55**, 11–17.
13. Fitzgerald A, Heary C, Kelly C *et al.* (2013) Self-efficacy for healthy eating and peer support for unhealthy eating are associated with adolescents' food intake patterns. *Appetite* **63**, 48–58.
14. Bruening M, Eisenberg M, MacLehose R *et al.* (2012) Relationship between adolescents' and their friends' eating behaviors: breakfast, fruit, vegetable, whole-grain, and dairy intake. *J Acad Nutr Diet* **112**, 1608–1613.
15. Croll JK, Neumark-Sztainer D & Story M (2001) Healthy eating: what does it mean to adolescents? *J Nutr Educ* **33**, 193–198.
16. Kristjansdottir AG, De Bourdeaudhuij I, Klepp KI *et al.* (2009) Children's and parents' perceptions of the determinants of children's fruit and vegetable intake in a low-intake population. *Public Health Nutr* **12**, 1224–1233.
17. Rasmussen M, Krølner R, Klepp KI *et al.* (2006) Determinants of fruit and vegetable consumption among children and adolescents: a review of the literature. Part I: Quantitative studies. *Int J Behav Nutr Phys Act* **11**, 22.
18. Vereecken CA, De Henauw S, Maes L *et al.* (2009) Reliability and validity of a healthy diet determinants questionnaire for adolescents. *Public Health Nutr* **12**, 1830–1838.
19. Vereecken CA, Covents M, Sichert-Hellert W *et al.* (2008) Development and evaluation of a self-administered computerized 24-h dietary recall method for adolescents in Europe. *Int J Obes (Lond)* **32**, Suppl. 5, S26–S34.
20. Dehne LI, Klemm C, Henseler G *et al.* (1999) The German food code and nutrient data base (BLS II.2). *Eur J Epidemiol* **15**, 355–359.
21. Béghin L, Dauchet L, Vriendt T *et al.* (2013) Influence of parental socio-economic status on diet quality of European adolescents: results from the HELENA study. *Br J Nutr* **13**, 1–10.
22. Vlaams Instituut voor Gezondheidspromotie (2012) *De actieve voedingsdriehoek: een praktische voedings – en beweeggids (The Active Food Pyramid: A Practical Guide to Diet and Physical Activity)*. Brussels: VIG.
23. Huybrechts I, Vereecken C, De Bacquer D *et al.* (2010) Reproducibility and validity of a diet quality index for children assessed using a FFQ. *Br J Nutr* **104**, 135–144.

24. Vyncke K, Cruz-Fernandez E, Fajo-Pascual M *et al.* (2013) Validation of the Diet Quality Index for Adolescents by comparison with biomarkers, nutrient and food intakes: the HELENA study. *Br J Nutr* **109**, 2067–2078.
25. Cole TJ, Bellizzi MC, Flegal KM *et al.* (2000) Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ* **320**, 1240–1243.
26. Nagy E, Vicente-Rodriguez G, Manios Y *et al.*; HELENA Study Group (2008) Harmonization process and reliability assessment of anthropometric measurements in a multi-center study in adolescents. *Int J Obes (Lond)* **32**, Suppl. 5, S58–S65.
27. Wuenstel JW, Wądołowska L, Słowinska MA *et al.* (2016) Intake of dietary fibre and its sources related to adolescents' age and gender, but not to their weight. *Cent Eur J Public Health* **24**, 211–216.
28. Winkvist A, Hultén B, Kim JL *et al.* (2016) Dietary intake, leisure time activities and obesity among adolescents in Western Sweden: a cross-sectional study. *Nutr J* **21**, 41.
29. Benjamini Y & Hochberg Y (1995) Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc B* **57**, 289–300.
30. Van Buuren S & Groothuis-Oudshoorn K (2011) MICE: multivariate imputation by chained equations in R. *J Stat Softw* **45**, 1–66.
31. Molaison EF, Connell CL, Stuff JE *et al.* (2005) Influences on fruit and vegetable consumption by low-income black American adolescents. *J Nutr Educ Behav* **37**, 246–251.
32. Neumark-Sztainer D, Story M, Perry C *et al.* (1999) Factors influencing food choices of adolescents: findings from focus-group discussions with adolescents. *J Am Diet Assoc* **99**, 929–937.
33. Wind M, Bobelijn K, De Bourdeaudhuij I *et al.* (2005) A qualitative exploration of determinants of fruit and vegetable intake among 10- and 11-year-old schoolchildren in the low countries. *Ann Nutr Metab* **49**, 228–235.
34. Faith MS, Scanlon KS, Birch LL *et al.* (2004) Parent-child feeding strategies and their relationships to child eating and weight status. *Obes Res* **12**, 1711–1722.
35. Sichert-Hellert W, Beghin L, De Henauw S *et al.* (2011) Nutritional knowledge in European adolescents: results from the HELENA (Healthy Lifestyle in Europe by Nutrition in Adolescence) study. *Public Health Nutr* **14**, 2083–2091.
36. Kim C, Hanjoo L & Marc A (2009) Adolescents' perceptions of family communication patterns and some aspects of their consumer socialization. *Psychol Mark* **26**, 888–907.
37. Iliescu C, Beghin L, Maes L *et al.* (2008) Socioeconomic questionnaire and clinical assessment in the HELENA Cross-Sectional Study: methodology. *Int J Obes (Lond)* **32**, Suppl. 5, S19–S25.
38. Sung-Chan P, Sung YW, Zhao X *et al.* (2013) Family-based models for childhood-obesity intervention: a systematic review of randomized controlled trials. *Obes Rev* **14**, 265–278.
39. Barr-Anderson DJ, Adams-Wynn AW, DiSantis KI *et al.* (2013) Family-focused physical activity, diet and obesity interventions in African-American girls: a systematic review. *Obes Rev* **14**, 29–51.
40. Hingle MD, O'Connor TM, Dave JM *et al.* (2010) Parental involvement in interventions to improve child dietary intake: a systematic review. *Prev Med* **51**, 103–111.
41. Kitzman-Ulrich H, Wilson DK, St George SM *et al.* (2010) The integration of a family systems approach for understanding youth obesity, physical activity, and dietary programs. *Clin Child Fam Psychol Rev* **13**, 231–253.
42. Leslie LK, Mehus CJ, Hawkins JD *et al.* (2016) Primary health care: potential home for family-focused preventive interventions. *Am J Prev Med* **51**, 4 Suppl. 2, S106–S118.
43. Prado G, Pantin H & Estrada Y (2015) Integrating evidence-based interventions for adolescents into primary care. *Am J Prev Med* **48**, 488–490.
44. Turner KM, Shield JP & Salisbury C (2009) Practitioners' views on managing childhood obesity in primary care: a qualitative study. *Br J Gen Pract* **59**, 856–862.
45. Barlow SE; Expert Committee (2007) Expert committee recommendations regarding the prevention, assessment, and treatment of child and adolescent overweight and obesity: summary report. *Pediatrics* **120**, Suppl. 4, S164–S192.
46. August GP, Caprio S, Fennoy I *et al.* (2008) Prevention and treatment of pediatric obesity: an endocrine society clinical practice guideline based on expert opinion. *J Clin Endocrinol Metab* **93**, 4576–4599.
47. Feinberg ME, Solmeyer AR & McHale SM (2012) The third rail of family systems: sibling relationships, mental and behavioral health, and preventive intervention in childhood and adolescence. *Clin Child Fam Psychol Rev* **15**, 43–57.
48. Quiles Marcos Y, Quiles Sebastián MJ, Pamies Aubalat L *et al.* (2013) Peer and family influence in eating disorders: a meta-analysis. *Eur Psychiatry* **28**, 199–206.
49. Larson R & Richards MH (1991) Daily companionship in late childhood and early adolescence: changing developmental contexts. *Child Dev* **62**, 284–300.
50. Levine MP, Smolak L, Moodey AF *et al.* (1994) Normative developmental challenges and dieting and eating disturbances in middle school girls. *Int J Eat Disord* **15**, 11–20.
51. Hill AJ & Franklin JA (1998) Mothers, daughters and dieting: investigating the transmission of weight control. *Br J Clin Psychol* **37**, 3–13.
52. Sawka KJ, McCormack GR, Nettel-Aguirre A *et al.* (2015) Associations between aspects of friendship networks and dietary behavior in youth: findings from a systematized review. *Eat Behav* **18**, 7–15.
53. Neumark-Sztainer D, Wall M, Guo J *et al.* (2006) Obesity, disordered eating, and eating disorders in a longitudinal study of adolescents: how do dieters fare 5 years later? *J Am Diet Assoc* **106**, 559–568.
54. Patton GC, Selzer R, Coffey C *et al.* (1999) Onset of adolescent eating disorders: population based cohort study over 3 years. *BMJ* **318**, 765–768.
55. Field AE, Austin SB, Taylor CB *et al.* (2003) Relation between dieting and weight change among preadolescents and adolescents. *Pediatrics* **112**, 900–906.
56. Stice E, Presnell K, Shaw H *et al.* (2005) Psychological and behavioral risk factors for obesity onset in adolescent girls: a prospective study. *J Consult Clin Psychol* **73**, 195–202.
57. Stice E, Presnell K & Spangler D (2002) Risk factors for binge eating onset in adolescent girls: a 2-year prospective investigation. *Health Psychol* **21**, 131–138.
58. Julián-Almárcegui C, Bel-Serrat S, Kersting M *et al.* (2015) Comparison of different approaches to calculate nutrient intakes based upon 24-h recall data derived from a multi-center study in European adolescents. *Eur J Nutr* **55**, 537–545.