

EPV0317

The emergence of new obsessions and compulsions after COVID-19: a case report

M. Abdelkefi*, R. Feki, I. Gassara, N. Smaoui, S. Omri, N. Charfi, L. Zouari, J. Ben thabet, M. Maalej bouali and M. Maalej

Psychiatry C department, Hedi Chaker university hospital, sfax, Tunisia

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1049

Introduction: The coronavirus pandemic has affected mental health since its outbreak in 2019 and several studies have revealed that obsessive-compulsive disorder (OCD) patients were adversely affected.

Objectives: The aim of our present report is to study the impact of the coronavirus infection on OCD.

Methods: We illustrate a case of new emerging obsessions and compulsions after a COVID-19 infection in a patient consulting at the Psychiatry C department of the Hedi Chaker University Hospital.

Results: Mr. HB first presented in our outpatient unit of the Psychiatry C department at the age of 32. His medical history was unremarkable except for a COVID-19 infection in June 2022 that did not cause any organic complications. His family history was negative without neurological and psychiatric diseases. Further history revealed that OCD symptoms already started before his infection with COVID-19. At that time, he suffered from compulsions with the urge to constantly wash his hands and check rituals, but his symptoms were not severe enough to make him seek a psychiatric consult or treatment. However, since July 2022, soon after his infection with COVID-19, he suffered from new obsessions and compulsions, he would spend hours calculating all the numbers he sees and counting the number of letters in the words he came across. Non-surprisingly, OCD symptoms caused relevant problems in social life, and at work, he had difficulties concentrating and working. Due to his symptoms, he had neglected hobbies, avoided social contact, spent less time with his family, and even had suicidal thoughts. Therefore, pharmacotherapy with clomipramine was initiated with a maximal dose of 150 mg/d. Because of adverse events and lack of efficacy, he stopped medication and decided to consult our department to seek a different treatment. Therefore, sertraline was started at the dose of 50 mg/d along with psychotherapy.

Conclusions: People with OCD are likely to be more susceptible to the mental health impact of COVID-19. Responses to the pandemic are not only associated with an increase in all Obsessive-Compulsive (OC) symptoms and their severity but also the emergence of new types of obsessions and compulsions. Consequently, therapists should consider the effects of the pandemic on all OC symptom dimensions and adjust their treatment plans accordingly.

Disclosure of Interest: None Declared

EPV0318

Psychopharmacological factor in the course of COVID-19 among psychiatric inpatients

M. Sorokin¹, A. Shabelnik¹, M. Bocharova^{1,2*}, N. Lutova¹, O. Limankin^{3,4}, L. Azarova³ and N. Neznanov^{1,5}

¹V.M.Bekhterev National Medical Research Centre for Psychiatry and Neurology, St. Petersburg, Russian Federation; ²King's College London, London, United Kingdom; ³St. Petersburg Psychiatric Hospital No. 1 named after P.P. Kashchenko; ⁴North-Western State Medical University named after I.I. Mechnikov and ⁵FSBEI HE I.P. Pavlov SPbSMU MOH Russia, St. Petersburg, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1050

Introduction: It is known that many psychopharmacological drugs have anti-inflammatory, as well as antibacterial and antiviral effects.

Objectives: To investigate the association between the severity and duration criteria of COVID-19 with psychopharmacotherapy in double-diagnosed patients.

Methods: A total of 169 case histories from a specialized infectious psychiatric department (May 2020 to January 2021) were evaluated. Progression indicators of severe and mild COVID-19, along with the duration of persistent SARS-CoV-2 viral shedding, were assessed in correlation with the administration of antidepressants, antipsychotics, and acid sphingomyelinase inhibitors (FIASMA-active drugs).

Results: The use of any psychotropic agents was associated with a 0.9% increase in the risk of severe course of COVID-19 for each unit increase in the systemic inflammation index PLR, specifically in patients with intellectual disability (ICD-10 codes F70-79), when compared to patients with schizophrenia (ICD-10 codes F20-29): $R^2McF=0.138$; $AIC=181$; $\chi^2=25.8$; $df=9$; $p=0.002$. High PLR values and the use of FIASMA-active drugs were associated with prolonged COVID-19 duration, while antidepressant therapy and elevated C-reactive protein levels were associated with a reduced predicted duration of viral shedding in 13.8% of variance: $R^2=0.0864$; $AIC=1299$; $F=5.2(3)$, $p=0.002$. Including the nosology of psychiatric disorders in the regression model increased the proportion of explained variance to 22.8%.

Conclusions: Thymoanaleptic therapy for individuals with psychiatric disorders may act as a protective factor against COVID-19. There is no evidence suggesting adverse effects of antipsychotics on the severity and duration of COVID-19. Further research is necessary to investigate the effects of FIASMA-active psychopharmacological agents within nosologically homogeneous groups.

Disclosure of Interest: None Declared

EPV0319

The factor structure the PHQ-9 in Russian patients recovered from COVID-19

M. Zinchuk^{1*}, G. Kustov¹, S. Popova¹, A. Razmakhnin¹, D. Zhuravlev¹, R. Akzhigitov¹ and A. Guekht^{1,2}

¹Moscow Research and Clinical Centre for Neuropsychiatry and ²Pirogov Russian National Research Medical University, Moscow, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1051

Introduction: The nine-item Patient Health Questionnaire-9 (PHQ-9) is the first choice for screening for depression in primary care and other medical settings. The PHQ-9 has been shown to be a reliable and valid measure of depression symptoms, but there is disagreement among researchers about the factor structure of this questionnaire. Recent systematic reviews have found four different factor models of the PHQ-9, with one- and two-factor models being the most common. This discrepancy may be due to linguistic, cultural and clinical differences between the populations studied. The factor structure of the Russian version of the PHQ-9 during the COVID-19 pandemic has not been examined in any study to date.

Objectives: The aim of our study was to determine the factorial structure and internal consistency of the Russian version of the PHQ-9 in COVID-19 survivors.

Methods: Fourteen thousand 725 (female - 11479 (78.0%), age - 18-79 years (M - 47.09, SD - 12.70) participants completed an online survey including the PHQ-9 and an ad hoc questionnaire focusing on sociodemographic and COVID-related characteristics. McDonald's omega coefficient was estimated to determine the internal consistency of the questionnaire. Exploratory structural equation modelling (ESEM) with weighted least squares mean and variance adjusted estimator and geomin rotation was performed in Mplus 7.

Results: ESEM provided evidence for a three-factor structure of the PHQ-9, representing affective (items 2, 6, 9), anergic (items 1, 3) and somatic (items 3, 5, 7, 8) dimensions of depression. These factors fit the data well (CFI - 0.998; TLI - 0.994; RMSEA (95% CI) - 0.028 (0.024 - 0.032)), better than a single factor (CFI - 0.955; TLI - 0.940; RMSEA (95% CI) - 0.089 (0.087 - 0.092)) and two-factor (CFI - 0.985; TLI - 0.971; RMSEA (95% CI) - 0.062 (0.059 - 0.065)). The McDonald's omega was 0.82.

Conclusions: Our study revealed a three-factor structure of the Russian version of the PHQ-9 in COVID-19 survivors. COVID-19. A high internal consistency of the Russian version of the instrument was confirmed.

Disclosure of Interest: None Declared

EPV0320

Investigation of the factor structure of GAD-7 in Moscow residents exposed to SARS-CoV2

M. Zinchuk^{1*}, G. Kustov¹, V. Nadezhda¹, A. Razmakhnin¹, D. Zhuravlev¹, R. Akzhigitov¹ and A. Guekht^{1,2}

¹Moscow Research and Clinical Centre for Neuropsychiatry and ²Pirogov Russian National Research Medical University, Moscow, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1052

Introduction: Rates of anxiety in the general population increased significantly during the COVID-19 pandemic. Several studies have shown that people exposed to SARS-CoV2 are at increased risk for both exacerbation and de novo development of anxiety disorders. Therefore, screening for anxiety disorders in this at-risk population is essential. In pre-pandemic studies, the 7-item Generalized Anxiety Disorder Questionnaire (GAD-7) was one of the most commonly used self-report instruments. Its validity has been demonstrated in several studies. However, there is no agreement among researchers about its underlying internal structure. Both

one-factor and two-factor solutions have been reported. This discrepancy may be due to linguistic, cultural, and clinical differences between the populations studied. To our knowledge, no studies have been conducted to investigate the factor structure of the GAD-7 in the Russian-speaking community sample and the psychometric properties of this questionnaire in SARS-CoV2 exposed individuals.

Objectives: The aim of the study was to determine the factorial structure and internal consistency of the Russian version of the GAD-7 in a large sample of Moscow residents exposed to SARS-CoV2.

Methods: Fourteen thousand 725 (male - 11479 (78.0%), age - 18-79 years (M - 47.09, SD - 12.70) Moscow residents exposed to SARS-CoV2 completed an online survey including the GAD-7 and an ad hoc questionnaire focusing on socio-demographic characteristics. McDonald's Omega was used to assess internal consistency. Exploratory structural equation modelling (ESEM) with weighted least squares means and variance adjusted estimator and geomin rotation was used to assess the factor structure of the Russian version of the GAD-7.

Results: The McDonald's Omega of the Russian version of the GAD-7 was 0.85, indicating a good internal consistency of the questionnaire. ESEM provided evidence for a one-factor solution that fits the data well (CFI - 0.996; TLI - 0.995; RMSEA (95% CI) - 0.041 (0.037 - 0.045)).

Conclusions: In Russian people exposed to SARS-CoV2, the GAD-7 showed good internal consistency. Our results are consistent with those of previous studies that reported a single-factor solution for the questionnaire.

Disclosure of Interest: None Declared

EPV0321

Coping strategies facing Covid-19, perceived social support, and trait anxiety among Tunisian caregivers

N. Smaoui^{1*}, B. Jallouli^{1,2}, I. Gassara¹, R. Feki¹, S. Omri^{1,3}, M. Bou Ali Maalej¹, N. Charfi¹, J. Ben Thabet¹, L. Zouari¹ and M. Maalej¹

¹Psychiatry C department, Hedi Chaker university hospital; ²Faculty of Medicine, Sfax and ³University of Sfax, SFAX, Tunisia

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.1053

Introduction: Caregivers in the Sfax region, Tunisia, having been at the forefront in the face of the Covid-19 pandemic, were therefore faced with intense stress. It seemed useful and interesting to us to study their adaptation strategies during this period of pandemic.

Objectives: The aims of our study were to identify the coping strategies used by Tunisian Healthcare workers (HCW) during the Covid-19 pandemic and to study the links of the different coping strategies with perceived social support and trait anxiety.

Methods: A cross-sectional, descriptive, and analytical study conducted among 254 Tunisian HCW working at the Habib Bourguiba and Hedi Chaker university hospitals in Sfax, during period from January 2021 to April 2021. the questionnaire used included an information sheet and three scales; "Social support questionnaire 6" (SSQ-6), "State Trait Inventory Anxiety Form Y2" (STAI-Y2), and "Ways of Coping Checklist" (WCC).