

ERRATUM

Adherence to a healthy Nordic food index is associated with a lower incidence of colorectal cancer in women: the Diet, Cancer and Health cohort study - ERRATUM

Cecilie Kyrø, Guri Skeie, Steffen Loft, Kim Overvad, Jane Christensen, Anne Tjønneland and Anja Olsen

(First published online 11 November 2013)

doi:10.1017/S0007114512002085, Published by Cambridge University Press, 3 July 2012.

We regret to announce that there Table 2 was published incorrectly in the article⁽¹⁾. All the data was correct, but the energy intake column was not rendered correctly. The publishers apologise for this error.

Reference

1. Kyrø C, Skeie G, Loft S, et al. (2013) Adherence to a healthy Nordic food index is associated with a lower incidence of colorectal cancer in women: the Diet, Cancer and Health cohort study. Br J Nutr 109, 920-927. Published by Cambridge University Press, 3 July 2012, doi:10.1017/S0007114512002085.



Table 2. Baseline characteristics of all participants in the Diet, Cancer and Health cohort and by healthy Nordic food index scores of 0-1 points (poorest adherence), 2-3 points and 4-6 points (best adherence)

(Medians, number of participants, percentiles and percentages)

Characteristic	Food index score															
	Men								Women							
	All		0-1		2-3		4-6		All		0-1		2–3		4-6	
	Median	P5-P95	Median	P5-P95	Median	P5-P95	Median	P5-P95	Median	P5-P95	Median	P5-P95	Median	P5-P95	Median	P5-P95
n	26 664		6608		11 966		8090		29216		5464		13 728		10 024	
%	100		25		45		30		100		19		47		34	
Cases																
n	567		142		273		152		458		108		203		147	
%	100		25		48		27		100		24		44		32	
Age (years)	56	50-64	55	50-64	56	50-64	56	50-64	56	50-64	55	50-64	56	50-64	56	50-64
Schooling (%)																
Short (≤7 years)		35		41		35		29		31		37		32		28
Medium (8-10 years)		42		42		42		40		50		49		51		50
Long (≥11 years)		24		17		23		31		19		15		17		22
Smoking status (%)																
Never		26		22		25		30		44		37		43		48
Past	35		28		35		39		24		19		23		27	
Current		40		50		40		31		33		44		35		24
BMI (kg/m ²)	26	21-33	26	22-34	26	22-33	26	21-32	25	20-34	25	20-34	25	20-34	25	20-34
Waist	95	81-114	96	82-116	95	82-113	94	81-112	80	67-103	81	67-105	80	67-103	80	67-103
circumference (cm)																
Alcohol intake (g/d)	19	2-80	21	1-91	19	2-78	19	2-67	9	0-42	8	0-48	10	0-43	9	1-39
Participate in sports (%)		49		38		48		58		58		45		57		68
Intake of red and processed meat (g/d)	139	67–254	131	67–237	140	67–252	144	65-272	83	35–157	78	35-145	84	36-154	87	34-166
Energy intake (kJ/d), excluding alcohol	9908	6414-14942	8492	5518-12542	9781	6666-14316	11 232	7898-16332	8092	5092-12318	6677	4159-10206	7800	5216-11 568	9155	6241-13411

P5, 5th percentile; P95, 95th percentile.