

## A community outbreak of *Campylobacter jejuni* infection from a chlorinated public water supply

G. RICHARDSON<sup>1</sup>, D. RH. THOMAS<sup>1</sup>, R. M. M. SMITH<sup>1</sup>, L. NEHAUL<sup>2</sup>,  
C. D. RIBEIRO<sup>3</sup>, A. G. BROWN<sup>4</sup> AND R. L. SALMON<sup>1\*</sup>

<sup>1</sup> National Public Health Service, Communicable Disease Surveillance Centre, Cardiff, UK

<sup>2</sup> National Public Health Service, South East Wales Health Protection Unit, Mamhilad House, Pontypool, UK

<sup>3</sup> National Public Health Service, Microbiology Cardiff, University of Wales, Cardiff, UK

<sup>4</sup> Caerphilly County Borough Council, Environmental Health, Council Offices, Blackwood, UK

(Accepted 16 December 2006; first published online 9 February 2007)

### SUMMARY

An outbreak of *Campylobacter jejuni* infection occurred in a South Wales Valleys housing estate. Illness in estate residents was associated with tap water consumption [population attributable risk (PAR) 50%, relative risk (RR) 2.53, 95% confidence interval (CI) 1.9–3.37] and residence in the upper estate (PAR 49%, RR 2.44, 95% CI 1.83–3.24). Amongst upper estate residents, rates of diarrhoeal illness increased with rates of water consumption (OR 18, 95% CI 3.5–92.4 for heaviest consumers,  $\chi^2$  trend  $P < 0.0001$ ). The upper estate received mains water via a covered holding reservoir. A crack in the wall of the holding reservoir was identified. Contamination with surface water from nearby pasture land was the likely cause of this outbreak. Service reservoirs are common in rural communities and need regular maintenance and inspection. The role of water in sporadic cases of campylobacter enteritis may be underestimated.

### INTRODUCTION

Campylobacters are important emerging pathogens worldwide causing a considerable health and economic burden [1]. Infections are usually sporadic and although outbreaks related to food, raw milk and untreated water have been described, they are unusual. Outbreaks in treated public water supplies are rare and only three have been previously documented, two in the United States [2, 3] and one in Sweden [4].

In September 2000, 15 cases of gastroenteritis due to *Campylobacter jejuni* were reported amongst residents

of an economically deprived housing estate located in the South Wales Valleys. The estate, comprising mainly local government-owned rented accommodation, is located on a steep hill and is surrounded by agricultural pasture land. Cases became ill between 17 and 24 September and only low numbers of *C. jejuni* (<5) had been notified for the area in the 6 months prior to the outbreak indicating an unusual event. Initial investigation by the local government environmental health department found that patients were clustered in the upper part of this housing estate and reported drinking large quantities of tap water.

Chlorinated mains water is supplied to the upper part of the hillside estate (442 properties) by pumping from the main supply to a small concrete service reservoir, submerged into the hillside. The lower part of the estate (248 properties) is supplied via a different

\* Author for correspondence: Dr R. L. Salmon, National Public Health Service Communicable Disease Surveillance Centre, Abton House, Wedal Rd, Cardiff, CF14 3QX, UK.  
(Email: Roland.Salmon@nphs.wales.nhs.uk)

route. This raised suspicion that the geographical clustering of cases in the upper estate may have been related to water supply.

## METHODS

### Cohort study

We conducted a retrospective cohort study of diarrhoeal illness in estate residents in all 690 households. Information was collected by postal questionnaire on individual's self-reported exposure to household pets and farm animals. An individual dietary history was recorded, including specific questions on meats, meat products, takeaway meals, milk, carbonated drinks, tea/coffee, tap water and bottled water. Residents were asked where they purchased milk. Where home delivery of milk was recorded, the dairy supplying the milkman was identified by follow-up investigation. Residents were assigned a source of mains tap water by whether they were resident in the upper or lower estate. Any history of diarrhoeal illness amongst residents or pets was noted and dates and symptoms recorded. A case was considered to be an individual residing on the estate, with diarrhoeal illness ( $\geq 3$  stools within 24 h) lasting over 1 day in September 2000.

### Nested case-control study

Using data collected from the cohort study we also carried out a nested case-control study of the 15 laboratory-confirmed cases and 93 estate resident controls that had not experienced diarrhoeal illness (simple random sample of 10% of cohort). There were no other exclusion criteria.

### Statistical analysis

Univariate analysis was performed by Mantel-Haenszel  $\chi^2$  test or Fisher's exact test (two-tailed) using Epi-Info, version 6 (CDC, Atlanta, GA, USA). Univariate analysis was repeated and multivariate analysis carried out by logistic regression using STATA software, version 6 (Stata Corp., College Station, TX, USA) with confidence intervals adjusted for household clustering using the Huber-White robust estimate of variance. Initially all exposures were controlled for age and sex. In the second model, all exposures were controlled for age, sex, household clustering and any variables statistically significantly associated with illness at the 5% level from the first model (water supply, water consumption, milkman delivery, cold takeaway

consumption and puppy/kitten in house with or without diarrhoea).

### Microbiological investigations

Serotyping of campylobacter isolates was not routinely carried out by local laboratories. However, as part of the outbreak investigation samples from cases were forwarded by the two local hospital laboratories to the Laboratory of Enteric Pathogens, Central Public Health Laboratory, London (now Centre for Infections, Health Protection Agency) for serotyping.

### Environmental investigations

The service reservoir outlet supplying the upper estate is sampled weekly by the water company on random days. Bacteriological and chlorination records of weekly sampling of the service reservoir were provided by the water company and evaluated by the outbreak control team. Sampling of the service reservoir supplying the upper estate was carried out daily from 30 September by the water company and samples were tested for indicator organisms (coliforms and *E. coli*) and free and total chlorine levels. Additional analyses for faecal streptococci and clostridia were also performed. All methods and standards used were as set out by the European Council Directive 98/83/EC of 3 November 1998 on the quality of water intended for human consumption [5].

The water company conducted a physical external inspection of the reservoir on 2 October and one compartment of the reservoir was taken out of service for internal inspection. The second compartment was taken out of service for inspection on 11 October. Rainfall records for the area, maintained by the national Meteorological Office, were obtained.

## RESULTS

### Cohort study

Questionnaires were received from 528 households (76.5%) with 1220 personal responses. Diarrhoeal illness was reported by 281 (23%) of respondents. Of those, 22 (7.8%) reported blood in stools and 102 (36.3%) contacted their family practitioner. The median length of illness was 4 days (mode 2 days) and mean daily stool frequency on the worst day of illness was 10 (mode 5). The attack rate for those resident in the upper estate was 232/802 (29.0%) compared with




Fig. 1. Epidemic curve for diarrhoeal illness in upper (□) and lower (■) estate residents.

49/413 (11.9%) for those in the lower estate [relative risk (RR) 2.44, 95% confidence interval (CI) 1.83–3.24,  $P < 0.001$ ]. An epidemic curve for diarrhoeal illness in estate residents (Fig. 1) indicated raised levels of diarrhoeal illness in upper estate residents during a 2-week window in September 2000 against a general background of low levels of diarrhoeal illness in the whole estate. Attack rates varied by age, with the highest attack rate in the  $\leq 4$  years age group (40%) and the lowest in those aged  $\geq 65$  years (13.4%).

Upper estate residence and tap water consumption were significantly associated with illness (Table 1) with a significant dose–response (Fig. 2). The odds ratio for illness after consumption of 1–3 glasses of upper estate tap water in 24 h was 2.42 (95% CI 1.59–3.70), with increasing odds ratios for increased levels of consumption to a maximum of  $\geq 11$  glasses (OR 18, 95% CI 3.51–92.40). The population attributable risk (PAR) for upper estate water supply was 49% and for tap water consumption 50%.

Other exposures associated with illness were: consumption of cold milk, fizzy drinks, cold takeaways for re-heating at home, home delivery of milk by a milkman, having a puppy or kitten in the house and having a puppy or kitten with diarrhoea (Table 1).

Of 231 individuals who had milk delivered at home 71 had experienced bottle-tops being pecked by birds (30.7%). This exposure was not significantly associated with illness (RR 0.85, 95% CI 0.57–1.40,  $P = 0.617$ ).

Stratified univariate analysis of milk consumption among persons reporting milkman delivery was

performed. There was no significant association between consuming milk and illness (OR 1.57, 95% CI 0.61–4.04,  $P = 0.351$ ).

Upper water supply (OR 2.97 95% CI 1.83–4.8,  $P < 0.0001$ ) and tap water consumption (OR 3.24 95% CI 2.18–4.8,  $P < 0.0001$ ) remained significant effects after controlling for other exposures (Table 2). However, one confounding factor was attendance at the estate primary school, which, although located in the lower estate received upper water supply. Many children aged between 4 and 11 years from both the upper and lower estate attended this school so that children living in the lower water supply area drank upper supply water during school hours.

Being aged  $< 11$  years showed increased risk of illness independent of residence in the upper or lower estate (RR for lower part of the estate 3.48, 95% CI 2.07–5.85,  $P < 0.001$ ; RR for upper estate 1.70, 95% CI 1.35–2.14,  $P < 0.001$ ). Consumption of cold takeaway meals re-heated at home was also significantly associated with illness but only 13 of the 281 possible cases had been exposed to these (PAR 2.3%, OR 3.45, 95% CI 1.47–8.06,  $P < 0.004$ ). Having home delivery of milk was also associated with illness but, again, would only account for a small proportion of diarrhoeal illness in residents (PAR 6.5%, OR 1.58, 95% CI 1.03–2.44,  $P < 0.037$ ).

### Nested case-control study (Table 3)

All laboratory-confirmed cases resided in the upper estate. Of the 93 controls, 39 (42%) lived in the lower

Table 1. Self-reported diarrhoeal illness in 1220 estate residents: univariate analysis by  $\chi^2$  test

Exposure	Exposed			Unexposed			Relative risk of illness with exposure, 95% CI and PAR (%)			
	Ill	Well	AR (%)	Ill	Well	AR (%)	RR	95% CI	P	PAR (%)
<b>Water and drinks</b>										
Upper water supply	232	802	28.9	49	413	11.9	2.44	1.83–3.24	0.000	49.0
Tap water	228	773	29.5	48	412	11.7	2.53	1.90–3.37	0.000	50.0
Bottled water	27	137	19.7	240	1024	23.4	1.24	1.00–1.54	0.330	
Fizzy drinks	157	621	25.3	113	556	20.3	1.24	1.01–1.54	0.043	11.4
Tea/Coffee	224	1019	22.0	52	189	27.5	0.80	0.62–1.03	0.096	
<b>Milk</b>										
Cold milk	188	741	25.4	87	443	19.6	1.29	1.03–1.62	0.024	15.4
Hot milk	50	155	24.4	220	749	22.7	1.07	0.82–1.40	0.602	
Untreated milk	6	46	13.0	269	1128	23.9	0.55	0.26–1.16	0.090	
Milkman delivery	67	223	30.0	212	965	22.0	1.37	1.08–1.73	0.010	6.5
Bottle-tops pecked	19	74	25.7	195	790	24.7	0.89	0.57–1.40	0.617	
<b>Food</b>										
Poultry	270	1165	23.2	10	46	21.7	1.07	0.61–1.86	0.820	
Red meat	255	1129	22.6	24	74	32.4	0.70	0.49–0.98	0.052	28.5
Pork	248	1092	22.7	31	114	27.2	0.84	0.61–1.15	0.280	
Cooked chicken	41	143	22.3	229	761	23.1	0.96	0.71–1.29	0.802	
Pies	110	461	23.9	166	728	22.8	1.04	0.84–1.29	0.673	
Hot takeaway	48	204	23.5	231	994	23.2	1.01	0.77–1.33	0.930	
Cold takeaway	13	29	44.8	265	1167	22.7	1.97	1.30–3.00	0.005	2.3
Vegetarian	2	17	11.8	273	1188	23.0	0.51	0.14–1.89	0.274	
<b>Animals</b>										
Farm animals	9	27	33.3	269	1179	22.8	1.46	0.85–2.52	0.20	
Puppy/kitten in house	38	94	40.4	241	1111	21.7	1.86	1.42–2.44	0.000	6.3
Puppy/kitten – diarrhoea	12	20	60.0	267	1185	22.5	2.66	1.83–3.87	0.0001	2.7

PAR, Population attributable risk; RR, relative risk; AR, attack rate; OR, odds ratio; CI, confidence interval.

estate, and 54 (58%) in the upper. The odds ratio for illness after drinking tap water was 8.35 with 95% CI 1.05–66.4 ( $P=0.045$ ). When the amount of tap water consumed was considered as an ordinal variable with consumption of 1–3 glasses as baseline, consuming 4–6 glasses of water daily gave an odds ratio for illness of 13.6 (95% CI 1.58–116.87,  $P=0.017$ ) and 7–10 glasses gave an odds ratio of 85 (95% CI 6.46–1118.9,  $P=0.001$ ). None of the 15 laboratory-confirmed cases had been exposed to cold takeaways, puppies, kittens, or farm animals. No other foods or drinks were associated with illness. As in the cohort study there was an association between home delivery of milk and illness (OR 12, 95% CI 3.5–40.8,  $P<0.001$ ). However, consumption of milk was not significantly associated with illness (OR 1.29, 95% CI 0.37–4.38,  $P=0.687$ ). No laboratory-confirmed case that received home-delivered milk had a history of bottle-tops being pecked by birds.

### Microbiological investigations

The Laboratory of Enteric Pathogens cultured two serotypes of *Campylobacter jejuni* (HS23 phage type 1 and HS8) from faecal specimens of cases identified following 4 October.

### Environmental investigations

Routine weekly sampling of the dual-chambered upper supply service reservoir from a common outlet by the water company had been satisfactory on 15 September. On 18 September low levels of contamination (10 coliforms and 8 *E. coli*/100 ml) were detected following heavy rainfall. Free chlorine level had been <0.1 mg/l and total chlorine 0.1 mg/l, below the target level of 0.2 mg/l. The water company had manually dosed the reservoir with sodium hypochlorite and flushed the associated mains before re-sampling the reservoir and distribution system. Sampling on 19–20 September did not reveal any

Table 2. *Self-reported diarrhoeal illness in 1220 estate residents: multivariate analysis by logistic regression*

Exposure	Unadjusted			Adjusted for age, sex (robust cluster)			Adjusted for age, sex, water supply, water consumption, milkman delivery, cold takeaway consumption, puppy/kitten in house, and puppy/kitten diarrhoea (robust cluster)		
	OR	95% CI	P	OR	95% CI	P	OR	95% CI	P
<b>Water and drinks</b>									
Upper water supply	3.02	2.16–4.23	0.000	2.97	1.87–4.74	0.000	2.97	1.83–4.80	0.000
Tap water	3.17	2.26–4.50	0.000	2.95	2.05–4.26	0.000	3.24	2.18–4.80	0.000
Bottled water	0.80	0.51–1.25	0.331	0.87	0.55–1.39	0.560	1.09	0.65–1.80	0.747
Fizzy drinks	1.33	1.01–1.75	0.044	1.20	0.88–1.64	0.241	1.10	0.79–1.52	0.582
Tea/Coffee	0.74	0.52–1.06	0.097	1.33	0.84–2.09	0.221	1.28	0.80–2.05	0.307
<b>Milk</b>									
Cold milk	1.39	1.04–1.85	0.024	1.18	0.85–1.65	0.318	1.00	0.71–1.41	0.989
Hot milk	1.10	0.77–1.56	0.602	1.11	0.75–1.64	0.586	1.22	0.75–1.69	0.578
Untreated milk	0.48	0.20–1.14	0.097	0.48	0.19–1.24	0.131	0.45	0.17–1.25	0.128
Milkman delivery	1.53	1.10–2.11	0.011	1.54	0.99–2.41	0.055	1.58	1.03–2.44	0.037
Bottle-tops pecked	0.85	0.46–1.60	0.617	0.60	0.23–1.58	0.301	0.43	0.16–1.21	0.112
<b>Food</b>									
Poultry	1.09	0.53–2.22	0.821	1.49	0.75–2.94	0.254	1.38	0.63–3.05	0.421
Red meat	0.61	0.37–1.01	0.054	0.85	0.46–1.56	0.605	0.64	0.34–1.23	0.182
Pork	0.79	0.51–1.22	0.281	0.92	0.51–1.67	0.786	0.78	0.43–1.40	0.403
Hot takeaway	1.02	0.71–1.45	0.929	1.09	0.72–1.65	0.667	1.07	0.69–1.66	0.753
Cold takeaway	2.77	1.31–5.82	0.007	3.33	1.61–6.90	0.001	3.45	1.47–8.06	0.004
Vegetarian	0.45	0.10–1.97	0.287	0.41	0.11–1.58	0.197	0.54	0.10–2.84	0.469
<b>Animals</b>									
Farm animals	1.69	0.75–3.81	0.204	1.54	0.43–5.53	0.512	1.51	0.44–5.15	0.513
Puppy/kitten in house	2.45	1.58–3.79	0.000	2.19	1.15–4.16	0.017	1.89	0.93–3.85	0.080
Puppy/kitten – diarrhoea	5.16	2.09–12.75	0.000	4.10	1.02–16.44	0.046	1.64	0.35–7.80	0.533

OR, Odds ratio; CI, confidence interval.

further indicator organisms from the reservoir but free chlorine levels were still low at <0.1 mg/l. A sample from a household tap supplied by the upper estate service reservoir contained 1 faecal coliform/100 ml leading to further mains flushing. The company introduced daily manual sodium hypochlorite dosing of the reservoir on 21 September and reverted to routine weekly sampling. A reservoir sample of 28 September showed no indicator organisms with free and total chlorine 0.1 mg/l. From 30 September the reservoir was sampled daily. The water company reported no documented bacteriological failures prior to 18 September. However, mean free chlorine in the reservoir since May 2000 was <0.1 mg/l.

Figure 3 shows the temporal association between chlorine levels in the upper supply service reservoir and numbers of cases of diarrhoeal illness reported by estate residents. Chlorine levels were rarely above 0.1 mg/l for weeks before the indicator organisms

were detected on 18 September. An increase in the frequency of diarrhoeal illness was reported from 11 September. The water company had not tested bacteria or chlorine levels for the period 20–28 September. Faecal streptococci and clostridia were detected on 4 October in the absence of indicator organisms. Despite boosting, chlorine levels had fallen once more to <0.1 mg/l by 21 November.

Inspection of the upper estate supply service reservoir site externally revealed that the reservoir lay beneath a sheep-grazing field shielded by a protective fence within which was a mature tree. The water company reported that physical external inspection appeared satisfactory. One compartment was internally inspected on 2 October and returned to use following cleaning on 9 October. The full internal inspection of the second reservoir compartment on 11 October revealed a crack in the roof, allowing seepage of surface water through the roof. Rainfall

Table 3. *Laboratory-confirmed campylobacter enteritis: nested case-control study*

Exposure	Proportion exposed		OR	95% CI	P
	Case	Control			
Residence in upper estate	15/15	54/93	—	—	0.000
Tap water	14/15	57/91	8.35	1.1–66.4	0.045
Milkman delivery to household	10/15	13/91	12.0	3.53–40.79	0.000
Cold milk consumption	11/15	62/93	1.29	0.37–4.38	0.687
Prior household contact with an ill person	3/15	17/93	1.12	0.28–0.44	0.874

OR, Odds ratio; CI, confidence interval.




Fig. 2. Association between volume of tap water consumed per day and rate of illness in upper estate residents.

for the month of September was 150% of the monthly average.

## DISCUSSION

This is the first documented waterborne outbreak of campylobacteriosis recorded in a chlorinated public water supply in the United Kingdom. Two waterborne outbreaks of campylobacteriosis in the United States were attributed to inadequate chlorination accompanied by heavy rainfall and a fault in the distribution system [2, 3]. Contaminated water from untreated private water supplies, lakes and streams has been previously identified as a vehicle of both sporadic cases and outbreaks of campylobacteriosis [6–9]. Contamination of surface and ground water occurs by run-off from agricultural land used by livestock and is exacerbated by heavy rainfall or melting snow [10–12].

This outbreak affected 281 residents of a South Wales housing estate. The demarcation of the estate


into two discrete water-supply areas allowed comparisons that confirmed the source. There was a dose-related association between diarrhoeal disease and exposure to upper estate tap water supply.

The cohort investigation provided strong epidemiological evidence of waterborne transmission according to US and UK classifications [13, 14]. The nested case-control study supported the cohort study and strengthened the conclusion that upper estate tap water consumption was responsible for the outbreak.

The epidemic curve for residents of the upper estate suggested a common source outbreak. A mechanism for transmission existed in that the concrete reservoir allowed seepage of surface water contaminated by agricultural waste following heavy rainfall on the surrounding hills. Free chlorine levels in the service reservoir were inadequate to deal with the bacterial load and difficult to sustain. Campylobacters are not routinely tested for by water companies and can be difficult to culture as stressed campylobacters may adopt a viable non-culturable form [15]. *Campylobacter* was not cultured from water samples. However this does not exclude the possibility that it was present.

Inadequate or failed milk pasteurization has been linked to previous outbreaks [16–19]. However processes at two independent dairies would need to have failed in this instance. Consumption of home-delivered milk was not a statistically significant risk factor for illness. Having the bottle-tops of home-delivered milk pecked by birds is known to account for some sporadic campylobacter infections [20, 21] but no confirmed case in this outbreak reported pecked bottle-tops.

Only 15 cases of *C. jejuni* infection were laboratory confirmed. It is likely that many outbreak cases were undiagnosed either because they did not attend their



**Fig. 3.** Temporal association between total chlorine count (—) in upper estate service tank and cases of diarrhoea (---) in estate residents.

doctor (62.8% of those with illness in this cohort) or because their doctor did not send stool specimens for culture. It is well known that there is generalized under-diagnosis, underreporting and under-identification of infectious gastrointestinal illness. A British study found that one case is reported to national surveillance for every 1.5 laboratory notifications, 3.6 cases in primary care and 8.7 cases of campylobacter infection in the community [22].

The findings of this outbreak investigation have implications for the UK water industry, for local authorities and health agencies that may investigate future potential waterborne outbreaks. The authors reported their findings to the Drinking Water Inspectorate (DWI) of England and Wales in June 2001. The DWI reported conclusions and recommendations arising from this outbreak in August 2004 [23]. In its assessment of the incident, the DWI accepted the epidemiological conclusions of the outbreak control team but was very critical of the role of the water company in the outbreak. It concluded that the company did not follow best practice for recognizing and dealing with microbiological incidents and emergencies. In the report the water company was criticized for delayed notification of the incident and for providing reports that contained inadequate and conflicting information. The company's sampling strategy during the early stages of the incident was too limited and was not escalated sufficiently upon detecting faecal coliforms in two spatially and temporally related samples.

It was concluded that the company could have identified which compartment was defective at an earlier stage and could have taken large volume

samples from each compartment by dipping, with enhanced analysis for faecal indicator organisms. It was concluded that the company did not seek appropriate or timely expert microbiological advice and did not act to collect and preserve water samples from the distribution system in anticipation that they may later be required for pathogen analysis. In this regard, outbreak control teams investigating possible waterborne outbreaks could consider obtaining advice and field investigation services, at an early stage, from an organization independent of the water company.

The service reservoir was repaired and permanent booster chlorination facilities commissioned on 29 November 2000. Whilst the DWI concluded that the company had supplied water that was not wholesome, it commended the company for rectifying the service reservoir defects once these had come to light. The DWI reported that the water company responded positively to its critical assessment of the incident and took steps to improve in the areas of concern highlighted, including the value to be gained from sharing the lessons learnt widely with public health professionals.

Campylobacters are the commonest bacterial cause of infectious intestinal disease causing considerable morbidity. They are ubiquitous in the animal kingdom. Minimizing transmission is important wherever animals or their products link with human food and water sources. Service reservoirs are common in rural communities and need a programme of regular maintenance and internal inspection. The role of water in the aetiology of sporadic campylobacteriosis may be underestimated.

## ACKNOWLEDGEMENTS

Robert Davis, Jennifer Harries, Richard Lewis and John Porter assisted with field work and data entry. We are grateful to the GPs and residents of the affected housing estate for their cooperation in the outbreak investigation. Department of Gastrointestinal Infections, Centre for Infections, Health Protection Agency provided support with microbiological investigations. Welsh Water Dwr Cymru provided the outbreak team with the results of their microbiological and environmental investigations.

## DECLARATION OF INTEREST

None.

## REFERENCES

1. **Havelaar AH, et al.** Health burden in the Netherlands due to infection with thermophilic *Campylobacter* species. *Epidemiology and Infection* 2000; **125**: 505–522.
2. **Vogt RL, et al.** *Campylobacter* enteritis associated with contaminated water. *Annals of Internal Medicine* 1982; **96**: 292–296.
3. **Sacks JJ, et al.** Epidemic campylobacteriosis associated with community water supply. *American Journal of Public Health* 1986; **6**: 424–428.
4. **Andersson Y, de Jong B, Studahl A.** Waterborne campylobacter in Sweden: the cost of an outbreak. *Water Science and Technology* 1997; **35**: 11–14.
5. **Council Directive 98/83/EC.** 3 November 1998 on the quality of water intended for human consumption. Official Journal L 330, 05/12/1998 P. 0032-0054 (<http://eurlex.europa.eu/LexUriServ/LexUriServ.do?uri=CELEX:31998L0083:EN:HTML>).
6. **Furtado C, et al.** Outbreaks of waterborne infectious intestinal disease in England and Wales, 1992–1995. *Epidemiology and Infection* 1998; **121**: 109–119.
7. **Levine W, Stephenson T, Craun GF.** Waterborne disease outbreaks, 1986–88. *Mortality and Morbidity Weekly Reports* 1990; **39**: 1–9.
8. **Aho M, et al.** Waterborne outbreak of campylobacter enteritis after outdoors infantry drill in Utti, Finland. *Epidemiology and Infection* 1989; **103**: 133–141.
9. **Palmer SR, et al.** Waterborne outbreak of *Campylobacter* gastroenteritis. *Lancet* 1983; **1**: 287–290.
10. **Melby K, et al.** Clinical and serological manifestations in patients during a waterborne epidemic due to *Campylobacter jejuni*. *Journal of Infection* 1990; **21**: 309–316.
11. **Melby K, et al.** Waterborne *Campylobacteriosis* in northern Norway. *International Journal of Food Microbiology* 1991; **12**: 151–156.
12. **Millson M, et al.** An outbreak of *Campylobacter jejuni* gastroenteritis linked to meltwater contamination of a municipal well. *Canadian Journal of Public Health* 1991; **82**: 27–31.
13. **Barwick RS, et al.** Surveillance for waterborne-disease outbreaks United States 1997–1998. *Morbidity and Mortality Weekly Report* 2000; **40**: 1–35.
14. **Stanwell-Smith R.** Public health and waterborne diseases. *Public Health* 1999; **2**: 53–60.
15. **Gondrosen B.** Survival of thermotolerant campylobacters in water. *Acta Veterinaria Scandinavica* 1986; **27**: 1–10.
16. **Jones PH, et al.** *Campylobacter* enteritis associated with the consumption of free school milk. *Journal of Hygiene* 1981; **87**: 155–162.
17. **Birkhead G, et al.** A multiple-strain outbreak of *Campylobacter* enteritis due to consumption of inadequately pasteurized milk. *Journal of Infectious Diseases* 1988; **157**: 1095–1097.
18. **Fahey T, et al.** An outbreak of *Campylobacter jejuni* enteritis associated with failed milk pasteurisation. *Journal of Infection* 1995; **31**: 137–143.
19. **Porter IA, Reid TM.** A milk-borne outbreak of *Campylobacter* infection. *Journal of Hygiene* 1980; **84**: 415–419.
20. **Riordan T, Humphrey TJ, Fowles A.** A point source outbreak of campylobacter infection related to bird-pecked milk. *Epidemiology and Infection* 1993; **110**: 261–265.
21. **Stuart J, et al.** Outbreak of campylobacter enteritis in a residential school associated with bird pecked bottle tops. *Communicable Disease Report. CDR Review* 1997; **7**: R38–40.
22. **Wheeler JG, et al.** Study of infectious intestinal disease in England: rates in the community, presenting to general practice, and reported to national surveillance. *British Medical Journal* 1999; **318**: 1046–1050.
23. **Drinking Water Inspectorate, 2005.** Drinking Water in Wales 2004 (<http://www.dwi.gov.uk/pubs/annrep04/welshpdf/Englishversion/part1.pdf>).