

Vitamin D status in pregnant Indian women across trimesters and different seasons and its correlation with neonatal serum 25-hydroxyvitamin D levels

R. K. Marwaha^{1*}, N. Tandon², S. Chopra³, N. Agarwal⁴, M. K. Garg⁵, B. Sharma⁴, R. S. Kanwar¹, K. Bhadra¹, S. Singh¹, K. Mani² and S. Puri⁴

¹Department of Endocrinology and Thyroid Research, Institute of Nuclear Medicine and Allied Sciences (INMAS), Brig. SK Mazumdar Marg, Timarpur, Delhi 110 054, India

²Department of Endocrinology and Metabolism, All India Institute of Medical Sciences (AIIMS), New Delhi, India

³Department of Gynecology and Obstetrics, Armed Forces Clinic, New Delhi, India

⁴Department of Food and Nutrition, Institute of Home Economics, University of Delhi, New Delhi, India

⁵Department of Endocrinology, Army Hospital (Research and Referral), Delhi Cantt, New Delhi, India

(Received 12 October 2010 – Revised 1 March 2011 – Accepted 4 March 2011 – First published online 31 May 2011)

Abstract

The present cross-sectional study was conducted to determine the vitamin D status of pregnant Indian women and their breast-fed infants. Subjects were recruited from the Department of Obstetrics, Armed Forces Clinic and Army Hospital (Research and Referral), Delhi. A total of 541 apparently healthy women with uncomplicated, single, intra-uterine gestation reporting in any trimester were consecutively recruited. Of these 541 women, 299 (first trimester, ninety-seven; second trimester, 125; third trimester, seventy-seven) were recruited in summer (April–October) and 242 (first trimester, fifty-nine, second trimester, ninety-three; third trimester, ninety) were recruited in winter (November–March) to study seasonal variations in vitamin D status. Clinical, dietary, biochemical and hormonal evaluations for the Ca–vitamin D–parathormone axis were performed. A subset of 342 mother–infant pairs was re-evaluated 6 weeks postpartum. Mean serum 25-hydroxyvitamin D (25(OH)D) of pregnant women was 23.2 (SD 12.2) nmol/l. Hypovitaminosis D (25(OH)D < 50 nmol/l) was observed in 96.3% of the subjects. Serum 25(OH)D levels were significantly lower in winter in the second and third trimesters, while serum intact parathormone (iPTH) and alkaline phosphatase levels were significantly higher in winter in all three trimesters. A significant negative correlation was found between serum 25(OH)D and iPTH in mothers ($r = -0.367$, $P = 0.0001$) and infants ($r = -0.56$, $P = 0.0001$). A strong positive correlation was observed between 25(OH)D levels of mother–infant pairs ($r = 0.779$, $P = 0.0001$). A high prevalence of hypovitaminosis D was observed in pregnancy, lactation and infancy with no significant inter-trimester differences in serum 25(OH)D levels.

Key words: Pregnancy; Trimesters; Mother–infant pairs; Serum 25-hydroxyvitamin D; Parathormone

High prevalence of hypovitaminosis D has been established in all age groups across the world⁽¹⁾. The problem of hypovitaminosis D is likely to worsen during pregnancy because of the active transplacental transport of Ca to the developing fetus. Mother–offspring studies in Western populations have confirmed that optimal vitamin D supply not only influences the course of pregnancy, but is also required for fetal and neonatal Ca homeostasis, bone maturation and mineralisation^(2–6). Breast-fed infants born to vitamin D-deficient mothers are at risk for developing vitamin D deficiency and its metabolic sequelae^(7–12).

Divergent data on the status of 25-hydroxyvitamin D (25(OH)D) levels in different trimesters of pregnancy are available, with different investigators reporting either a decline⁽¹³⁾ or an increase⁽¹⁴⁾ or absence of change with progression of

pregnancy^(15,16). Furthermore, most studies have evaluated mothers in the third trimester and correlated their serum vitamin D levels with the newborn's cord blood 25(OH)D levels^(17–19). In view of the aforementioned facts, we have (1) evaluated maternal 25(OH)D levels in different trimesters, (2) assessed the impact of seasonal variation on serum vitamin D status, and (3) correlated maternal and newborn vitamin D status by concurrent evaluation of serum 25(OH)D levels in mother–infant pairs at 6–8 weeks postpartum.

Methods

Setting

Subjects were recruited between April 2006 and October 2007, from the obstetrics outpatient department of the Armed Forces

Abbreviations: 25(OH)D, 25-hydroxyvitamin D; ALP, alkaline phosphatase; iPTH, intact parathormone.

* **Corresponding author:** Dr R. K. Marwaha, fax +91 11 23939684, email marwaha_ramank@hotmail.com

Clinic and Army Hospital (Research and Referral), Delhi, which is a primary care provider for families of armed forces personnel currently residing in Delhi. The present study was conducted according to the guidelines laid down in the Declaration of Helsinki, and all procedures involving human subjects/patients were approved by the Institutional Human Ethics Committee at Army Hospital (Research and Referral). Written informed consent was obtained from all subjects/patients.

Subjects

Healthy women (n 541) with uncomplicated, single, intra-uterine gestation in any trimester were consecutively recruited, and anthropometric, nutritional, biochemical and hormonal investigations were carried out once at the time of first contact. All women who were approached agreed to participate in the study. These women, all of whom were housewives, belonged to lower–middle socio-economic strata, with 85% having completed 12 years of schooling. Fasting blood samples were drawn without venostasis under basal conditions. Serum was separated in a cold centrifuge, and three aliquots were made, one of which was used immediately to measure ionised and total Ca, inorganic P and serum alkaline phosphatase (ALP), while the other two were stored at -80°C for assessing 25(OH)D and intact parathormone (iPTH). Women with any chronic hepatic or renal illness, malabsorption syndrome, medications (current and past) and vitamin supplements that can affect the Ca–vitamin D–parathormone axis were excluded from the study.

All women who completed their pregnancy were invited 6–8 weeks postpartum for clinical and biochemical evaluation of mother–infant pairs. Of the 541 recruited pregnant women, only 342 mother–infant pairs could be studied. The remaining mothers were unavailable for comment, as they had gone back to their native villages after delivery, which is a common local tradition.

Hormonal assays

Serum 25(OH)D was measured by RIA using a commercial kit (Diasorin, Stillwater, MN, USA). The normal range for 25(OH)D was 22.5–92.5 nmol/l (9–37 ng/ml), with analytical sensitivity being 3.75 nmol/l (1.5 ng/ml). Serum iPTH was measured by immunoradiometric assay with a commercial kit (Diasorin). The normal range for iPTH was 13–54 pg/ml, with analytical sensitivity being 0.7 pg/ml. Commercial kits (Roche Diagnostics GmbH, Mannheim, Germany) were used to measure serum Ca, P and ALP. Total Ca was estimated by the colorimetric method. The normal range for total Ca was 2.24–2.74 mmol/l (90–110 mg/l) in infants (2 d–2 years old) and 2.09–2.54 mmol/l (84–102 mg/l) in adults, with analytical sensitivity being 2 mg/l. Serum P and ALP were determined by photometric analysis. The normal range for P was 0.97–2.25 mmol/l (30–70 mg/l) in infants and 0.87–1.45 mmol/l (27–45 mg/l) in adults, with analytical sensitivity being 3 mg/l. The normal upper limit of ALP was 1076 IU/l in infants and \leq 240 IU/l in non-pregnant women.

The analytical sensitivity of ALP was 5 IU/l. Serum ionised Ca was estimated by the ion exchange method and its normal range was 1.12–1.32 mmol/l (44.8–5.280 mg/l) in adults and 1–1.25 mmol/l (40–50 mg/l) in infants. Vitamin D deficiency was classified using Lips criteria⁽²⁰⁾ based on 25(OH)D levels as mild (25–50 nmol/l (10–20 ng/ml)), moderate (12.5–25 nmol/l (5–10 ng/ml)) and severe ($<$ 12.5 nmol/l (5 ng/ml)) hypovitaminosis D.

Dietary analysis

Nutrient intake was calculated using the 24 h dietary recall method. During pre-testing, three separate 24 h dietary recalls were recorded from fifty subjects (two on weekdays and one on a weekend). Since no difference was found between weekday and weekend intakes, only one 24 h dietary recall was taken during the final study. Detailed descriptions of foods consumed along with their quantities, as estimated by standardised household measures, were noted. Raw weights were then calculated and used to estimate nutrient intake using the Nutritive Value of Indian foods (National Institute of Nutrition, 2001)⁽²¹⁾.

Statistical analysis

Data were analysed using STATA-9.0 (Stata Corp LP, College Station, TX, USA). Descriptive statistics are expressed as numbers (percentages) or means and standard deviations/medians (ranges) as appropriate. Seasonal differences in biochemical parameters were tested using Student's *t* test and Wilcoxon's rank-sum test for non-normal data. Spearman's rank correlation coefficient was used to determine the strength of the relationship between variables, since data were non-normal. $P < 0.05$ was considered significant.

Results

The basic characteristics of women are given in Table 1. The mean age of pregnant women was 24.6 (SD 2.8) (range 19–30) years. The mean age at marriage was 20.3 (SD 1.5) years. There were 219 (40.5%) women with their first pregnancy.

Vitamin D status of pregnant women

A total of 521 women (96.3%) were found to be vitamin D deficient (25(OH)D $<$ 50 nmol/l), with 36.8, 41.8 and 17.7% falling into the mild (25–50 nmol/l), moderate (12.5–25 nmol/l) and severe ($<$ 12.5 nmol/l) hypovitaminosis D categories, respectively. Mean serum total Ca, ionised Ca, P, alkaline phosphate, 25(OH)D and iPTH were 2.33 (SD 0.09) mmol/l, 1.19 (SD 0.05) mmol/l, 1.22 (SD 0.15) mmol/l, 182.07 (SD 40.51) IU/l, 23.2 (SD 12.2) nmol/l and 649 (SD 44) pg/l, respectively. A highly significant negative correlation was observed between vitamin D and iPTH (r -0.317 , $P=0.001$) and between vitamin D and ALP (r -0.232 , $P=0.0001$). Seasonal differences observed in various biochemical and hormonal parameters during the three trimesters are shown in Table 2. In comparison with the values reported in summer, serum 25(OH)D levels were

Table 1. Basic parameters of pregnant women(Mean values, standard deviations, medians, ranges, number of subjects and percentages, *n* 541)

Parameters	First trimester (<i>n</i> 156)			Second trimester (<i>n</i> 218)			Third trimester (<i>n</i> 167)		
	Mean	SD	Range	Mean	SD	Range	Mean	SD	Range
Age (years)	24.4	2.67	19–30	25.04	2.94	19–30	24.26	2.82	19–30
Age of marriage (years)	20.35	1.62	18–26	20.35	1.54	18–26	20.05	1.46	18–26
BMI (kg/m ²)	20.25	3.17	14.27–26.31	22.25	3.16	15.31–30.68	25.19	3.16	19.14–31.8
Income (rupees)	7762	1469	5000–12 000	7959	1464	5000–12 000	7829	1464	5000–12 000
Hb (g%)	10.3	1.26	8.4–14.6	10.36	1.55	8.2–14.6	11.05	7.86	8.2–14.5
Sun exposure (min)									
Median	15		10–60	25		10–60	9		10–20
UV score									
Median	7.5		3.75–45	15		3.75–45	15.6		3–45
	<i>n</i>	%		<i>n</i>	%		<i>n</i>	%	
Members in the family									
Frequency									
2	91	58.3		90	41.2		82	49.1	
3	61	39.1		113	51.8		74	44.3	
> 3	4	2.5		15	6.8		11	6.5	
Parity									
1	74	47.4		78	35.7		67	40.1	
2	50	32.0		83	38.0		62	37.1	
> 2	32	20.5		57	26.1		38	22.1	

significantly lower in winter in the second and third trimesters, while iPTH and ALP levels were significantly higher in winter in all the three trimesters.

Among the women studied either in summer or winter, while there was no significant difference in mean serum 25(OH)D levels between the three trimesters, iPTH levels were significantly higher in the first trimester compared with values in both the second and third trimesters. The prevalence of maternal hypovitaminosis D was not different in the three trimesters whether studied in summer (96.9 *v.* 92 *v.* 98.7%) or winter (100 *v.* 97.9 *v.* 95.6%) in the first, second and third trimesters, respectively. No significant difference was observed in mean serum phosphate levels in the three different trimesters both in summer and winter. Mean serum ALP showed a progressive decline in summer, whereas in winter, it decreased in the second trimester and then increased in the third trimester, the difference being non-significant in both seasons.

There was no difference in serum Ca, phosphate, ALP and 25(OH)D levels between primigravida and multigravida. However, serum iPTH levels were marginally higher in primigravida (69.57 (SD 43.95) pg/ml) compared with those in multigravida (61.72 (SD 43.94) pg/ml; *P*=0.023).

Diet

The dietary intake of energy (5300 (SD 1130) kJ), total Ca (408 (SD 160) mg; range 38–1024 mg) and Ca from dairy sources (271 (SD 154) mg) in pregnant women was significantly lower when compared with the RDA for Indians⁽²²⁾. The mean vitamin D intake was similar in all trimesters (0.2 (SD 0.4) µg, *P*=0.16), being higher in winter than in summer, but did not reach statistical significance (0.3 (SD 0.4) *v.* 0.1 (SD 0.3) µg), *P*=0.06). Although mean energy intake increased from 5154 (SD 1087) kJ in the first trimester to 5314 (SD 1132) kJ in the second trimester and to 5450 (SD 1185) kJ in the third

trimester, these differences were not statistically significant (*P*=0.06). Percentage energy contribution was highest from carbohydrates (181 (SD 49) g; 64%) followed by fat (45 (SD 10) g; 32%) and protein (35 (SD 12) g; 11%), which were well in the recommended range.

Vitamin D status of lactating mothers

The biochemical profile of lactating mothers is presented in Table 3. A total of 341 (99.7%) lactating mothers had serum 25(OH)D levels < 50 nmol/l, with 19.3, 51.2 and 29.2% suffering from mild (25–50 nmol/l), moderate (12.5–25 nmol/l) and severe (< 12.5 nmol/l) hypovitaminosis D, respectively. A highly significant negative correlation was found between 25(OH)D and iPTH (*r* −0.310, *P*=0.0001) and between 25(OH)D and ALP (*r* −0.217, *P*=0.0001), respectively, in lactating mothers.

Vitamin D status of exclusively breast-fed infants

The biochemical profile of infants is shown in Table 3. No significant difference was observed in serum 25(OH)D levels in infants born in summer and winter (data not shown). A total of 338 infants (98.8%) had serum 25(OH)D levels < 50 nmol/l, with 38.0, 44.5 and 16.3% classified as mild (25–50 nmol/l), moderate (12.5–25 nmol/l) and severe (< 12.5 nmol/l) hypovitaminosis D, respectively. A highly significant negative correlation was also observed between 25(OH)D and iPTH levels of infants (*r* −0.56, *P*=0.0001; data not shown).

Correlations between the vitamin D status of mother–infant pairs

As shown in Fig. 1, a strong positive correlation was found between 25(OH)D, (*r* 0.779, *P*=0.0001), ionised Ca (*r* 0.166,

Table 2. Seasonal differences between biochemical parameters in the three trimesters of pregnancy (Mean values, standard deviations, medians, ranges and number of subjects)

Parameters	Summer		Winter		P
	Mean	SD	Mean	SD	
First trimester					
<i>n</i>	97		59		
Ca (mmol/l)*	2.37	0.09	2.36	0.10	0.547
PO ₄ (mmol/l)*	1.25	0.14	1.20	0.16	0.049
ALP (IU/l)*	177.45	43.78	195.88	45.00	0.012
Ionised Ca (mmol/l)*	1.190	0.045	1.200	0.059	0.205
25(OH)D (nmol/l)†	23.4	11.3	19.6	9.2	0.085
iPTH (pg/l)‡	687	502	872	445	0.003
Median	730		810		
Range	71.5–3611		134.7–2200		
Second trimester					
<i>n</i>	125		93		
Ca (mmol/l)*	2.38	0.09	2.38	0.09	0.884
PO ₄ (mmol/l)*	1.24	0.14	1.17	0.15	0.0003
ALP (IU/l)*	176.28	41.23	189.12	40.02	0.022
Ionised Ca (mmol/l)*	1.170	0.044	1.180	0.050	0.009
25(OH)D (nmol/l)†	25.7	15.1	20.2	10.6	0.0009
iPTH (pg/l)‡	509	370	763	444	0.0001
Median	361		738.6		
Range	96.5–1393.6		91.2–2180		
Third trimester					
<i>n</i>	77		90		
Ca (mmol/l)*	2.39	0.09	2.38	0.09	0.318
PO ₄ (mmol/l)*	1.21	0.16	1.22	0.15	0.83
ALP (IU/l)*	165.91	23.41	192.58	38.65	0.0001
Ionised Ca (mmol/l)*	1.19	0.05	1.20	0.06	0.053
25(OH)D (nmol/l)†	27.7	9.2	21.1	12.4	0.0001
iPTH (pg/l)‡	482	351	682	413	0.0009
Median	380		595		
Range	100–1688		75.5–1650		

ALP, alkaline phosphatase; 25(OH)D, 25-hydroxyvitamin D; iPTH, intact parathormone.

* Mean values were significantly different as tested by the independent *t* test between summer and winter: $P < 0.001$, $P < 0.01$, $P < 0.05$.

† Mean values were not significantly different between trimesters: $P = 0.81$ (first v. second trimester), $P = 0.12$ (second v. third trimester), $P = 0.07$ (first v. third trimester).

‡ Mean values were significantly different as tested by Wilcoxon's rank-sum test between summer and winter: $P < 0.001$ (first trimester), $P < 0.01$ (second trimester), $P < 0.05$ (third trimester).

$P = 0.0001$) and iPTH ($r = 0.534$, $P = 0.0001$) levels of mothers and infants.

Discussion

We have reported vitamin D status of pregnant women hailing from lower–middle socio-economic strata. The nutritional, educational and obstetric data of these women were consistent with that described for this socio-economic class⁽²³⁾, thereby making the information generated generalisable for this group. In the present study, 96% of pregnant women had hypovitaminosis D, which is the highest reported prevalence in the literature. Several other studies from developing and developed nations across the world have reported that the prevalence of hypovitaminosis D (25(OH)D < 25 nmol/l) in pregnancy ranged from 18 to 84%^(24–26,10,27–32). South Asians, both in their country of origin and after migration to Europe or the UK, have been found to have lower serum 25(OH)D concentrations than white Caucasians^(26,33–35) due to a range of factors including skin pigmentation, covered-up clothing (especially common in women), restricted outdoor physical activity and low dietary vitamin D intake^(25,36).

The present findings have shown the status of 25(OH)D, Ca, ALP and iPTH during different trimesters of pregnancy. There was no significant difference in the prevalence of 25(OH)D deficiency (25(OH)D < 50 nmol/l) among pregnant women in the three different trimesters, both in summer and winter. A study of pregnant Iranian women has shown that

Table 3. Biochemical profile of mothers and their infants at 6 weeks postpartum (Mean values, standard deviations, medians and ranges, *n* 342)

Parameters	Mothers		Infants	
	Mean	SD	Mean	SD
Ca (mmol/l)	2.37	0.09	2.39	0.09
PO ₄ (mmol/l)	1.25	0.13	1.26	0.13
ALP (IU/l)	213.20	42.41	702.39	158.07
Ca ²⁺ (mmol/l)	1.18	0.05	1.18	0.05
25(OH)D (nmol/l)	19.6	8.3	22.3	10.5
iPTH (pg/l)				
Median	655		583	
Range	102–1672		150–1863.2	

ALP, alkaline phosphatase; 25(OH)D, 25-hydroxyvitamin D; iPTH, intact parathormone.




Fig. 1. Relationship between the serum 25-hydroxyvitamin D (25(OH)D) levels of mother–infant pairs, (r 0.779, P =0.0001).

60% of the women in the first trimester, 48% in the second trimester and 47% in the third trimester had either severe or moderate vitamin D deficiency⁽³⁷⁾. In an earlier study conducted in Asian women residing in London, it has been found that 25(OH)D concentration was <25 nmol/l in 25% of subjects in the first trimester, which reduced in the third trimester⁽¹⁷⁾.

Total serum Ca and ionised Ca values showed no variation across trimesters in the present study. Studies in the literature have shown either no change in serum Ca values^(15,16,37) or an increase^(17,38–40) or a decrease^(41,42) with progression of pregnancy. The constancy of serum ionised Ca values has also been reported by other investigators^(15,43,44). Similar to earlier reports^(15,37,43,44), there was no change in serum P during the course of pregnancy both in summer and winter. Mean serum ALP showed a non-significant decline in the present study. This is in contrast with the findings of Ainy *et al.*⁽³⁷⁾ who attributed the increment in ALP, related to placental production, to lack of vitamin D supplementation and insufficient dietary intake. We observed no significant difference in mean serum 25(OH)D concentration in the three trimesters, both in summer and winter, which is in concordance with the reports from Reddy *et al.*⁽¹⁵⁾ and Selly *et al.*⁽¹⁶⁾. In contrast, Sanchez *et al.*⁽¹⁴⁾ found that 25(OH)D concentration increased in the second and third trimesters, the increase being attributable to food and supplement intake and sun exposure. In a longitudinal study of pregnant women, Ardawi *et al.*⁽¹³⁾ showed a moderate, but statistically significant, decrease towards the end of pregnancy and at term. The decrease was attributed to the particular dietary and cultural habits followed by the subjects. In India, there is no fortification of food products with vitamin D, and there is no clear guidelines recommending mandatory vitamin D supplementation during pregnancy. These factors could partly explain the absence of variation in serum 25(OH)D levels during the three trimesters of pregnancy.

Several reasons for change in serum 25(OH)D levels in pregnancy have been postulated. These include altered hepatic 25-hydroxylase activity, change in iPTH levels and

increased fetal metabolic activity^(14,43,45–47). Another study has suggested that rise in iPTH was responsible for the increased absorption of vitamin D in mothers⁽⁴⁸⁾.

There is no consistent pattern in the change in serum iPTH levels during the different trimesters of pregnancy. Most studies conducted in populations replete with Ca and vitamin D have reported a gradual decline in serum iPTH levels with evolution of pregnancy. In contrast, studies from the Gambia, Asia and other regions with low Ca and vitamin D intake often do not report any decline in iPTH levels during pregnancy^(49,50). Other causes of varying results could include methodological differences in assays resulting in the measurement of multiple different immunoreactive but biologically inactive fragments of parathormone⁽⁴⁹⁾. In addition, the contribution of placenta-derived parathormone-related peptide to the different aspects of bone mineral metabolism, including renal 1 α hydroxylation of 25(OH)D, may also be partly responsible for the variation in iPTH values in the three trimesters reported in different studies^(49,51–53).

Marked seasonal variation in serum 25(OH)D levels was observed in the present study. A progressive fall in 25(OH)D levels in pregnant women during winter months due to the reduced availability of sunshine has been described in European and US populations^(54–57), as well as from India and other Asian countries^(32,34,37,58,59). In addition, serum 25(OH)D concentration also depends on the extent of the body surface area exposed, which is likely to be reduced due to the style of dressing in winter^(8,60).

High prevalence of vitamin D deficiency in apparently healthy lactating mothers (99.7%) and exclusively breast-fed infants (98.8%) observed in the present study only reiterates our earlier observation⁽¹²⁾ as well as those of other workers^(37,60–63). Mothers with suboptimal vitamin D status have offspring with reduced intra-uterine and postnatal skeletal development^(7,10). The impact of maternal vitamin D status on the neonate's serum 25(OH)D levels is apparent from the strong correlation reported by us in the present study. Although a similar correlation between 25(OH)D levels of mothers and newborns has been reported earlier, most investigators have measured cord blood 25(OH)D levels to establish the relationship^(18,19,31).

Mother–offspring studies in Western populations have shown associations of maternal body build, diet, nutritional status, smoking and physical activity with bone mass in newborns and children^(2–4,5,7,10). The importance of nutrition, mainly Ca, has been acknowledged with regard to pregnancy outcome⁽⁶⁴⁾. Greater maternal consumption of Ca and Ca-rich foods, especially milk and milk products, in mid- to late pregnancy has been associated with improved bone outcomes in children⁽⁹⁾. In the present study, mothers had low Ca intakes, consistent with other low-income groups in India⁽⁶⁵⁾. The mean dietary Ca intake (408.11 (SD 167.2) mg/d) of mothers was just 60% of the RDA given by the Indian Council of Medical Research⁽²²⁾, and the intake of other macronutrients was also far below the RDA. Also, vitamin D supplementation is not a part of antenatal care programmes in India, which worsens the situation further.

These data reinforce the need to provide greater emphasis on maternal nutrition to improve neonatal and childhood bone health.

Conclusion

We conclude that there is a high prevalence of hypovitaminosis D among pregnant women and their infants in India. Serum 25(OH)D levels were uniformly low across all three trimesters, with a tendency to decline in winter. There was a strong positive correlation between maternal and infant serum 25(OH)D levels. Further research, preferably by randomised controlled trials, is needed to establish the effects of vitamin D supplementation during pregnancy on the bone health of women and their children.

Acknowledgements

The present study was funded through project no. INM305, from the Defence Research and Development Organization, Ministry of Defence, Government of India. The authors would like to acknowledge the assistance provided by Madan Prasad, MI Beg, Abhishek Kaushik, Amit Panwar, Pramod Kumar and Neeta Rautela for the conduct of the present study. We would also like to express our gratitude to the study volunteers and staff of the Armed Forces Clinic, Dalhousie Road, New Delhi. None of the authors has a conflict of interests to declare. The authors' contributions were as follows: R. K. M. and N. T. contributed to the conceptualisation of the study, clinical evaluation and preparation of the manuscript. S. C. supervised the recruitment and clinical evaluation of the subjects. N. A. was involved in the data analysis and preparation of the manuscript. M. K. G. was responsible for conceptualising the study and clinical evaluation of the subjects. B. S. contributed to conceptualisation of the study and data collection. R. S. K. assisted in the clinical evaluation. K. B. and S. S. performed the collection of biochemical samples and laboratory evaluation. K. M. analysed the data. S. P. conducted the data collection and dietary analysis. The study has been approved by the Institutional Ethics Committee of Army Hospital (Research and Referral), Delhi Cantt, India.

References

- Mithal A, Wahi DA, Bonjour JP, *et al.* (2009) Global vitamin D status and determinants of hypovitaminosis D. *Osteoporos Int* **20**, 1807–1829.
- Jones G, Riley M & Dwyer T (1999) Maternal smoking during pregnancy, growth and bone mass in prepubertal children. *J Bone Miner Res* **14**, 146–151.
- Jones G, Riley MD & Dwyer T (2000) Maternal diet during pregnancy is associated with bone mineral density in children: a longitudinal study. *Eur J Clin Nutr* **54**, 749–756.
- Godfrey K, Walker-Bone K, Robinson S, *et al.* (2001) Neonatal bone mass: influence of parental birthweight and maternal smoking, body composition and activity during pregnancy. *J Bone Miner Res* **16**, 1694–1703.
- Tobias JH, Steer CD, Emmett PM, *et al.* (2005) Bone mass in childhood is related to maternal diet in pregnancy. *Osteoporos Int* **16**, 1731–1741.
- Walicka M & Marcinowska-Suchowierska E (2008) Vitamin D deficiency during pregnancy and lactation. *Ginek Pol* **79**, 780–784.
- Pawley N & Bishop NJ (2004) Prenatal and infant predictors of bone health: the influence of vitamin D. *Am J Clin Nutr* **80**, Suppl., S1748–S1751.
- Hatun S, Ozkan B, Orbak Z, *et al.* (2005) Vitamin D deficiency in early infancy. *J Nutr* **135**, 279–282.
- Ganpule A, Yajnik CS, Fall CHD, *et al.* (2006) Bone mass in Indian children – relationships to maternal nutritional status and diet during pregnancy: the Pune Maternal Nutrition Study. *J Clin Endocrinol Metab* **91**, 2994–3001.
- Javaid MK, Crozier SR, Harvey NC, *et al.* (2006) Maternal vitamin D status during pregnancy and childhood bone mass at age nine years: a longitudinal study. *Lancet* **367**, 36–43.
- Dijkstra SH, Beek AV, Janssen JW, *et al.* (2007) High prevalence of vitamin D deficiency in newborn infants of high-risk mothers. *Arch Dis Child* **92**, 750–753.
- Seth A, Marwaha RK, Singla B, *et al.* (2009) Vitamin D nutrition status of exclusively breast fed infants and their mothers. *J Pediatr Endocrinol Metab* **22**, 241–246.
- Ardawi MSM, Nasrat HAN & BA'Aqueel HS (1997) Calcium regulating hormones and parathyroid hormone-related peptide in normal human pregnancy and postpartum: a longitudinal study. *Eur J Endocrinol* **137**, 402–409.
- Sanchez PA, Idrisa A, Bobzom DN, *et al.* (1997) Calcium and vitamin D status of pregnant teenagers in Maiduguri Nigeria. *J Natl Med Assoc* **89**, 805–811.
- Reddy SG, Norman AW, Willis DM, *et al.* (1983) Regulation of vitamin D metabolism in normal human pregnancy. *J Clin Endocrinol Metab* **56**, 363–370.
- Selly EW, Brown EM, DeMaggio DM, *et al.* (1997) A prospective study of calcitropic hormones in pregnancy and post partum: reciprocal changes in serum intact parathyroid hormone and 1,25-dihydroxyvitamin D. *Am J Obstet Gynecol* **176**, 214–217.
- Brooke OG, Brown IRF, Cleeve HJW, *et al.* (1981) Observations on the vitamin D state of pregnant Asian women in London. *Br J Obstet gynaecol* **88**, 18–26.
- Marya RK, Rathee S, Dua V, *et al.* (1988) Effect of vitamin D supplementation during pregnancy on foetal growth. *Ind J Med Res* **88**, 488–492.
- Bhalala U, Desai M, Parekh P, *et al.* (2007) Subclinical hypovitaminosis D among exclusively breastfed young infants. *Indian Pediatr* **44**, 897–901.
- Lips P (2001) Vitamin D deficiency and secondary hyperparathyroidism in the elderly, consequences for bone loss and fractures and therapeutic implications. *Endocr Rev* **22**, 477–501.
- Gopalan C, Ramasastry BV & Balasubramaniam SC (2001) *Nutritive Value of Indian Foods*. Hyderabad: Indian Council of Medical Research (ICMR) Publication.
- Indian Council of Medical Research (1990) *Nutrient Requirements and Recommended Dietary Allowances for Indians*. New Delhi: ICMR Publication.
- Sachar RK, Kaur Navjeet, Soni RK, *et al.* (2000) Energy consumption during pregnancy and its relationship to birth weight. A population based study from rural Punjab. *Indian J Commun Med* **25**, 166–169.
- Bassir M, Laborie S, Lapillonne A, *et al.* (2001) Vitamin D deficiency in Iranian mothers and their neonates: a pilot study. *Acta Paediatr* **90**, 577–579.
- Sachan A, Gupta R, Das V, *et al.* (2005) High prevalence of vitamin D deficiency among pregnant women and their newborns in northern India. *Am J Clin Nutr* **81**, 1060–1064.

26. Scroth RJ, Lavelle CLB & Moffatt MEK (2005) A review of vitamin D deficiency during pregnancy: who is affected? *Int J Circumpolar Health* **64**, 112–120.
27. Judkins A & Eagleton C (2006) Vitamin D deficiency in pregnant New Zealand women. *N Z Med J* **119**, U2144.
28. Van der Meer IM, Karamali NS, Boeke JP, *et al.* (2006) High prevalence of vitamin D deficiency in pregnant non-western women in The Hague, Netherlands. *Am J Clin Nutr* **84**, 350–353.
29. Cavalier E, Delanaye P, Morreale A, *et al.* (2008) Vitamin D deficiency in recently pregnant women. *Rev Med Liege* **63**, 87–91.
30. O'Riordan MN, Kiely M, Higgins JR, *et al.* (2008) Prevalence of suboptimal vitamin D status during pregnancy. *Ir Med J* **101**, 240–243.
31. Farrant HJW, Krishnaveni GV, Hill JC, *et al.* (2009) Vitamin D insufficiency is common in Indian mothers but is not associated with gestational diabetes or variation in newborn size. *Eur J Clin Nutr* **63**, 646–652.
32. Sahu M, Bhatia V, Aggarwal A, *et al.* (2009) Vitamin D replacement in pregnant women in rural north India: a pilot study of vitamin D replacement in pregnant Indian women. *Eur J Clin Nutr* **63**, 1157–1159.
33. Awumey EMK, Mitra DA, Hollis BW, *et al.* (1998) Vitamin D metabolism is altered in Asian Indians in the Southern United States: a clinical research center study. *J Clin Endocrinol Metab* **83**, 169–173.
34. Goswami R, Gupta N, Goswami D, *et al.* (2000) Prevalence and significance of low 25-hydroxyvitamin D concentrations in healthy subjects in Delhi. *Am J Clin Nutr* **72**, 472–475.
35. Hamson C, Goh L, Sheldon P, *et al.* (2003) Comparative study of bone mineral density, calcium, and vitamin D status in the Gujarati and white populations of Leicester. *Postgrad Med J* **79**, 279–283.
36. Masood SH & Iqbal MP (2008) Prevalence of vitamin D deficiency in South Asia. *Pak J Med Sci* **24**, 891–897.
37. Ainy E, Ghazi AAM & Azizi F (2006) Changes in calcium, 25(OH) vitamin D₃ and other biochemical factors during pregnancy. *J Endocrinol Invest* **29**, 303–307.
38. Brooke OG, Brown IRF, Bone CDM, *et al.* (1980) Vitamin D supplements in pregnant Asian women: effects on calcium status and fetal growth. *Br Med J* **80**, 751–754.
39. Kohlmeier L & Marcus R (1995) Calcium disorders of pregnancy. *Endocrinol Metab Clin North Am* **24**, 15–39.
40. Brunvand L, Quigstad E, Urdal P, *et al.* (1996) Vitamin D deficiency and fetal growth. *Early Hum Dev* **45**, 27–33.
41. Polanska N, Dale RA & Wills MR (1976) Plasma calcium levels in pregnant Asian women. *Ann Clin Biochem* **13**, 339–344.
42. Henriksen C, Brunvand L, Stoltenberg C, *et al.* (1995) Diet and vitamin D status among pregnant Pakistani women in Oslo. *Eur J Clin Nutr* **49**, 211–218.
43. Pitkin RM (1985) Calcium metabolisms in pregnancy and the perinatal period: a review. *Am J Obstet Gynecol* **151**, 99–109.
44. Saggese G, Baroncelli GI, Bertelloni S, *et al.* (1991) Intact parathyroid hormone levels during pregnancy, in healthy term neonates and in hypocalcemic preterm infants. *Acta Paediatr Scand* **80**, 36–41.
45. Cushard WG, Creditor M, Canterbury JM, *et al.* (1972) Physiologic hyperparathyroidism in pregnancy. *J Clin Endocrinol Metab* **34**, 767–771.
46. Turton CWG, Stamp TCB, Stanley P, *et al.* (1977) Altered vitamin D metabolism in pregnancy. *Lancet* **i**, 222–225.
47. Marya RK, Rathee S, Lata V, *et al.* (1981) Effects of vitamin D supplementation in pregnancy. *Gynecol Obstet Invest* **12**, 155–161.
48. Bruinse HW & Van den Berg H (1995) Changes of some vitamin levels during and after normal pregnancy. *Eur J Obstet Gynecol Reprod Biol* **61**, 31–37.
49. Kovacs CS & Kronenberg HM (1997) Maternal–fetal calcium and bone metabolism during pregnancy, puerperium and lactation. *Endocr Rev* **18**, 832–872.
50. Kovacs CS (2008) Vitamin D in pregnancy and lactation: maternal, fetal, and neonatal outcomes from human and animal studies. *Am J Clin Nutr* **88**, 520S–528S.
51. Gallacher SJ, Fraser WD, Owens OJ, *et al.* (1994) Changes in calcitropic hormones and biochemical markers of bone turnover in normal human pregnancy. *Eur J Endocrinol* **131**, 369–374.
52. Hosking DJ (1996) Calcium homeostasis in pregnancy. *Clin Endocrinol* **45**, 1–6.
53. Tobias JH & Cooper C (2004) PTH/PTHrP activity and the programming of skeletal development *in utero*. *J Bone Miner Res* **19**, 177–182.
54. MacLaughlin M, Fairney A, Lester E, *et al.* (1974) Seasonal variations in serum 25-hydroxy-cholecalciferol in healthy people. *Lancet* **i**, 536–538.
55. Kuoppala T, Tuimala R, Parviainen M, *et al.* (1986) Serum levels of vitamin D metabolites, calcium, phosphorus, magnesium and alkaline phosphatase in Finnish women throughout pregnancy and in cord serum at delivery. *Hum Nutr Clin Nutr* **40**, 287–293.
56. Sherman SS, Hollis BW & Tobin JD (1990) Vitamin D status and related parameters in a healthy population: the effects of age, sex, and season. *J Clin Endocrinol Metab* **71**, 405–413.
57. Van der Wielen RP, Lowik MR, Van den Berg H, *et al.* (1995) Serum vitamin D concentrations among elderly people in Europe. *Lancet* **345**, 207–210.
58. Kim JH & Moon SJ (2000) Time spent outdoors and seasonal variation in serum concentrations of 25-hydroxyvitamin D in Korean women. *Int J Food Sci Nutr* **51**, 439–451.
59. Nakamura K, Nashimoto M & Yamamoto M (2000) Summer/winter differences in the serum 25 hydroxyvitamin D₃ and parathyroid hormone levels of Japanese women. *Int J Biometeorol* **44**, 186–189.
60. Puri S, Marwaha RK, Agarwal N, *et al.* (2008) Vitamin D status of apparently healthy schoolgirls from two different socioeconomic strata in Delhi: relation to nutrition and lifestyle. *Br J Nutr* **99**, 876–882.
61. MacLennan WJ, Hamilton JC & Darmady JM (1980) The effect of season and stage of pregnancy on plasma 25-hydroxy-vitamin D concentrations in pregnant women. *Postgrad Med J* **56**, 75–79.
62. Dawodu A, Agarwal M, Hossain M, *et al.* (2003) Hypovitaminosis D and vitamin D deficiency in exclusively breast-feeding infants and their mothers in summer: a justification for vitamin D supplementation of breast-feeding infants. *J Pediatr* **142**, 169–173.
63. Dawodu A & Wagner CL (2007) Mother–child vitamin D deficiency: an international perspective. *Arch Dis Child* **92**, 737–740.
64. Balasubramanian K, Rajeshwari J & Gulab (2003) Varying role of vitamin D deficiency in the etiology of rickets in young children vs. adolescents in northern India. *J Trop Pediatr* **49**, 201–206.
65. Shatruguna V, Kulkarni B, Kumar PA, *et al.* (2005) Bone status of Indian women from a low-income group and its relationship to nutritional status. *Osteoporos Int* **16**, 1827–1835.