

# A *Spitzer*-IRS search for the galaxies that re-ionized the Universe

Mark Lacy<sup>1</sup>, Andrew J. Bunker<sup>2</sup>, Jean-Paul Kneib<sup>3</sup>  
and Harry I. Teplitz<sup>1</sup>

<sup>1</sup>Spitzer Science Center, California Institute of Technology, Pasadena, CA 91125, USA  
email: mlacy,hit@ipac.caltech.edu

<sup>2</sup>School of Physics, University of Exeter, Stocker Road, Exeter, EX4 4QL, UK  
email: bunker@astro.ex.ac.uk

<sup>3</sup>Observatoire Midi-Pyrénées, UMR 5572, 14 Avenue Edouard Belin, F-31000 Toulouse, France  
email: jean-paul.kneib@oamp.fr

**Abstract.** We describe an observation designed to find H $\alpha$  emission from galaxies at  $z \simeq 7-12$  made using the Infrared spectrograph (IRS) on the *Spitzer Space Telescope*.

**Keywords.** galaxies: formation,galaxies: clusters: individual (Abell 2218)

---

Spectra of the most distant quasars at  $z \simeq 6.5$  show the Gunn-Peterson trough caused by absorption by neutral hydrogen in the ISM (White *et al.* 2003). The most recent analysis of the WMAP dataset is consistent with re-ionization at  $z \simeq 11$  (Page *et al.* 2007). These observations suggest that starburst galaxies or quasars with large amounts of escaping UV emission were present at  $z \simeq 7-12$ , but such a population is inconsistent with a straightforward extrapolation of the evolution of known galaxy or quasar populations at  $z \simeq 6$  (e.g., Bunker *et al.* 2004), unless the dominant stellar populations at  $z \simeq 6$  is very metal poor (Stiavelli *et al.* 2004). Prior to re-ionization, Ly $\alpha$  emission is expected to be extinguished by a large (though uncertain) factor  $\sim 10-100$  by the damping wing of the Gunn-Peterson trough (e.g., Santos 2006). This makes blank field H $\alpha$  searches with *Spitzer* competitive with ground-based near-infrared searches for Ly $\alpha$ .

We used the on *Spitzer*-IRS spectrograph to observe a single slit position aligned along the critical line for  $z \gtrsim 7$  galaxies in the cluster Abell 2218 to demonstrate the feasibility of such observations. We used the second order short-low spectrum to search for H $\alpha$  emission lines redshifted into the  $5-9 \mu\text{m}$  range. We achieved a  $3\sigma$  detection limit  $\sim 2 \times 10^{-19} \text{ WHz}^{-1} \text{ m}^{-2}$ , corresponding to star formation rates  $\sim 1000/\mu\text{M}_{\odot} \text{ yr}^{-1}$  where  $\mu \simeq 10-100$  is the magnification factor. We have one candidate emission line detection at this level, which, if it is indeed H $\alpha$ , is at a wavelength corresponding to  $z = 9.8$ .

## Acknowledgements

This work is based on observations made with the *Spitzer Space Telescope*, which is operated by the Jet Propulsion Laboratory, California Institute of Technology under a contract with NASA.

## References

- Bunker, A. J., Stanway, E. R., Ellis, R. S., & McMahon, R. G. 2004, *MNRAS*, 355, 374  
Page, L., Hinshaw, G., Komatsu, E., *et al.* 2007, *ApJS*, 170, 335  
Santos, M. R. 2006, *MNRAS*, 349, 1137  
Stiavelli, M., Fall, S. M., & Panagia, N. 2004, *ApJ* (Letters), 610, L1  
White, R. L., Becker, R. H., Fan, X., & Strauss, M. A. 2003, *AJ*, 126, 1