

Calculation of Perturbations in the Stellar Three-Body Problem

J. A. DOCOBO, C. PRIETO, & J. F. LING

Observatorio Astronómico Ramón M. Aller, Universidad de Santiago,
P.O. Box 197, Santiago de Compostela, Spain

ABSTRACT: The method of perturbations of Deprit has been applied to the stellar three-body problem. The calculation, developed previously by the authors in other articles, is here systematized for easy use and applied to various triple systems.

1. INTRODUCTION

The stellar three-body problem (Harrington 1968a,1969; Docobo 1977; Abad 1984; Docobo & Prieto 1988; Ling 1989) is concerned with the motion of three comparable masses m_0 , m_1 and m_2 , such that m_2 lies far enough from the others to be able to take the motion of m_1 about m_0 and that of m_2 about the center of mass of m_0 and m_1 as perturbed Keplerian orbits.

This problem has lately been approached in several ways: hypothesis of Keplerian outer orbit; study of the perturbations of the outer orbit; integration of the equations of motion in various systems of variables; use of methods of analytical, semianalytical or numerical integration, etc.

In this paper, following R.S. Harrington, we have operated with the equations of motion in their Hamiltonian form, while their integration was approached by using the perturbation theory of Deprit (1969), in which the application of the Lie transformation to the initial canonical system gives canonical systems more amenable to integration. There is no unique way to apply this method, since the angular variables that appear in the Hamiltonian function may be eliminated with a single canonical transformation, or one by one using two or three transformations. The constants of integration may, furthermore, be recalculated at fixed time intervals or those obtained initially can be used throughout the process.

Experience has shown us that the results obtained using these forms of applying the method are in close agreement, therefore we decided to jointly eliminate the angular variables and recalculate the constants of integration at intervals of time coinciding with the non-perturbed period of the inner orbit. We have considered up to the third-order term in the small parameter in powers from which the Hamiltonian function is developed; that is, we have only used the term that most contributes to the perturbations of both orbits.

The software obtained, which we call Program Pertur, is here applied to the well-known systems ξ Ursae Majoris (ADS 8119 — WDS 11182N 3133) and ζ Aquarii (ADS 15971 — WDS 22288S 0002).

2. APLICATION TO ξ URSAE MAJORIS

This system is really a quadruple star formed of two subsystems Aa and Bb . Stars B and b form a spectroscopic binary of period 3.98 days and so, to a

first approximation, this subsystem can be considered as a single star, the total system becoming a triple star with two close members A and a , and a distant member Bb .

ζ Ursae Majoris is probably the system which best satisfies the requirements to be able to study perturbations in its motion. Because of this, it was used in earlier studies, chief among these being those of Brown (1936-37), Heintz (1967), Harrington (1968a), and Ling (1989).

In order to apply our theory to this star we have used the elements and masses published by W.D. Heintz to initialize the calculation process.

2.1. Results

The results obtained are shown in Tables 1 and 2, in which we give the values of the orbital elements calculated 25, 50, 75 and 100 years after the initial moment $t_0 = 1900.0$. (Masses: $m_A = 0.83M_\odot$, $m_a = 0.31M_\odot$, $m_{Bb} = 0.92M_\odot$).

TABLE 1. Inner orbit

$t - t_0$	P_1	T_1	e_1	a_1	i_1	Ω_1	ω_1
0 ^y	1 ^y 832	1935.41	0.560	0 ^{''} .202	83°6	326°0	146°0
25 ^y	1 ^y 833	1936.748	0.558	0 ^{''} .202	86°27	323°86	143°05
50 ^y	1 ^y 827	1934.477	0.560	0 ^{''} .201	88°55	321°63	140°53
75 ^y	1 ^y 827	1935.799	0.555	0 ^{''} .202	91°13	319°36	138°16
100 ^y	1 ^y 827	1935.242	0.554	0 ^{''} .202	94°03	316°24	135°02

3. APPLICATION TO ζ AQUARI

This triple star was previously studied by R. S. Harrington (1968b) and W. D. Heintz (1984). This system closely resembles the previous one except that the orbital periods are much longer (about 10 times), therefore the perturbations are observed with difficulty in the approximately 200 years for which there are observations.

In this case the elements used to initialize the calculation process were those published by Heintz in 1984. The variations of the elements with secular perturbations in 10 centuries were:

$$\begin{aligned} \Delta i_1 &= 14^{\circ}3 & \Delta i_2 &= -1^{\circ}2 \\ \Delta \Omega_1 &= -10^{\circ}2 & \Delta \Omega_2 &= 0^{\circ}9 \\ \Delta \omega_1 &= -5^{\circ}0 & \Delta \omega_2 &= 0^{\circ}7 \end{aligned}$$

TABLE 2. Outer orbit

$t - t_0$	P_2	T_2	e_2	a_2	i_2	Ω_2	ω_2
0 ^y	59 ^y 84	1935.17	0.414	2 ^{''} 53	122°65	101°59	127°53
25 ^y	59 ^y 85	1935.64	0.414	2 ^{''} 53	122°35	101°54	127°90
50 ^y	59 ^y 887	1936.08	0.415	2 ^{''} 531	122°15	101°30	127°84
75 ^y	59 ^y 911	1936.54	0.414	2 ^{''} 532	121°80	101°30	127°66
100 ^y	59 ^y 912	1937.16	0.415	2 ^{''} 533	121°58	100°94	127°32

It is foreseeable that high resolution techniques, such as speckle interferometry, permit the discovery of new triple systems of the ξ Ursae Majoris type or even faster which will facilitate the comparison of the different approaches to the stellar three-body problem.

4. REFERENCES

- Abad, A.J. 1984, *Estudio de Sistemas Estelares Múltiples*. Pub. Seminario García Galdeano, Ser. II, Sec. 2, num. 2, Universidad de Zaragoza. Spain
- Brown, E.W. 1936-37, The Stellar Problem of Three Bodies, *MNRAS*, **97**, num. 7, 25
- Deprit, A. 1969, Canonical Transformations Depending on a Small Parameter, *Celes. Mech.*, **1**, 12
- Docobo, J.A. 1977, *Aplicación de la teoría de Perturbaciones al estudio de sistemas estelares triples*. Tesis Doctoral. Universidad de Zaragoza
- Docobo, J.A. & Prieto, C. 1988, The Stellar Three-body Problem. Explicit Formulation for Computing Perturbations. *Pub. Obs. Ast. R.M. Aller*, **46**
- Harrington, R.S. 1968a, Dynamical Evolution of Triple Stars, *AJ*, **73**, 190
- Harrington, R.S. 1968b, Dynamics of the Triple System Zeta Aquarii, *AJ*, **73**, 508
- Harrington, R.S. 1969, The Stellar Three-body Problem, *Celes. Mech.*, **2**, 200
- Heintz, W.D. 1967, Storungen und Bahnelemente im System ξ Ursae Majoris, *Astrom. Nach.*, **289-6**, 269
- Heintz, W.D. 1984, The Triple Star Zeta Aquarii, *ApJ*, **284**, 806
- Ling, J.F. 1989, *Aplicación del método estroboscópico al problema estelar de tres cuerpos*. Tesis Doctoral. Universidad de Santiago