



Erratum

Latent subtypes of manic and/or irritable episode symptoms in two population-based cohorts – ERRATUM

Ryan Arathimos, Chiara Fabbri, Evangelos Vassos, Katrina A. S. Davis, Oliver Pain, Alexandra Gillett, Jonathan R. I. Coleman, Ken Hanscombe, Saskia Hagenaars, Bradley Jermy, Anne Corbett, Clive Ballard, Dag Aarsland, Byron Creese and Cathryn M. Lewis

Keywords

Bipolar affective disorders; epidemiology; genetics; mood disorders; subtypes; erratum.

Copyright and usage

© The Author(s), 2022. Published by Cambridge University Press on behalf of the Royal College of Psychiatrists. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

DOI: https://doi.org/10.1192/bjp.2021.184 Published online by Cambridge University Press, 4 January 2022

This article was published with an error in Figure 1 Section b, which accidentally duplicated the percentage values of Figure 1 Section a. The correct version of the figure is below.

The publisher apologises for the error.

Ryan Arathimos; Chiara Fabbri; Evangelos Vassos; Katrina A. S. Davis; Oliver Pain; Alexandra Gillett; Jonathan R. I. Coleman; Ken Hanscombe; Saskia Hagenaars; Bradley Jermy; Anne Corbett; Clive Ballard; Dag Aarsland; Byron Creese; Cathryn M. Lewis

Reference

Arathimos R, Fabbri C, Vassos E, Davis KAS, Pain O, Gillett A, Coleman JRI, Hanscombe K, Hagenaars S, Jermy B, Corbett A, Ballard C, Aarsland D, Creese B. and Lewis CM. Latent subtypes of manic and/or irritable episode symptoms in two population-based cohorts. *Br J Psychiatry* 2022; 1–10. doi: 10.1192/bjp.2021.184.








Fig. 1. Conditional probabilities of each response (symptom) in (a) the UK Biobank optimum five-class latent model (N = 42 183) and (b) the Platform for Research Online to Investigate the Genetics and Cognition in Aging (PROTECT) replication study optimum five-class latent model (N = 4445).