

## A CLASS OF SPACES WITH WEAK NORMAL STRUCTURE

BRAILEY SIMS

It has recently been shown that a Banach space enjoys the weak fixed point property if it is  $\varepsilon_0$ -inquadrate for some  $\varepsilon_0 < 2$  and has WORTH; that is, if  $x_n \xrightarrow{w} 0$  then,  $\|x_n - x\| - \|x_n + x\| \rightarrow 0$ , for all  $x$ . We establish the stronger conclusion of weak normal structure under the substantially weaker assumption that the space has WORTH and is ' $\varepsilon_0$ -inquadrate in every direction' for some  $\varepsilon_0 < 2$ .

A Banach space  $X$  is said to have the *weak fixed point property* if whenever  $C$  is a nonempty weak compact convex subset of  $X$  and  $T : C \rightarrow C$  is a *nonexpansive* mapping; (that is,  $\|Tx - Ty\| \leq \|x - y\|$  for all  $x, y \in C$ ), then  $T$  has a fixed point in  $C$ .

It is well known that if  $X$  fails to have the weak fixed point property then it fails to have *weak normal structure*; that is,  $X$  contains a weak compact convex subset  $C$  with more than one point which is *diametral* in the sense that, for all  $x \in C$

$$\sup\{\|y - x\| : y \in C\} = \text{diam } C := \sup\{\|y - z\| : y, z \in C\}.$$

Further, if  $X$  fails to have weak normal structure then there exists a sequence,  $(x_n)$ , satisfying;

$$(S1) \quad x_n \xrightarrow{w} 0$$

and for  $C := \overline{\text{co}}\{x_n : n \in \mathbb{N}\}$

$$(S2) \quad \lim_n \|x - x_n\| = \text{diam } C = 1, \quad \text{for all } x \in C.$$

That is,  $(x_n)$  is a non-constant weak null sequence which is 'diameterising' for its closed convex hull. In particular, since  $0 \in C$ , we have  $\|x_n\| \rightarrow 1$ .

Details of these and related results may be found in the monograph by Goebel and Kirk [7] for example.

---

Received 6th September, 1993

Copyright Clearance Centre, Inc. Serial-fee code: 0004-9729/94 \$A2.00+0.00.

For the Banach space  $X$  we define  $\delta : [0, 2] \times X \setminus \{0\} \rightarrow |\mathbb{R}|$  by

$$\delta(\varepsilon, x) := \inf \left\{ 1 - \left\| y + \frac{\varepsilon}{2\|x\|}x \right\| : \|y\| \leq 1 \text{ and } \left\| y + \frac{\varepsilon}{\|x\|}x \right\| \leq 1 \right\}.$$

We refer to  $\delta(\varepsilon, x)$  as the *modulus of convexity in the direction  $x$* .  $X$  is *uniformly convex in every direction* (UCED) if  $\delta(\varepsilon, x) > 0$ , for all  $x \neq 0$  and all  $\varepsilon > 0$  (Day, James and Swaminathan [2]).

The *modulus of convexity* of  $X$  is given by

$$\delta(\varepsilon) := \inf_{x \neq 0} \delta(\varepsilon, x),$$

and  $X$  is *uniformly convex* if  $\delta(\varepsilon) > 0$ , for all  $\varepsilon > 0$ .

Following Day, given  $\varepsilon_0 \in (0, 2]$  we say  $X$  is  $\varepsilon_0$ -*inquadrate* if  $\delta(\varepsilon_0) > 0$ .

By analogue with this last definition, for  $\varepsilon_0 \in (0, 2]$  we shall say  $X$  is  $\varepsilon_0$ -*inquadrate in every direction* if  $\delta(\varepsilon_0, x) > 0$ , for all  $x \neq 0$ .

It is readily verified that  $X$  is  $\varepsilon_0$ -inquadrate in every direction if and only if whenever  $\limsup_n \|x_n\| \leq 1$ ,  $\limsup_n \|x_n + \lambda_n x\| \leq 1$ , and  $\|x_n + (\lambda_n/2)x\| \rightarrow 1$  we have  $\limsup_n |\lambda_n| \|x\| \leq \varepsilon_0$ .

Note there are also the weaker notions, of  $X$  being  $\varepsilon_0$ -inquadrate in some subset of directions, and for each  $x \neq 0$  there being an  $\varepsilon_x \in [0, 2)$  with  $\delta(\varepsilon_x, x) > 0$ , however; these will not concern us.

Garkavi [5] showed that spaces which are UCED have weak normal structure and hence enjoy the weak fixed point property. An essentially similar argument establishes the result for spaces which are  $\varepsilon_0$ -inquadrate in every direction for some  $\varepsilon_0 \in (0, 1)$ . To see this, suppose that  $X$  fails weak normal structure and so contains a sequence  $(x_n)$  satisfying (S1) and (S2). Choose  $m$  so that  $\|x_m\| > \varepsilon_0$ , then putting  $x = x_m$  we have  $\|x_n\| \rightarrow 1$ ,  $\|x_n - x\| \rightarrow 1$  and, since  $0 \in C$ , so  $x/2 \in C$ ,  $\|x_n - x/2\| \rightarrow 1$  contradicting the assumption that  $X$  is  $\varepsilon_0$ -inquadrate in every direction.

In general the situation when  $1 \leq \varepsilon_0 < 2$  remains unresolved, even in the  $\varepsilon_0$ -inquadrate case.

Two other 'classical' conditions known to be sufficient for weak normal structure are:

- (1) *The condition of Opial*, whenever  $x_n \xrightarrow{w} 0$  and  $x \neq 0$  we have

$$\limsup_n \|x_n\| < \limsup_n \|x_n - x\|.$$

The condition was introduced by Opial [10], and shown to imply weak normal structure by Gossez and Lami Dozo [8]. The condition is unchanged if both lim sups are

replaced by  $\liminf$  s. We say  $X$  satisfies the *non-strict Opial condition* if the condition holds with strict inequality replaced by ' $\leq$ '.

(2)  $\varepsilon_0$ -Uniform Radon-Reisz ( $\varepsilon_0$ -URR), for some  $\varepsilon_0 \in (0, 1)$ ; there exist  $\delta > 0$  so that whenever  $x_n \xrightarrow{w} x$ , with  $\|x_n\| \leq 1$  and  $sep(x_n) := \inf\{\|x_n - x_m\| : n \neq m\} > \varepsilon_0$  we have  $\|x\| < 1 - \delta$ . When the condition holds for all  $\varepsilon_0 > 0$  we say  $X$  is URR. The condition is essentially due to Huff [9], and was shown to imply weak normal structure by van Dulst and Sims [4].

Gossez and Lami Dozo [8] showed that Opial's condition follows from the non-strict version in the presence of uniform convexity, however a careful reading of their argument establishes the following.

**PROPOSITION 1.** *If  $X$  is UCED and satisfies the non-strict Opial condition then  $X$  satisfies the Opial condition.*

PROOF: Suppose  $X$  fails the Opial condition, then there exists a sequence  $x_n \xrightarrow{w} 0$  and  $x \neq 0$  with

$$\liminf_n \|x_n\| \not\leq \liminf_n \|x_n - x\|.$$

By the non-strict Opial condition we must have equality, and without loss of generality we may assume that  $\|x_n\| \rightarrow 1$ . Let  $y_n := x_n - x$ , then  $\|x_n\|, \|y_n\| \rightarrow 1$  and  $x_n - y_n = x$ . Thus, by UCED we must have that

$$\liminf_n \|x_n - x/2\| = \liminf_n \|(x_n + y_n)/2\| < 1 = \liminf_n \|x_n\|,$$

contradicting the non-strict Opial condition. □

In Sims [12], the notion of *weak orthogonality* (WORTH); if  $x_n \xrightarrow{w} 0$  then for all  $x \in X$  we have

$$\|x_n - x\| - \|x_n + x\| \rightarrow 0,$$

was introduced (also see Rosenthal [11]), and it was asked whether spaces with WORTH have the weak fixed point property. WORTH generalises the lattice theoretic notion of 'weak orthogonality' introduced by Borwein and Sims [1] and shown to be sufficient for the weak fixed point property in Sims [12].

**PROPOSITION 2.** *The non-strict Opial condition is entailed by WORTH.*

PROOF: If  $x_n \xrightarrow{w} 0$  then for any  $x \in X$  we have

$$\begin{aligned} \limsup_n \|x_n\| &\leq \frac{1}{2} (\limsup_n \|x_n - x\| + \limsup_n \|x_n + x\|) \\ &= \limsup_n \|x_n - x\|, \quad \text{as } \lim_n \|x_n - x\| - \|x_n + x\| = 0, \text{ by WORTH.} \end{aligned}$$

□

Combining this with proposition 1 we obtain the following.

**COROLLARY 3.** *A Banach space which has UCED and WORTH satisfies the Opial condition.*

Recently García Falset [6] working through the intermediate notion of the ACM-property has shown that spaces which are  $\varepsilon_0$ -inquadrate for some  $\varepsilon_0 < 2$  and have WORTH have the weak fixed point property.

We give a direct and elementary proof that the stronger conclusion of weak normal structure follows from the substantially weaker premises of WORTH and  $\varepsilon_0$ -inquadrate in every direction for some  $\varepsilon_0 < 2$ . That  $\varepsilon_0$ -inquadrate in every direction is genuinely weaker than  $\varepsilon_0$ -inquadrate follows since spaces which are  $\varepsilon_0$ -inquadrate, for an  $\varepsilon_0 < 2$  are necessarily superreflexive (van Dulst [3]) while every separable Banach space has an equivalent norm which is UCED [2], and hence  $\varepsilon_0$ -inquadrate in every direction for  $0 < \varepsilon_0 < 2$ .

**DEFINITION:** We say a Banach space  $X$  has *property (k)* if there exists  $k \in [0, 1)$  such that whenever  $x_n \xrightarrow{w} 0$ ,  $\|x_n\| \rightarrow 1$  and  $\liminf_n \|x_n - x\| \leq 1$  we have  $\|x\| \leq k$ . Note: By considering subsequences we see that the property remains unaltered if in the definition we replace  $\liminf$  by  $\limsup$ .

Property (k) is an interesting condition which clearly exposes the uniformity in Opial's condition. Indeed Opial's condition corresponds to property (k) with  $k = 0$ .

**PROPOSITION 4.** *If  $X$  has property (k) then  $X$  has weak normal structure.*

**PROOF:** Suppose  $X$  fails weak normal structure, then there is a sequence  $(x_n)$  satisfying (S1) and (S2). Choosing  $m$  sufficiently large so that  $\|x_m\| > k$  (see the remark following S2) and taking  $x := x_m$  we have that property (k) is contradicted by the sequence  $(x_n)$ .  $\square$

We now turn to conditions sufficient for property (k), and hence also for weak normal structure.

**PROPOSITION 5.** *If  $X$  is  $\varepsilon_0$ -URR, for some  $\varepsilon_0 \in (0, 1)$  then  $X$  has property (k).*

**PROOF:** Suppose  $x_n \xrightarrow{w} 0$ ,  $\|x_n\| \rightarrow 1$  and  $\limsup_n \|x_n - x\| \leq 1$ . Choose  $m$  so that  $\|x_m\| > \varepsilon_0$ , then, since  $\liminf_n \|x_n - x_m\| \geq \|x_m\|$ , we may extract a subsequence, which we continue to denote by  $(x_n)$ , with  $\|x_n - x_m\| > \varepsilon_0$  for all  $n$ . Continuing in this way we obtain a subsequence, still denoted by  $(x_n)$ , with  $\text{sep}(x_n) > \varepsilon_0$ . But, then  $x_n - x \xrightarrow{w} x$  is a sequence in the unit ball with a separation constant greater than  $\varepsilon_0$  and so  $\|x\| \leq 1 - \delta$ , where  $\delta$  is given by the definition of  $\varepsilon_0$ -URR. Thus  $X$  has property (k) with  $k = 1 - \delta$ .  $\square$

**PROPOSITION 6.** *If  $X$  is  $\varepsilon_0$ -inquadrate in every direction for some  $\varepsilon_0 \in (0, 1)$  and satisfies the non-strict Opial condition then  $X$  has property (k).*

PROOF: Suppose  $x_n \xrightarrow{w} 0$ ,  $\|x_n\| \rightarrow 1$  and  $\limsup_n \|x_n - x\| \leq 1$ . If  $x = 0$  there is nothing to prove, so we assume that  $x \neq 0$ . Then  $x_n$  and  $y_n := x_n - x$  are two sequences in the unit ball with  $x_n - y_n = x$  a fixed direction, so

$$\left\| \frac{x_n + y_n}{2} \right\| \leq 1 - \delta(\|x\|, x) < 1, \text{ if } \delta(\|x\|, x) > 0.$$

But, then  $\limsup_n \|x_n - x/2\| < 1 = \limsup_n \|x_n\|$ , contradicting the non-strict Opial condition. Thus we must have  $\delta(\|x\|, x) = 0$  and this requires  $\|x\| \leq \varepsilon_0$ . So  $X$  has property (k) with  $k = \varepsilon_0$ . □

The case when  $X$  is  $\varepsilon_0$ -inquadrate in every direction for an  $\varepsilon_0 \in [1, 2)$  is handled by the following proposition which in conjunction with proposition 3 yields our main result.

**PROPOSITION 7.** *If  $X$  is  $\varepsilon_0$ -inquadrate in every direction for some  $\varepsilon_0 \in (0, 2)$  and has WORTH then  $X$  has property (k).*

PROOF: Suppose  $x_n \xrightarrow{w} 0$ ,  $\|x_n\| \rightarrow 1$  and  $\limsup_n \|x_n - x\| \leq 1$ . Let  $a_n := x_n - x$  and  $b_n := x_n + x$ . Then by WORTH  $\|a_n\| - \|b_n\| \rightarrow 0$ , so  $\limsup_n \|b_n\| = \limsup_n \|a_n\| = \limsup_n \|x_n - x\| \leq 1$ , and  $b_n - a_n = 2x$ . Therefore we have

$$\limsup_n \|x_n\| = \limsup_n \|(a_n + b_n)/2\| \leq 1 - \delta(2\|x\|, x) < 1,$$

a contradiction, unless  $2\|x\| \leq \varepsilon_0$ . Thus  $X$  has property (k) with  $k = \varepsilon_0/2$ . □

If  $X$  is  $\varepsilon_0$ -inquadrate then the calculations of the previous proof allow some flexibility. If we measure the ‘degree of WORTHwhileness’ of a Banach space  $X$  by

$$w := \sup\{\lambda : \lambda \liminf_n \|x_n + x\| \leq \liminf_n \|x_n - x\|, \text{ whenever } x_n \xrightarrow{w} 0 \text{ and } x \in X\},$$

(so  $X$  has WORTH if and only if  $w = 1$ ) then we can adapt the above calculations to verify the following.

**PROPOSITION 8.**  *$X$  has property (k) if*

$$w > \max\{\varepsilon_0/2, 1 - \delta(\varepsilon_0)\}$$

for some positive  $\varepsilon_0$ .

We close by noting that many spaces have WORTH, including all Banach lattices which are ‘weakly orthogonal’ in the sense introduced by Borwein and Sims [1]. In particular for  $0 < \alpha < 1$  the space  $\ell_2$  with the equivalent norm  $\|x\| := \max\{\alpha \|x\|_2, \|x\|_\infty\}$

has WORTH and hence satisfies the non-strict Opial condition, but fails to satisfy the Opial condition for any  $\alpha$ .

However, many important spaces do not enjoy WORTH, for example with the exception of  $p = 2$  all the spaces  $\mathcal{L}_p[0, 1]$  fail to satisfy the non-strict Opial condition (see the details of the example given in Opial [10]) and hence also fail to have WORTH. They do however enjoy property (k); for example, when  $p > 2$  it follows from Clarkson's inequality that we may take  $k = (1 - 2^{-p})^{1/p}$ .

#### REFERENCES

- [1] J.M. Borwein and B. Sims, 'Non-expansive mappings on Banach lattices and related topics', *Houston J. Math* **10** (1984), 339–355.
- [2] M.M. Day, R.C. James and S. Swaminathan, 'Normed linear spaces that are uniformly convex in every direction', *Canad. J. Math.* **23** (1971), 1051–1059.
- [3] D. van Dulst, *Reflexive and superreflexive Banach spaces*, M.C. Tract **102** (Mathematical Centre, Amsterdam, 1978).
- [4] D. van Dulst and B. Sims, 'Fixed points of nonexpansive mappings and Chebyshev centres in Banach spaces with norms of type KK', in *Banach space theory and its applications*, Lecture Notes in Maths. **991** (Springer Verlag, Berlin, Heidelberg, New York, 1983), pp. 35–43.
- [5] A.L. Garkavi, 'The best possible net and the best possible cross-section of a set in a normed linear space', *Izv. Akad. Nauk. SSSR Ser. Mat* **26** (1962), 87–106. *Amer. Math. Soc. Trans. Ser.* **39** (1964), 111–131.
- [6] J. García Falset, 'The fixed point property in Banach spaces whose characteristic of uniform convexity is less than 2', *J. Austral. Math. Soc.* **54** (1993), 169–173.
- [7] K. Goebel and W.A. Kirk, *Topics in metric fixed point theory*, Cambridge Studies in Advanced Mathematics **28** (Cambridge University Press, New York, 1990).
- [8] J.-P. Gossez and E. Lami Dozo, 'Some geometric properties related to the fixed point theory for nonexpansive mappings', *Pacific J. Math.* **3** (1972), 565–573.
- [9] R. Huff, 'Banach spaces which are nearly uniformly convex', *Rocky Mountain J. Math.* **10** (1980), 743–749.
- [10] Z.O. Opial, 'Weak convergence of the sequence of successive approximations for non-expansive mappings', *Bull. Amer. Math. Soc* **73** (1967), 591–597.
- [11] H.P. Rosenthal, 'Some remarks concerning unconditional basic sequences', Longhorn Note – Uni of Texas, Texas Functinal Analysis Seminar (1982–3), pp. 15–47.
- [12] B. Sims, 'Orthogonality and fixed points of nonexpansive maps', *Proc. Centre Math. Anal. Austral. Nat. Univ.* **20** (1988), 178–186.

Department of Mathematics  
The University of Newcastle  
Newcastle NSW 2308  
Australia