

SEROTONINERGIC ANTIDEPRESSANTS AND ANTICOAGULANTS IN PSYCHOGERIATRIC WARDS, MENTAL HOSPITAL KROMERIZ, CZECH REPUBLIC - WHAT SHALL WE DO TO PREVENT FROM SERIOUS DRUG INTERACTIONS?

J. Luzny, M. Blazkova, F. Honzak, V. Bican, O. Vyrostkova, A. Indrakova, I. Pospisilova

Mental Hospital Kromeriz, Kromeriz, Czech Republic

Introduction: Both serotonergic antidepressants and anticoagulants are widely used in psychogeriatrics. These drugs can develop drug interactions with subsequent risk of bleeding.

Objectives, aims: To determine risk of potential drug interactions between serotonergic antidepressants and anticoagulants serotonergic antidepressants and anticoagulants with subsequent risk of bleeding.

Methods: Retrospective analysis of medical records and laboratory checks (INR values) in senior patients treated with serotonergic antidepressants and anticoagulants (Mental hospital Kromeriz, Czech republic, year 2011).

Results: In 15.5% of patients treated with combination serotonergic antidepressant / anticoagulant the dosage of anticoagulant had to be decreased due to unwanted elevation of INR value. In 0.6% of patients the risk of bleeding was clinically evident.

Conclusion: Serotonergic antidepressants and anticoagulants are common drugs used in treating seniors with mental disorder and potential drug interactions between them should be taken into account.