

Observable Impacts of Exoplanets on Stellar Hosts – An X-Ray Perspective

Scott J. Wolk¹, Ignazio Pillitteri^{1,2} and Katja Poppenhaeger^{1,3}

¹Harvard Smithsonian Center for Astrophysics, 60 Garden Street,
Cambridge MA 02138, USA
email: swolk@cfa.harvard.edu

²Osservatorio Astronomico di Palermo, piazza del Parlamento 1, I-90134 Palermo, Italy
email: pilli@astropa.unipa.it

³Queen's University Belfast, University Road, Belfast, BT7 1NN, Northern Ireland, UK
email: K.Poppenhaeger@qub.ac.uk

Abstract. Soon after the discovery of hot Jupiters, it was suspected that interaction of these massive bodies with their host stars could give rise to observable signals. We discuss the observational evidence for star-planet interactions (SPI) of tidal and magnetic origin observed in X-rays. Hot Jupiters can significantly impact the activity of their host stars through tidal and magnetic interaction, leading to either increased or decreased stellar activity – depending on the internal structure of the host star and the properties of the hosted planet. We provide several examples of these interactions. In HD 189733, the strongest X-ray flares are preferentially seen in a very restricted range of planetary phases. Hot Jupiters, can also obscure the X-ray signal during planetary transits. Observations of this phenomena have led to the discovery of a thin upper atmospheres in HD 189733A. On the other hand, WASP-18 – an F6 star with a massive hot Jupiter, shows no signs of activity in X-rays or UV. Several age indicators (isochrone fitting, Li abundance) point to a young age ($\sim 0.5 - 1.0$ Gyr) and thus significant activity was expected. In this system, tidal SPI between the star and the very close-in and massive planet appears to disrupt the surface shear layer and thus nullify the stellar activity.

Keywords. X-rays: stars, magnetic fields, exoplanets, interactions

1. Introduction

Over the two decades, the discovery of exoplanets has fundamentally changed our perception of the universe and humanity's place within it. The role of X-rays in the study of exoplanets is subtle, but recent work indicates exoplanets, especially hot Jupiter systems, are unique X-ray environments and the impact of X-rays may be significant for the evolution of the system. The effects can work several ways; the intense high energy flux alters the thermal budget of the upper atmosphere of planet, the angular momentum and magnetic field of the planet can induce more activity on the star and the enhanced X-rays are absorbed by the transiting planet, which, in turn, act as a probe of the planetary upper atmosphere. In addition, an overall enhancement of the stellar host activity can significantly influence the chemistry of any additional planet in the habitable zone of the same star and thus the evolution of life in the system.

X-rays play a significant role in the evolution of close-in exoplanets. X-rays can modify the chemistry of exoplanet upper atmospheres and which can become over-inflated and evaporate because of the strong UV/X-ray flux to which they are exposed (see also presentations in these proceedings by Airapetian and Vidotto). As first noted by Lammer *et al.* (2003), inclusion of stellar X-ray and EUV flux in irradiance calculations leads to energy-limited escape and atmospheric expansion not found in models incorporating stellar UV/optical/IR insulation alone. The increased mass loss rates are of order 10^{12} g s⁻¹,

implying hydrogen-rich exoplanets may evaporate and shrink to levels at which heavier atmospheric constituents may prevent hydrodynamic escape. The generation of an exosphere due to local X-ray luminosity has been directly detected in the case of the planets HD 209458b and HD 189733b. Absorption by atmospheric gas has been used to probe the layer where the gas escapes in the upper atmosphere (e.g. Vidal-Madjar 2003, Ballester *et al.* 2007 and Poppenhaeger *et al.* 2013).

Modeling by Penz *et al.* (2008) shows evolution of close-in exoplanets strongly depends on the detailed X-ray luminosity history of their host stars. Stars located at the high end of the X-ray luminosity distribution evaporate most of their planets' atmospheres within 0.05 AU, while a significant fraction of planets can survive if exposed to a moderate X-ray luminosity. At lower X-ray luminosities, they find that the mass loss is negligible for hydrogen-rich Jupiter-mass planets at orbits >0.02 AU, while Neptune-mass planets are influenced up to 0.05 AU (see also Murray-Clay *et al.* 2009).

Analytic models and MHD simulations show that the X-ray environment can be modified by Star-Planet Interaction (SPI) when Jupiter mass planets are very close to their parentstars (< 0.1 AU; c.f. Lanza 2008, 2009, Cohen *et al.* 2009) SPI arises through two mechanisms, either gravitational tides, or interactions between magnetic fields. Both processes should increase in strength as a^{-3} (where a is the separation distance; Cuntz *et al.* 2000).

SPI not only proceeds from star to the planet, but theoretical arguments demonstrate that hot Jupiters (HJs; planets with masses approximately equal to or greater than Jupiter with an orbital semi-major axis of less than 0.1 AU) can also influence their hosts. Cuntz *et al.* (2000) showed that energy generation due to tidal perturbations is also proportional to a^{-3} . More detailed work, demonstrates HJs induce tidal bulges on the host star. Cool stars can dissipate the energy contained in the bulges much more effectively than hot stars due to turbulent eddies in the convective envelopes (Zahn 2008). Similarly, Saar *et al.* (2004) estimated The energy released via reconnection during an interaction of the planetary magnetosphere with the stellar magnetic field is estimated as $F_{int} \propto B_* \times B_p v_{rel} a_p^n$ (with $n \sim -3$).

The manifestation of SPI in X-ray band is matter of debate. Kashyap *et al.* (2008) show that stars with hot Jupiters are statistically brighter in X-rays than stars without hot Jupiters. On the other hand, Poppenhaeger *et al.* (2010) have found no statistical evidence of X-ray SPI. Sample selection seems to be at the heart of the issue. Local solar analogs selected for inactivity (perhaps introducing a fundamental bias in the method) showed no significant correlation between common proxies for interaction strength (M_p/a^2 or M_p/a) versus coronal activity (L_x or L_x/L_{bol}), but a full sample of ~ 200 FGK stars does (Miller *et al.* 2015). Miller *et al.* argue that there is a threshold for SPI and the effect is driven by hot Jupiters ($M_p/a^2 > 450 M_{Jup}/AU^2$). In both analysis, a bias was noted in the data that stars hosting hot Jupiters were more active, but in both cases the effect was related to a limited number of extreme outliers in the sample, not a general trend.

The focus of this presentation is to discuss signs of feedback between the stars and planets in some of these extreme examples. In some cases there is a detectable enhancement, other times the high energy signal appear nullified. These cases may prove crucial for evolution of planets as well as our estimation of their habitability. For our purposes, star-planet interaction (SPI) is driven by magnetic interaction between the stellar and planetary magnetic fields, or by tidal interaction (Cuntz *et al.*, 2000). Both effects strongly depend on the planet-star separation, which is directly measurable. But the observation result is also a function of the intensity and topology of the magnetic fields, and the internal structure of the star – which are less apparent.

2. Observations

For the purposes of this contribution we will focus on observations of three systems with HJs. The second and third cases are edge cases representing a highly eccentric system and a high mass planet respectively. The first case, on the other hand, is the gold standard of HJs.

2.1. *HD 189733*

HD 189733 is one of the best studied systems with a transiting HJ. It is in a binary system with a quite inactive and old ($\tau > 5$ Gyr) M4 secondary star. This system was originally thought to be about 600 Myr, based on a relatively high activity level of the primary which led to the assumption it was a Hyades member. Pillitteri *et al.* (2010) noted they did not detect the M4 companion in the XMM image although it should have been bright enough if it were indeed 600 Myr. They speculated that the activity in the primary was enhanced by the interaction with the exoplanet. Indeed, when Poppenhaeager, Schmitt & Wolk (2013) detected the secondary star with the *Chandra X-ray Observatory*, they found an activity level consistent with an age of about 5 Gyr. This latter finding has led to the conclusion that the primary has been spun up by the hot Jupiter (Poppenhaeager & Wolk 2014).

2.1.1. *In Eclipse*

In addition to the age anomaly, Pillitteri *et al.* (2014) discuss three eclipse observations of HD 189733b (Fig. 3). Each time they noted a significant flare within hours after the eclipse. They speculated this was due to a hot spot forward phased from the sub planetary point by about 90° , consistent with analytic predictions by Lanza (2008). Using HST, Pillitteri *et al.* (2015) acquired high quality COS spectra in the wavelength range 1150–1450 Å. Again, flares were observed just after the eclipse. They found two episodes of strong variability of the line fluxes of ions of Si, C, and N that had not been observed in planetary transits. The details of the flares were consistent with an MHD model (Matsakos *et al.* 2015). The flow morphology in the model provides a natural explanation of the FUV line and X-ray variability of HD 189733. Specifically, the plasma is liberated from the upper atmosphere of the planet and funneled by the magnetized stellar wind in an almost radial trajectory close to the star. The flow forms a knee structure that consists of hot and dense plasma which then accretes in a region of the star fixed with the synodic phase (Fig. 2).

On the other hand, the X-ray flare observed in 2012, had aspects of reverberation. During the decay of the flare, three successively smaller peaks are observed separated by about 4 ks. This appears to have been a damped magneto acoustic oscillation in a flaring loop (e.g. Mitra-Kraev *et al.* 2005). In such loop the change of the intensity of the successive peaks can be described as:

$$\frac{\Delta I}{I} \sim \frac{4\pi n k_B T}{B^2}$$

Since fits to the XMM spectra can be used to determine the temperature and the density, this formulation can be used to measure the magnetic field – if magneto acoustic oscillation is the cause of the flare structure. In this case, the derived B field (~ 40 G) is consistent with results of spectropolarimetry (Fares *et al.* 2010). With the reverberation hypothesis thus supported, the length of the loop can be calculated by simple arguments about the sound speed, a function of the density, and the travel time (the time between the successive peaks). The result is a loop length of about $4 R_*$ indicating a flare covering half the distance to the planet. While there is no evidence that the flare is actually




Figure 1. Light curves for the XMM-Newton EPIC/PN observations of three eclipses of HD 189733b. The top X-axis indicates phase. The bottom X-axis indicates time (normalized to the phase start of the 2011 observation). First through fourth contacts are indicated by the vertical lines. In all cases, the quiescent count rates are within 20% of 100 cnt/ks.

directed to the planet, the result is reminiscent of Favata *et al.* (2005) and McCleary & Wolk (2011). Both groups found that for flares in PMS stars, long length flares only occurred in cases where the star is surrounded by a disk. Both groups concluded that the flares stretched from the star to the disk in these cases. By analogy the magnetic field of the planet could be acting as a footpoint for occasional, massive, flares. Such flares would have significant impacts on the planet, ionizing material in its upper atmosphere. Since it appears to be tidal interaction between the star and the planet is the *prima facie* cause of the enhanced stellar field, planetary orbital energy is the ultimate source of the flare that scorches the planet's atmosphere.

2.1.2. *In Transit*

Transit observations of HD 189733b across HD 189733A have been performed with both XMM and *Chandra*. Popenhaeger *et al.* (2013) reported a detection of the planetary transit in soft X-rays using several coadded observations mostly relying on five clean (flare-free) *Chandra* observations. They noted a significantly deeper X-ray transit depth than observed in the optical. The X-ray data favor a transit depth of 6% – 8%, versus a broadband optical transit depth of 2.41%. Because several observation were co-added they were able to exclude transits of active regions and other possible stellar origins for this deep transit. They interpret the deep X-ray transit to be caused by a thin




Figure 2. Particle density contours of an MHD simulation that models star–planet interactions between a HJ and its host (polar view). The star rotates counterclockwise, and the planet orbits the star along the same direction. The two “+” symbols shown on the left panel indicate the location of the star (red disk) and the planet (green disk). Right panel: a close-up of the impact region, where the motion of the accreting plasma is marked with arrows. Specifically, the shocked plasma is funneled by the magnetized stellar wind in an almost radial trajectory close to the star (A), forms a “knee” structure that consists of hot and dense plasma (B), and then accretes in a spot ahead of the orbital phase (C). The precise details of accretion are not investigated by the simulation (zone marked with ?). The knee (B) of the stream and the active spot upon impact on the surface (C) are the main sites of production of the enhanced flux observed in the FUV and X-ray bands and phased with the orbital motion (from Pillitteri *et al.* 2015).

outer planetary atmosphere extended to about 1.75 times the nominal radius which is transparent at optical wavelengths, but dense enough to be opaque to X-rays. The X-ray radius appears to be larger than the radius observed at far-UV wavelengths, most likely due to high temperatures in the outer atmosphere at which hydrogen is mostly ionized.

2.2. HD 17156

In addition to HJs close to their host stars, eccentric systems – in which the Jupiter is “hot” only a fraction of the time – represent another type of extreme system. These *should* be a good test bed for the existence of SPI. The prediction is that SPI impacts only occur when the star and planet are close. One candidate for such a test is HD 17156.

HD 17156 is a G0 star with a HJ in a 21 day orbit (Barbieri *et al.* 2007). The eccentricity of the orbit is 0.68. The planet reaches a minimum separation of $\sim 15 R_*$. XMM-Newton observed HD 17156 when the planet was at the periastron and a second time, when the planet was more distant. HD 17156 was not detected by XMM when the two were separated. However, just after periastron passage, there was a marked rise in the X-ray luminosity with a corresponding rise in the chromospheric activity. Maggio *et al.* (2015) suggest that this could have been either due a magnetic reconnection and or flaring activity when the planet was at its minimum separation or due to material stripped from the planet and falling onto the star. The excess of X-rays has a soft spectrum and could favor the tidal stripping hypothesis.

2.3. WASP-18

In a sense, WASP-18 (F6V) is the most extreme exoplanet system. It hosts one of the fastest orbiting and most massive HJs. The planet has a mass of over $10M_{Jup}$ and the




Figure 3. The co-added lightcurve of 5 Chandra X-ray Observatory observations (without flares) of a transit of HD 189733A by HD 189733b. The X-axis indicates phase. The orange line indicates typical optical data and an eclipse depth of 2-3 %. The dots with the error bars are the co-added and normalized X-ray data. The purple line is a best fit to the X-ray data using a limb-brightened model appropriate of an optically thin stellar corona. The best fit X-ray eclipse depth is about 7%.




Figure 4. X-ray images of HD 17156 taken far from the planetary periastron (left panel) and near the periastron (right panel). The image is colored by photon energy (red = 0.3-1.0 keV, green = 1.0-2.5 keV, blue=2.5-5.0 keV). Smoothing is applied to the images, with a Gaussian $\sigma = 4''$ kernel. Positions of the only two objects in the SIMBAD catalog are shown with small squares. Circle sizes indicate the wavelet detection scales of HD 17156 and of an unrelated background object with a harder spectrum (From Maggio *et al.* 2015).

period is about 22 hours. The estimate of the age for this system is about 600 Myrs. This is based on both isochrone fitting and the strong Li absorption. Indeed, the observed Li strength is a near perfect match for Hyades and Beehive cluster stars of similar temperature. The expected X-ray luminosity from such a star should be at least $\log L_X \sim 28.5$. This is nearly the same as observed from HD 189733 and *a priori* might indicate strong magnetic re-connection events.

In fact, exactly the opposite is seen. No X-rays were detected with a flux limit of $\log L_X \sim 26.5$. This result is part of a trend. Recently, Staab *et al.* (2016) showed that a quarter of known short period planet hosts exhibit anomalously *low* activity levels. Fossati *et al.* (2014) asserted that the lack of chromospheric activity detected from WASP-18

(and WASP-12) may have been due to local absorption of UV by atmospheric material stripped off of the planet by tidal forces. We find this unlikely for several reasons: First, given the high mass of WASP-18b, atmospheric stripping should be 100 times less effective in this case than HD 189733. Second, the calcium absorption line is observed. It is only the emission reversal which is not observed. As this should occur above the Ca absorption feature it is hard to understand how you can observe the lower lying feature while absorbing the higher altitude feature. Third, no X-rays are detected at all, not even high energy X-rays which might be expected from the Planck tail of a 1 keV thermal distribution. Pillitteri *et al.* (2014) conclude the the star is X-ray dark and suggest tidal interaction with the planet must have a role in destroying the dynamo efficiency and the overall activity of the star. Based on the formulae given in Cuntz *et al.* (2000), tides on WASP-18 are the largest of any known exoplanet host, of order of 500 km because of planet proximity and its mass. However, a complete break down in convection is not possible as this is key to energy transport in the star. On the other hand, the tidal wave on the surface may reach heights of ~ 500 km every ~ 10 hours. This could disrupt the shear layer at the top of the convection zone. This would prevent the build-up and concentration of magnetic energy close to the surface. Meanwhile the convective thermal transport from the core to near the surface is free to occur. Near the surface, radiative cooling processes dominate.

3. Conclusions

Over the past five years there has been a great deal of case specific evidence gathered that stars can interact with their HJ companions via tides and magnetic fields. The level of confidence in each kind of evidence varies.

One example that has not been touched on here comes from binary stars. Poppenhaeger & Wolk (2014) discussed using binaries to test for activity induced by HJs. They presented at least 3 examples of binaries in which the HJ host had an activity level incompatible with the age indicated by the activity level of the non-hot Jupiter host. They concluded that **very close in HJs spin-up stars when the HJ host has a large convective zone**. All the observed data are consistent with this hypothesis and it is strongest direct evidence of SPI. The corollary to this is an uncertainty when using the activity–age relation to date stars with close in planets.

Prior to eclipse observations, analytical calculations predicted the existence of an active region on HD 189733 forward phased by about 90° from the sub-planetary point. This would imply enhanced activity between phases 0.5 and 1.0. The propensity for the star to flare between phase 0.55 and 0.65 has been taken as evidence of the active region, perhaps being observed over the limb. In total there have been 5 high energy flares observed in four observations (including the FUV observations). The significance of this is unclear and fraught with over-interpretation. So while the result is suspicious, it is not yet compelling.

This could be an observational bias, certainly some (albiet smaller) flares have been noted in other phases and to be fair any flare observed between phase 0.45 and 0.6 would probably have been counted by Pillitteri *et al.* as evidence of this effect. Still, the odds of all the strong flares being observed to happen on one half of the star and not the other half is about 10% and this is taken as support of the hot spot hypothesis.

The two remaining pieces of evidence, the long length flare in HD 189733 and enhanced activity and periapses for HD 17156 are tantalizing. On the other hand, each of these has only been seen to occur once. Until additional flares are seen in HD 17156 or another eccentric system in periapses passage this will stand as a one-time occurrence.

The expected frequency for such long length flares is unknown, but it appears long. In 850 ks observing the ONC only 25 such flares were observed among about 400 disked stars (Favata *et al.* 2005). The situation is similar for the long flare seen on HD 189733. Overall the System has been monitored for several hundred kiloseconds in X-rays. While this is a long time, we expect such flares to be exceptionally rare. Simply based on the COUP, Favata *et al.* (2005) reported 25 loops longer than 10^{11} cm. In that project there was a total of 850 ks observing time put in on about 400 stars known to possess disks. From this, the derived “long flare” rate is about 1 per 13 megaseconds or about 2 per year among active young stars with disks. The occurrence rate is probably somewhat higher than this. The Favata study focused on luminous flares and the one observed on HD 189733 would not have met this criterion. Still, it appears such long flares are not the most common.

While the result of the extreme example appear compelling, statistical evidence is a different story. Several groups (Kayshap *et al.* 2008, Poppenhaeger *et al.* 2011 and Miller *et al.* 2015) have looked for statistical evidence of SPI, with mixed results. The problem is that many parameters effect the eventual outcome of the star-planet interaction. Mass and distance ratios are the obvious parameters, but depth of convection and relative field orientation and strength are clearly others. The cautionary tale of WASP-18 (and others such as WASP-12) indicates that tidal forces can be constructive or destructive when it comes to modifying stellar activity. Any statistical test needs to account properly for outliers. Both the high and low outliers may be the result of SPI while the median and mean may not be very affected. The coming few years should prove very interesting as we continue to gather evidence.

References

- Barbieri, M., *et al.* 2007, *A&A*, 476, L13
Favata, F., Flaccomio, E., Reale, F., *et al.* 2005, *ApJS*, 160, 469
Fares, R., *et al.* 2010, *MNRAS*, 406, 409
Fossati, L., Ayres, T. R., Haswell, *et al.* 2014, *APSS*, 354, 21
Kashyap, V. L., Drake, J. J., & Saar, S. H. 2008, *ApJ*, 687, 1339
Lanza, A. F. 2008, *A&A*, 487, 1163
Maggio, A., *et al.* 2015, *ApJL*, 811, L2
McCleary, J. E. & Wolk, S. J. 2011, *AJ*, 141, 201
Miller, B. P., Gallo, E., Wright, J. T., & Pearson, E. G. 2015, *ApJ*, 799, 163
Mitra-Kraev, U., Harra, L. K., Williams, D. R., & Kraev, E. 2005, *A&A*, 436, 1041
Murray-Clay, R. A., Chiang, E. I., & Murray, N. 2009, *ApJ*, 693, 23
Pillitteri, I., Wolk, S. J., Cohen, O., Kashyap, V., *et al.* 2010, *ApJ*, 722, 1216
Pillitteri, I., Wolk, S. J., Lopez-Santiago, J., Günther, H. M., *et al.* 2014, *ApJ*, 785, 145
Pillitteri, I., May A., Micela, G., Sciortino, S., Wolk, S. J., & Matsakos, T. 2015 *ApJ*, 805, 52
Poppenhaeger, K. & Wolk, S. J. 2014, *A&A*, 565, L1
Poppenhaeger, K., Schmitt, J. H. M. M., & Wolk, S. J. 2013, *ApJ*, 773, 62
Poppenhaeger, K. & Schmitt, J. H. M. M. 2011, *ApJ*, 735, 59
Staab, D., Haswell, C. A., Smith, G. D., Fossati, L., Barnes, J. R., Busuttil, R., & Jenkins, J. S. 2016, arXiv:1612.01739
Vidal-Madjar, A., Lecavelier des Etangs, A., Désert, *et al.* 2003, *Nature*, 422, 143

Discussion