

## A NOTE ON SPECIAL INVOLUTIONS

W. D. MUNN

(Received 19 December 1997; revised 7 June 1998)

Communicated by D. Easdown

### Abstract

The algebra consisting of those linear transformations of a complex inner product space that have a formal adjoint is shown to possess a special involution. Two earlier results concerning special involutions are then generalized.

1991 *Mathematics subject classification* (*Amer. Math. Soc.*): 16S50, 16W10.

For a given complex inner product space  $V$ , the set of all linear transformations of  $V$  that have a formal adjoint constitutes a complex algebra and is shown to possess an involution that is special in the sense of Easdown and Munn (Theorem 1). It follows that the standard involution on a  $C^*$ -algebra is special (Corollary 1) – a result first noted by Hofmann – and that hermitian conjugation is a special involution on the algebra of all  $I \times I$  row-finite and column-finite complex matrices, where  $I$  is an arbitrary nonempty set (Corollary 2). By combining Corollary 1 with a result of Barnes, it is proved that the natural involution on the  $l^1$ -algebra of an inverse semigroup is special (Theorem 2).

Let  $*$  be an involution on a semigroup  $S$  (that is, a permutation of  $S$  such that, for all  $a, b$  in  $S$ ,  $(ab)^* = b^*a^*$  and  $a^{**} = a$ ). We say that  $*$  is *special* if and only if, for each nonempty finite subset  $T$  of  $S$ ,

$$(\exists t \in T)(\forall u, v \in T) \quad t^*t = u^*v \Rightarrow u = v.$$

This definition is clearly equivalent to the one given originally in [2]. An involution [a special involution] on a complex algebra  $R$  is a mapping  $*$  :  $R \rightarrow R$  that is an automorphism of  $(R, +)$  and an involution [a special involution] on  $(R, \cdot)$ , with the further property that, for all  $a \in R$  and all  $\lambda \in \mathbb{C}$  (the complex field),  $(\lambda a)^* = \bar{\lambda}a^*$ ,

where  $\bar{\lambda}$  denotes the complex conjugate of  $\lambda$ . Examples of special involutions include hermitian conjugation on the algebra of all  $n \times n$  complex matrices [2] and the natural involution on the complex semigroup algebra of an inverse semigroup [3]. Both of these are generalized here. By a *star subalgebra* of a complex algebra  $R$  with an involution  $*$  we mean a subalgebra  $S$  of  $R$  such that  $a^* \in S$  for all  $a \in S$ . Observe that if  $*$  is special and  $S$  is a star subalgebra of  $R$  then  $*$  induces a special involution on  $S$ .

In the theorem below we examine a certain subalgebra of the algebra of all linear transformations of a complex inner product space, namely the subalgebra consisting of all elements that possess a 'formal adjoint'.

**THEOREM 1.** *Let  $V$  be a complex vector space that admits an inner product  $\langle \cdot | \cdot \rangle$ , let  $L(V)$  denote the algebra of all linear transformations of  $V$  and let*

$$A(V) := \{a \in L(V) : (\exists b \in L(V))(\forall x, y \in V) \langle ax|y \rangle = \langle x|by \rangle\}.$$

*Then*

- (i)  $A(V)$  is a subalgebra of  $L(V)$ ,
- (ii) to each  $a \in A(V)$  there corresponds a unique  $a^* \in A(V)$  such that, for all  $x, y \in V$ ,  $\langle ax|y \rangle = \langle x|a^*y \rangle$ ,
- (iii) the mapping  $*$  :  $A(V) \rightarrow A(V)$ ,  $a \mapsto a^*$  is a special involution.

**PROOF.** (i) This is routine.

(ii) Let  $a \in A(V)$ . Suppose that  $b, c \in L(V)$  are such that, for all  $x, y \in V$ ,  $\langle ax|y \rangle = \langle x|by \rangle = \langle x|cy \rangle$ . Then, for all  $y \in V$ ,  $\langle (b - c)y|(b - c)y \rangle = 0$  and so  $\langle (b - c)y|0 \rangle = 0$ . Thus  $b = c$ . This establishes the existence of a unique  $a^* \in L(V)$  such that, for all  $x, y \in V$ ,  $\langle ax|y \rangle = \langle x|a^*y \rangle$ . Then, for all  $x, y \in V$ ,

$$\langle a^*x|y \rangle = \overline{\langle y|a^*x \rangle} = \overline{\langle ay|x \rangle} = \langle x|ay \rangle$$

and so  $a^* \in A(V)$ . (This argument shows also that  $a^{**} = a$ .)

(iii) It is easily checked that  $*$  is an involution on  $A(V)$ : we must prove that it is special.

Let  $T$  be a nonempty finite subset of  $A(V)$ . Write

$$U := \{a - b : a, b \in T\}.$$

We show first that there exists a linear functional  $\chi$  on  $A(V)$  such that

- (1)  $(\forall a \in A(V)) \quad \chi(a^*a)$  is real and nonnegative,
- (2)  $(\forall u \in U) \quad \chi(u^*u) = 0$  implies  $u = 0$ .

If  $U = \{0\}$  we take  $\chi$  to be the zero mapping. Suppose, therefore, that  $U \neq \{0\}$ . Let  $u_1, u_2, \dots, u_n$  be the nonzero elements of  $U$ . For each  $r \in \{1, 2, \dots, n\}$ , choose

$x_r \in V$  such that  $u_r x_r \neq 0$ . We define  $\chi : A(V) \rightarrow \mathbb{C}$  by

$$(\forall a \in A(V)) \quad \chi(a) := \sum_{i=1}^n \langle ax_i | x_i \rangle.$$

It is clear that  $\chi$  is linear. To establish (1), we simply note that, for any  $a \in A(V)$ ,

$$\chi(a^*a) = \sum_{i=1}^n \langle a^*ax_i | x_i \rangle = \sum_{i=1}^n \langle ax_i | ax_i \rangle;$$

further, (2) holds, since, for  $r \in \{1, 2, \dots, n\}$ ,

$$\chi(u_r^*u_r) = \sum_{i=1}^n \langle u_r x_i | u_r x_i \rangle \geq \langle u_r x_r | u_r x_r \rangle > 0.$$

Choose  $t \in T$  such that  $\chi(t^*t) = \max\{\chi(a^*a) : a \in T\}$ . Suppose that  $t^*t = a^*b$ , where  $a, b \in T$ . We complete the proof by showing that  $a = b$ . Since  $t^*t = (a^*b)^* = b^*a$ , we have that  $(a - b)^*(a - b) = a^*a + b^*b - 2t^*t$ . Hence, by (1) and the choice of  $t$ ,

$$0 \leq \chi((a - b)^*(a - b)) = \chi(a^*a) + \chi(b^*b) - 2\chi(t^*t) \leq 0$$

and so  $\chi((a - b)^*(a - b)) = 0$ . But  $a - b \in U$ . Hence, by (2),  $a = b$ . □

**COROLLARY 1 (Hofmann).** *The standard involution on a  $C^*$ -algebra is special.*

**PROOF.** It is sufficient to consider the case of the  $C^*$ -algebra  $B(V)$  of all bounded linear operators on a complex Hilbert space  $V$ . Clearly  $B(V)$  is a star subalgebra of  $A(V)$  and the standard involution on  $B(V)$  is the restriction of the involution  $*$  on  $A(V)$ . Since  $*$  is special, the result follows. □

Let  $I$  be a nonempty set. An  $I \times I$  complex matrix  $[\alpha_{ij}]$  is said to be *row-finite* if and only if, for all  $i \in I$ , the set  $\{j \in I : \alpha_{ij} \neq 0\}$  is finite (possibly empty). Similarly,  $[\alpha_{ij}]$  is *column-finite* if and only if, for all  $j \in I$ ,  $\{i \in I : \alpha_{ij} \neq 0\}$  is finite. The set  $\mathbb{C}_I$  of all  $I \times I$  complex matrices that are both row-finite and column-finite is a complex algebra under the usual matrix operations and is closed under hermitian conjugation.

**COROLLARY 2.** *Let  $I$  be a nonempty set. Then hermitian conjugation is a special involution on  $\mathbb{C}_I$ .*

**PROOF.** Let  $V$  denote the complex vector space consisting of all  $I \times \{1\}$  ‘column’ vectors with at most finitely many nonzero entries. Then the mapping  $\theta : \mathbb{C}_I \rightarrow L(V)$  defined by  $\theta(a)x = ax$  ( $x \in V$ ), where  $ax$  is the usual matrix product, is an injective

homomorphism. Moreover,  $V$  admits an inner product  $\langle \cdot | \cdot \rangle$  defined by  $\langle x | y \rangle = \sum_i \xi_i \bar{\eta}_i$ , where  $\xi_i$  and  $\eta_i$  denote the  $i$ th components of  $x$  and  $y$  respectively; and it is easily seen that, for all  $a \in \mathbb{C}_I$  and all  $x, y \in V$ ,  $\langle ax | y \rangle = \langle x | a^\dagger y \rangle$ , where  $a^\dagger$  denotes the hermitian conjugate of  $a$ . Thus, for all  $a \in \mathbb{C}_I$ ,  $\theta(a) \in A(V)$  and  $(\theta(a))^* = \theta(a^\dagger)$ . But, by the theorem,  $*$  is special. Hence, since  $\text{im } \theta$  is a star subalgebra of  $A(V)$  and  $\theta$  is injective, it follows that hermitian conjugation is a special involution on  $\mathbb{C}_I$ .  $\square$

Observe that if  $I$  is infinite then  $\text{im } \theta$  above contains unbounded linear operators on  $V$ .

Each of these corollaries generalizes the result, due to Lavers [2, Example 4], that, for any positive integer  $n$ , hermitian conjugation is a special involution on the algebra of all  $n \times n$  complex matrices.

A further application of Theorem 1 arises in the context of certain Banach algebras. Let  $S$  be a semigroup. We denote by  $l^1(S)$  the Banach algebra consisting of all functions  $a : S \rightarrow \mathbb{C}$  of countable support such that  $\sum_{x \in S} |a(x)| < \infty$ , where addition and scalar multiplication are the usual pointwise operations, multiplication is convolution, and the norm  $\| \cdot \|$  is defined by

$$(\forall a \in l^1(S)) \quad \|a\| := \sum_{x \in S} |a(x)|.$$

Now suppose that  $S$  is an inverse semigroup; thus, to each  $x \in S$  there corresponds a unique element  $x^{-1} \in S$  (the ‘inverse’ of  $x$ ) such that  $xx^{-1}x = x$  and  $x^{-1}xx^{-1} = x^{-1}$ . It is well known that inversion ( $x \mapsto x^{-1}$ ) is an involution on  $S$ . As is readily checked, inversion on  $S$  induces an involution  $^\dagger$  on  $l^1(S)$  by the rule that

$$(\forall x \in S) \quad a^\dagger(x) := \overline{a(x^{-1})}.$$

We now combine Corollary 1 with a result of Barnes [1] to show that  $^\dagger$  is special.

**THEOREM 2 (Crabb).** *Let  $S$  be an inverse semigroup. Then the involution on  $l^1(S)$  induced by inversion on  $S$  is special.*

**PROOF.** As above, let  $^\dagger$  denote the involution on  $l^1(S)$  induced by inversion on  $S$ . By [1, Theorem 2.3], there exists a Hilbert space  $V$  and a (continuous) injective homomorphism  $\theta : l^1(S) \rightarrow B(V)$ , the algebra of all bounded linear operators on  $V$ , such that, for all  $a \in l^1(S)$ ,  $(\theta(a))^* = \theta(a^\dagger)$ , where  $*$  denotes the standard involution on  $B(V)$ . But, by Corollary 1,  $*$  is special. Hence, since  $\text{im } \theta$  is a star subalgebra of  $B(V)$  and  $\theta$  is injective, it follows that  $^\dagger$  is special.  $\square$

This extends [3, Theorem 5.1].

## Acknowledgments

I am grateful to K. H. Hofmann and M. J. Crabb for supplying the results in Corollary 1 and Theorem 2 respectively.

## References

- [1] B. A. Barnes, 'Representations of the  $l^1$ -algebra of an inverse semigroup', *Trans. Amer. Math. Soc.* **218** (1976), 361–396.
- [2] D. Easdown and W. D. Munn, 'On semigroups with involution', *Bull. Austral. Math. Soc.* **48** (1993), 93–100.
- [3] D. Easdown and W. D. Munn, 'Trace functions on inverse semigroup algebras', *Bull. Austral. Math. Soc.* **52** (1995), 359–372.

Department of Mathematics  
University of Glasgow  
Glasgow G12 8QW  
Scotland  
U.K.  
e-mail: wdm@maths.gla.ac.uk