

A Tidal Dwarf Galaxy in the Hercules Cluster?

W. van Driel¹

¹ *Observatoire de Paris, GEPI, Meudon, France*

P.-A. Duc², P. Amram³, F. Bournaud², C. Balkowski¹, V. Cayatte¹,
J.M. Dickey⁴, H. Hernández⁵, J. Iglesias-Páramo⁶, K. O'Neil⁶,
P. Papaderos⁷, J.M. Vílchez⁸

² *CEA, Service d'Astrophysique, Gif-sur-Yvette, France*

³ *Laboratoire d'Astrophysique de Marseille, Marseille, France*

⁴ *Dept. of Astronomy, Univ. of Minnesota, Minneapolis, MN, USA*

⁵ *Arecibo Observatory, Arecibo, PR, USA*

⁶ *NRAO, Green Bank, WV, USA*

⁷ *Universitäts-Sternwarte, Göttingen, Germany*

⁸ *Instituto de Astrofísica de Andalucía (CSIC), Granada, Spain*

Abstract. A candidate Tidal Dwarf Galaxy, ce-61, was identified in the merger system IC 1182 in the Hercules supercluster. The multi-wavelength data we obtained so far do not prove, however, that it is kinematically detached from the IC 1182 system and gravitationally bound.

The Hercules supercluster ($D=150$ Mpc, $H_0=75$) is one of the most massive structures in the nearby Universe. We studied (Iglesias-Páramo et al. 2003) 22 H I-selected galaxies in this cluster, from the blind VLA H I survey by Dickey (1997), obtaining: deep CCD B , V and I -band surface photometry of 10 galaxies, optical spectroscopy of 8 of these, Arecibo H I observations of all 22 galaxies and H α line Fabry-Perot observations of the IC 1182 merger system.

Based on these multi-wavelength observations, the object ce-61 was identified as a candidate Tidal Dwarf Galaxy (TDG) in a tidal tail of the peculiar IC 1182 system. IC 1182 ($B_T=15.4$, $V=10,223$ km s⁻¹) shows several characteristics typical of a merger system, e.g., a blue optical jet-like structure towards the East and tidal debris towards the NW, and an extended H I distribution with two tidal tails, towards the E and the NW. The candidate TDG ce-61 ($M_B = -18.24$ mag) lies at the tip of the eastern optical/H I tail, at about 1.5 arcmin (65 kpc) projected distance from the centre of the parent system. Its CCD image shows two distinct peaks and the maximum in the H I tail coincides with the easternmost optical peak. Its metallicity (8.41) is on the high side for a dwarf galaxy of its luminosity, but typical for a TDG. It is a very gas-rich system, with an estimated M_{HI}/L_B ratio of $6 M_\odot/L_{\odot,B}$; its H I line width is about 220 km s⁻¹. CO line observations (Braine et al. 2001) show about $7 \times 10^9 M_\odot$ of H₂ in a resolved distribution in IC 1182, but none was detected in ce-61, putting an upper limit to its H₂ mass of about $6 \times 10^7 M_\odot$.

In order to study whether the TDG candidate is already kinematically detached from the IC 1182 system and gravitationally bound, we obtained and analyzed H α line Fabry-Perot observations and found (Bournaud, Duc & Amram 2003, in prep.) that: (1) the brightest knot at the tip of the tail, coinciding with ce-61, seems to be kinematically detached from the overall velocity field along the tail (which is governed by streaming motions); the offset is about 30 km s $^{-1}$. However, our numerical simulations show that this offset can be consistent with a projection effect along the line of sight (with the tail, seen edge-on, being bent in 3D space), (2) along direction 2 (see Fig.), there is a hint of an internal velocity gradient, of 70 km s $^{-1}$ maximum, associated with one of the knots. We lack the spatial resolution to confirm it, however.




Figure 1. H α line Fabry-Perot observations of the candidate TDG ce-61 in the merger system IC 1182: (top) the clumpy distribution of the H α gas in the tidal tail of IC 1182, superimposed on a V-band optical image of the system + HI column density contours; (bottom) an H α line position-velocity diagram along the tidal tail (following the lines marked '1' in the top panel).

References

- Braine J., et al. 2001, *A&A*, 378, 51
 Dickey J.M. 1997, *AJ*, 113, 1939
 Iglesias-Pramo J., van Driel W., Duc P.-A. 2003, *A&A*, 406, 453
 van Driel W., et al. 2003, *A&A*, 399, 433