

Exploring the transition into an asymmetric planetary nebulae

Mónica W. Blanco Cárdenas¹, Roberto Vázquez¹,
Hans Ulrich Kaeuff², Marco A. Gómez-Muñoz¹, Pedro F. Guillén¹,
Martín A. Guerrero³ and Luis F. Miranda³

¹Instituto de Astronomía, Universidad Nacional Autónoma de México,
Km 103 Carretera Tijuana-Ensenada 22860, Ensenada, Mexico
email: mwblanco@astro.unam.mx

²European southern Observatory,
Karl Schwarzschild-Str 2 85748, Garching bei Muenchen, Germany

³Instituto de Astrofísica de Andalucía,
Glorieta de la Astronomía S/N 18008, Granada, Spain

Abstract. We present studies using different observational techniques, along different frequencies, aiming to resolve and investigate jets, outflows, as well as compact and innermost regions of asymmetric planetary nebulae (PNe) and objects in transition to PN. All the information gathered allow us to explore the kinematics and other important properties of the structures that play a crucial role in the shaping of complex PNe morphologies, in particular, we explore the role of disks/tori as collimating engine of extreme axisymmetric PNe.

Keywords. Stars, circumstellar mater, outflows.

1. The complex sculpting of an asymmetric planetary nebula

Several planetary nebulae (PNe) are asymmetric and high velocity jets and outflows are common features in these evolved objects. Furthermore, during the last years, steady and/or slowly rotating disks have been discovered in the cores of extreme axisymmetric PN. Infrared observations provide the opportunity to investigate the innermost dusty regions that usually are related with torii and disks. In this work, we present proto-PNe and young PNe with at least one observational proof of torus or disk, and we explore these structures using other techniques: spectro-astrometry (SA) using CRIRES-VLT (Blanco Cárdenas *et al.* 2014) and analysis of its SED using available photometry of several missions from the visible to radio frequencies (2MASS, *MSX*, *WISE*, *IRAS*, *AKARI*, etc.) and the SED building and analysis tool IRIS (cxc.cfa.harvard.edu/iris/v2.1).

2. Overview and preliminary results

Proto-PN AFGL 915. This source (a.k.a The Red Rectangle) has a well-known rotating disk of few hundreds of AU (Bujarrabal *et al.* (2013)) and references therein) surrounded by a steady disk of silicates (Waters *et al.* (1998)). Blanco Cárdenas *et al.* (2014) detected a smaller spectral signature of this rotating disk (few tens of AU, ~ 100 mas) using the SA technique. As for the SED, three components fit well: a 7000 K Blackbody (BB), a rotating disk in mid-IR wavelengths, and a 170 K BB for the larger wavelengths.

IRAS 18454+0001: an extremely young PN? Given its compact size ($\sim 1''$) and the lack of an optical counterpart this source must be a bipolar PN in the making.




Figure 1. SEDs of the sample. From left to right and up to bottom: AFGL 915, IRAS 18454+0001, M2-9, Mz3, and M1-16.

Results outstanding the equatorial structure clearly dominated by the $\lambda 8.6 \mu\text{m}$ PAH filter of VISIR-VLT (Blanco Cárdenas *et al.* 2013), whereas the spitzer spectrum shows the presence of carbon-rich material (mid-IR PAH features) and confirms the presence of ionised material ([NeII] $\lambda 12.8 \mu\text{m}$). According to the SED fitting, a very hot star (83,000 K) and a disk beyond the mid-IR (larger dust grains) are needed, among other components of warmer material in the IR.

Young PNe: M2-9, Mz3 and M1-16. Silicate disks were found in the central regions of M2-9 and Mz3 (Chesneau *et al.* 2007 and Lykou *et al.* 2011). On the other hand, CO ($J=2-1$) observations confirms the presence of two concentric rings in M2-9 (Castro-Carrizo *et al.* 2012) and a torus in M1-16 (Huggins *et al.* 2000). These three sources have fast collimated outflows (up to $\sim 300 \text{ km s}^{-1}$). Our CRIFRES spectra does not show any evidence of disks in M2-9 and Mz3, only the origin of the bipolar lobes of M2-9 is likely detected. The SEDs of M2-9 and Mz3 are alike. However, for M2-9 two BB of 96,000 K and 3,200 K fit the optical data, whereas only one BB of 30,000 K fits the SED of Mz3. A rotating disk in the mid-IR and components around 100 K fit the longer wavelengths. M1-16 has a prominent optical component fitted by 32,000 K BB. A rotating disk fits well the data, nevertheless, this disk may have a hotter component in the near IR. Once again, colder BBs are needed to fit the data beyond mid-IR wavelengths.

References

- Blanco Cárdenas, M., Kaeulf, H. U., Guerrero, M. A. *et al.* 2014, *A&A*, 566, A133
 Blanco Cárdenas, Guerrero, M. A., Ramos-Larios, G *et al.* 2013, *A&A*, 551, A64
 Bujarrabal, V., Castro-Carrizo, A., Alcolea, J., *et al.* 2013, *A&A*, 557, L11
 Castro-Carrizo, A., Neri, R., Bujarrabal, V. *et al.* 2012, *A&A*, 545, A1
 Chesneau, O., Lykou, B., Balick, B. *et al.* 2007, *A&A*, 473, L29
 Huggins, P. J., Forveille, T., Bachiller, R., & Cox, P. 2000, *ApJ*, 544, 889
 Lykou, B., Chesneau, O., Zijlstra, A. A. *et al.* 2011, *A&A*, 527, A105
 Waters, L. B. F. M., Waelkens, C., van Winckel, H. *et al.* 1998, *Nature*, 67, 4679