

($p = 0.162$) or the nature of admission ($p = 0.586$) (Table 1). When analyzing the voluntary nature of admission based on age and sex, statistically significant differences were only found in the female group ($p = 0.01$), resulting in a 9.18 year age difference among those admitted voluntarily (Table 2). The model that best predicted the probability of involuntary admission in individuals with psychotic disorders included the sex variable (OR = 4.88) and the interaction between sex and age (OR = 0.97) (Table 3).

Table 1: Differences between sex regarding voluntariness of patients with psychotic disorders.

	Male	Female	<i>p</i> value
N (%)	122 (48%)	132 (52%)	
Age, m (SD)	38.39 (16.64)	44.15 (18.44)	0.162
Admissions, N (%)			
Voluntary	38 (31.1%)	37 (28.0%)	0.586
Involuntary	84 (68.9%)	95 (72.0%)	

Table 2: Analysis of voluntariness by sex and age.

Age, m (SD)	Voluntary	Involuntary	<i>p</i> value
Male	37.45 (16.38)	38.81 (16.84)	0.677
Female	50.76 (18.19)	41.58 (17.98)	0.01*
Total	44.01 (18.44)	40.28 (17.46)	0.127

Table 3: Predictors of involuntariness in psychotic patients: Logistic regression model (ENTER METHOD).

Predictor	-2log likelihood	Nagelkerke		OR* (95% CI)*	<i>p</i> value
		R2	χ^2 (df*)		
	301.22	0.039	0.03 (1)		
Age				1.01 (0.98; 1.03)	0.674
Sex				4.88 (1.15; 20.72)	0.032*
Age x Sex Interaction				0.97 (0.94; 0.99)	0.046*

Conclusions: Young women with psychotic disorders face a higher risk of involuntary admissions, emphasizing the need for gender-specific strategies to improve care of these patients.

Disclosure of Interest: None Declared

Others

EPV1135

Current situation regarding psychedelics and magic mushroom in Korea

J. S. Seo

Psychiatry, Gwangmyeong Hospital, Chung-Ang University, Korea, Gwangmyeong-si, Korea, Republic Of
doi: 10.1192/j.eurpsy.2024.1695

Introduction: Recently, the pros and cons have been debating in Korea even before the approval of use of medical marijuana with very strict limitations. And the next controversial topic is psychedelics. In 1890, when mescaline was first isolated from peyote cactus, clinical researches began, but due to its harmful effects, it was thereafter legally prohibited in 1970 in USA. However, a pernicious debate over the medical efficacy of psychedelic drugs has begun again with the release of a study that uses psychedelic mushrooms to be effective against treatment-resistant depression, alcohol dependence, and depression and anxiety in terminal cancer patient.

Objectives: To make a consensus on the medical use of these, we reviewed wild mushrooms containing hallucinogenic ingredients living in Korea.

Methods: To make a consensus on the medical use of these, we reviewed wild mushrooms containing hallucinogenic ingredients living in Korea.

Results: Mushrooms have long been popular as a food ingredient in Korea. Psilocybin, a classical psychedelic, can be obtained from magic mushroom (*Psilocybe cubensis*). The psilocybin on the CNS and causes hallucinations. Intoxication symptoms include pleasant or nervousness, sudden laughter, hallucinations, visual impairment, tachycardia and hypertension, reflexes, agitation, cognitive impairment, confusion, and aggressive behavior. These symptoms last for 2-4 hours after ingestion, and most disappear within six hours.

Among 114 species of *Psilocybe* containing psilocybin around the world, only five wild mushrooms found in Korea that cause nervous system hallucinations are as follows: *P. argenteipes*, *P. coprophila*, *P. perdaria*, and *P. subcarulipes*.

In Korea, there is acute poisoning case suffering with GI symptoms caused by mushrooms, but it is difficult to find records of abuse or dependences case caused by psychedelic mushrooms. In addition, although oriental medicine treatment is relatively active, it is not used as an herbal medicine.

Conclusions: Currently, the Korean government classifies psychedelic mushroom-derived substances, Psilocybin and Psilocin, as psychotropic drugs by law. If researcher intends to clinical trial with even very small amount of it for academic purpose, it is only possible after obtaining approval from Korean FDA. In order to determine the usefulness of psychedelics, many clinical studies are needed in Korea.

Disclosure of Interest: None Declared