

Serum *n*-3 long-chain PUFA differ by sex and age in a population-based survey of New Zealand adolescents and adults

Francesca L. Crowe¹, C. Murray Skeaff^{1*}, Timothy J. Green¹ and Andrew R. Gray²¹Department of Human Nutrition, University of Otago, Dunedin, New Zealand²Department of Social and Preventive Medicine, University of Otago, Dunedin, New Zealand

(Received 1 February 2007 – Revised 16 May 2007 – Accepted 31 May 2007)

A higher proportion of *n*-3 long-chain PUFA in tissue lipids has been associated with a lower risk of CVD and some cancers. Diet is an important predictor of *n*-3 long-chain PUFA composition; however, the importance of non-dietary factors such as sex and age is unclear. We measured the proportion of *n*-3 long-chain PUFA in serum phospholipid, cholesterol ester and TAG of 2793 New Zealanders 15 years or older who participated in the 1997 National Nutrition Survey to determine differences by sex and age. Women had lower proportions of EPA and docosapentaenoic acid in phospholipid, by 0.07 ($P=0.004$) and 0.10 ($P<0.001$) mol%, respectively, and a higher proportion of DHA by 0.16 mol% ($P=0.001$) compared with men. Intake of fish fat did not differ between men and women. There was a positive association between age and the proportion of EPA and DHA in phospholipid ($P<0.001$). The sex differences in EPA and DHA were similar at all ages. Similar sex and age differences in serum cholesterol ester *n*-3 long-chain PUFA were found; only age differences were found in serum TAG. Sex and age differences in *n*-3 long-chain PUFA occur in the general population. Men and women may need to be considered separately when examining the association between disease risk and biomarkers of *n*-3 fatty acids.

Nutrition surveys: *n*-3 fatty acids: Docosahexaenoic acid: New Zealanders

An inverse association between *n*-3 long-chain PUFA in tissues and risk of CVD and some cancers is a consistent finding of observational studies^{1–4}. An important element in judging the causal nature of this association is to understand the factors that influence the *n*-3 long-chain PUFA composition of tissue lipids.

Increased dietary intake of *n*-3 long-chain PUFA – EPA and DHA – increases the proportion of these fatty acids in plasma, blood cells and adipose tissue^{5–8}. Furthermore, in cross-sectional studies, the proportion of tissue fatty acids as *n*-3 long-chain PUFA is correlated with *n*-3 long-chain PUFA intake^{9,10}. A number of non-dietary determinants such as sex and age may also affect *n*-3 long-chain PUFA composition. However, the extent to which non-dietary determinants influence tissue *n*-3 long-chain PUFA is not well described. A consequence of not considering the effect of non-dietary determinants of *n*-3 long-chain PUFA concentrations is that erroneous conclusions about *n*-3 biomarker–disease risk associations may be reached. For example, if sex, in addition to being an independent predictor of CVD risk, is also a biological determinant of tissue *n*-3 long-chain PUFA composition it may be more appropriate to examine the *n*-3 biomarker–disease risk association separately in men and women to minimise bias.

Evidence of sex differences in *n*-3 fatty acid metabolism has been reported. Results from two studies with small sample

sizes showed that women, when fed ¹³C-labelled α -linolenic acid, had a greater capacity than men to synthesise ¹³C-labelled DHA^{11–14}. The extent to which this sex difference in *n*-3 fatty acid metabolism, assessed under controlled conditions, is reflected in non-controlled conditions at a population level where dietary intake of preformed DHA may be the predominant determinant of tissue DHA composition is unclear. In three population-based surveys women had a higher percentage of total fatty acids as DHA in plasma^{15,16} and adipose¹⁷ lipids than men; however, in another survey no sex difference was reported¹⁸.

Age-related increases in *n*-3 long-chain PUFA composition of tissues have also been reported^{15–19}; however, in three of these studies in which fish intake was reported, the age-related changes in plasma DHA paralleled increases in fish consumption^{15,16,18}.

The objective of the present study was to determine in a population-based sample of people living in New Zealand if *n*-3 long-chain PUFA differed between men and women and with age.

Subjects and methods

The sample from the 1997 National Nutrition Survey (NNS97) was taken from the 1996/97 New Zealand Health Survey, a population-based survey that assessed the health status of non-institutionalised New Zealanders aged 15 years and

Abbreviations: NNS97, 1997 National Nutrition Survey.

* Corresponding author: Dr C. Murray Skeaff, P.O. Box 56, Dunedin, New Zealand, fax +64 3 479 7958, email murray.skeaff@otago.ac.nz

older. More detailed descriptions of the methods for the NNS97 have been published^{20–22}. In brief, the New Zealand Health Survey used an area-based sampling frame with a three-staged stratified design that consisted of a selection of a set of 18 800 geographical areas (the primary sampling units), households within these primary sampling units and then a randomly chosen respondent within a household. The response rate for the New Zealand Health Survey was 74% (*n* 7862); of these participants, 4636 provided dietary information in the NNS97 and 3223 gave a blood sample. The fatty acid composition of at least one serum lipid fraction was available for 2793 participants; thus, the fatty acid analysis represents 36% of the original New Zealand Health Survey. Fourteen ethics committees throughout New Zealand approved the survey and all participants or the legal guardian of those under 18 years provided written informed consent. The procedures followed were in accordance with the ethical standards of the New Zealand Health Research Council.

The data collection for the NNS97 was undertaken over a 12-month period from December 1996 to November 1997. During a home visit, dietary information was collected from each participant using a computer-assisted 24 h recall method²¹. Intake of meat and fish fats were calculated from the food group analysis used in the NNS97. Freshwater fish is not sold in New Zealand, therefore all fish consumed by New Zealanders was assumed to be of marine origin. Age and ethnicity were self-reported and ethnicity was categorised into three groups: New Zealand Maori, Pacific People, and New Zealand European and other. The term 'Pacific' includes those who identify as Samoan, Tongan, Cook Island Maori, Niuean, Tokelauan, Fijian and other Pacific ethnic groups. All remaining participants were assigned to the New Zealand European and other group. Height and weight were measured according to standard techniques and BMI (kg/m²) was calculated. Participants were asked three questions pertaining to their smoking and categorised as smokers or non-smokers, the latter including former smokers. At the end of the home visit, blood was drawn from the participants from an antecubital vein into vacuum evacuated tubes with no anticoagulant. Participants were not required to fast. The tubes were centrifuged and the serum aliquoted into cryovials for storage at –80°C.

Laboratory analysis

Details of the fatty acid analysis have been published elsewhere^{5,23}. In brief, lipids were extracted from 400 µl serum according to the method of Bligh & Dyer²⁴. Serum phospholipid, cholesterol ester and TAG were isolated using TLC and fatty acids were analysed using a DB-225 narrowbore column (30 m × 0.25 mm internal diameter; film thickness 0.25 µm; J&W Scientific, Deerfield, IL, USA) on a HP-6890 gas chromatograph with flame ionisation detection (Agilent, Palo Alto, CA, USA). The fatty acid results are reported as percentage of total fatty acids, on a molar basis (i.e. mol%). Precision of the fatty acid measurement was determined by repeat analysis of a pooled serum sample, one pooled sample was analysed for every twenty NNS97 samples. The CV for EPA (20:5*n*-3), docosapentaenoic acid (22:5*n*-3) and DHA (22:6*n*-3) in serum phospholipid (*n* 102) were 14, 12 and 13%, respectively. The docosapentaenoic acid composition of serum cholesterol ester and TAG was not reported because the

amount of this fatty acid in many serum samples was below the detection limits of the analytical methods.

Statistical analysis

All statistical analyses were carried out on Stata Statistical Software, Release 9 (Statacorp LP, College Station, TX, USA) using the survey commands where appropriate to control for the survey design or using weighted regression and adjusting the standard errors to account for the clustering. For the unadjusted analysis, age was coded as a categorical variable (15–24 years, 25–44 years, 45–64 years, 65+ years) and the survey commands were used to calculate the unadjusted means for the *n*-3 long-chain PUFA composition of serum phospholipid, cholesterol ester and TAG by sex and by sex and age category. Survey regression was used to test for sex differences and linear trends across the age categories for the *n*-3 long-chain PUFA. The adjusted means for each of the *n*-3 long-chain PUFA by sex and by sex and age were estimated using multiple fractional polynomial regression²⁵ controlling for potential confounders. All variables were entered into the model simultaneously using fractional polynomials for the continuous variables (age, BMI and fish fat (g/d)) or as indicator variables for the categorical variables (sex, ethnicity and smoking). The adjusted means were calculated for 20, 35, 54 and 73 years; the mean age of each age category. Sex-by-age and sex-by-BMI interactions were also tested in the models where appropriate and included if *P* < 0.05. The effects of sex and age on *n*-3 long-chain PUFA composition were considered statistically significant if *P* < 0.05. Similar analyses were conducted for the mean fish and meat fat intake by sex and by sex and age. All tests of significance were two-sided.

Results

The fatty acid composition of serum phospholipid, cholesterol ester and TAG were available for 2416, 2393 and 2402 participants, respectively, of the 3223 stored serum samples from the NNS97. The fatty acid composition of at least one serum lipid fraction was analysed in 2793 participants. The distribution of sex, ethnicity and smoking as well as the mean age, BMI, and intake of fish and meat fats were similar between participants for whom a fatty acid measurement was made (*n* 2793), and participants in the 1996/97 New Zealand Health Survey (*n* 7862) and NNS97 (*n* 4636) (Table 1).

The mean percentage of total fatty acids in serum phospholipid as EPA was 0.07 mol% (95% CI 0.02, 0.12) lower in women compared with men; in cholesterol ester, EPA was 0.09 mol% (95% CI 0.04, 0.14) lower in women (Table 2). Women also had a lower proportion of docosapentaenoic acid in serum phospholipid than men, by 0.10 mol% (95% CI 0.07, 0.13). In contrast, the proportion of DHA in serum phospholipid and cholesterol ester was higher in women than in men, by 0.16 mol% (95% CI 0.07, 0.25) and 0.02 mol% (95% CI 0.01, 0.04), respectively. The sex differences in fatty acid composition were increased slightly after adjustment of the regression model for age, BMI, ethnicity and smoking. There were no significant differences in the fatty acid composition of serum TAG between men and women.

Table 1. Characteristics of participants

	NZHS		NNS97		NNS97 fatty acid	
	(<i>n</i> 7862)	%	(<i>n</i> 4636)	%	(<i>n</i> 2793)	%
Percentage of original NZHS cohort		–		59		36
Sex <i>n</i> (%)						
Male	3258	41	1927	42	1246	45
Female	4604	59	2709	58	1547	55
Age category (y)						
15–24	1050	13	600	13	332	12
25–44	3221	41	1964	42	1227	44
45–64	2063	26	1255	27	765	27
65 +	1528	19	817	18	469	17
Ethnicity						
NZEO	5896	75	3627	78	2286	82
NZ Maori	1321	17	703	15	368	13
Pacific	645	8	306	7	139	5
Smoking category						
Never/former	5582	71	3365	73	2030	73
Current	2280	29	1271	27	763	27
			Mean	SD	Mean	SD
BMI (kg/m ²)	–	–	26.7	5.4	26.5	5.1
Fish fat (g/d)	–	–	1.6	7.8	1.6	6.9
Meat fat (g/d)	–	–	13.3	21.8	12.8	20.7

NNS97, 1997 National Nutrition Survey; NZEO, New Zealand European and others; NZHS, 1996/97 New Zealand National Health Survey.

The unadjusted means for the *n*-3 long-chain PUFA composition of serum phospholipid, cholesterol ester and TAG by sex and age are shown in Table 2. The proportion of EPA in serum phospholipid, cholesterol ester and TAG increased significantly across the age categories for both men and women ($P < 0.05$). The proportion of DHA in phospholipid for men and women and in cholesterol ester for men only increased across the age categories ($P < 0.05$), as did the proportion of docosapentaenoic acid in phospholipid for women only ($P < 0.001$).

The multivariate-adjusted means, after controlling for BMI, ethnicity and smoking, for *n*-3 long-chain PUFA in serum phospholipid by sex and age are shown in Fig. 1. Including fish fat intake (g/d) in the regression model had little influence on the adjusted means; therefore, it was omitted from the model. The age-related increase in the proportion of total fatty acids as EPA (sex by age interaction, $P = 0.186$) and DHA (sex by age interaction, $P = 0.188$) in serum phospholipid was similar in men and women. The proportion of EPA in serum phospholipid increased by 0.31 mol% between the ages of 20 and 73 years. The increase in DHA between 20 and 73 years was 0.28 mol%. At all ages, the proportion of EPA was lower and DHA higher in women compared with men. The docosapentaenoic acid composition of serum phospholipid increased with age in women ($P < 0.001$) but did not change with age in men ($P = 0.591$), the interaction of sex by age was significant ($P = 0.002$). The difference in docosapentaenoic acid composition of serum phospholipid between 20 and 73-year-old women was 0.18 mol%. Similar age-related increases for EPA and DHA in serum cholesterol ester and TAG were also found (results not shown).

The mean intake of fish fat and meat fat for men and women who had a fatty acid value for at least one serum lipid fraction are presented in Table 3. There was no significant difference in the intake of fish fat between men and women, before ($P = 0.106$) or after ($P = 0.106$) adjusting for

age, BMI, ethnicity and smoking. The age-related increase in fish fat intake was similar in men and women. The difference in fish fat intake between ages 20 and 73 years was 1.1 g/d for men and women. Intake of meat fat (unadjusted) was 8.2 (95% CI 6.2, 10.1) g/d higher in men than women ($P < 0.001$). Age was a significant predictor of meat fat intake ($P = 0.029$). Mean intake of meat fat was 1.0 g/d higher in participants aged 20 years compared with 73 years.

Discussion

We have shown that men and women differ in the relative proportions of *n*-3 long-chain PUFA in serum lipids in a population-based survey of New Zealanders. The most striking result was that women had lower proportions of EPA and docosapentaenoic acid than men, whereas, for DHA, the sex difference was reversed, and women had higher levels than men. This differential pattern of *n*-3 long-chain PUFA in serum lipids between men and women suggests sex differences in fatty acid metabolism. One would expect that a higher consumption of foods supplying *n*-3 long-chain PUFA in one sex or the other would be associated with higher proportions of all three *n*-3 long-chain PUFA not just one or two fatty acids as we found. Intake of individual *n*-3 long-chain PUFA was not estimated in the NNS97 because the New Zealand food composition database does not have this information. However, fish is the richest source of *n*-3 long-chain PUFA and the results of the 24 h diet recall showed no differences in fish fat intake between men and women. Meat fat is also a source of *n*-3 long-chain PUFA^{26,27}, and men consumed more meat fat than women. Thus, intake of all three *n*-3 long-chain PUFA was probably slightly higher in men than women. The higher intake of *n*-3 PUFA in men is consistent with a higher proportion of EPA and docosapentaenoic acid in serum phospholipid of men

Table 2. *n*-3 Long-chain PUFA composition of serum lipids by sex and age

	Unadjusted means			Adjusted means*			Age group (years), unadjusted means								<i>P</i> for trend†
	15 + (years)		<i>P</i> †	15 + (years)		<i>P</i>	15–24		25–44		45–64		65 +		
	Mean	SEM		Mean	SEM		Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM	
EPA (mol%)															
Phospholipid															
Men (<i>n</i> 1079)	1.05	0.02		1.07	0.02		0.86	0.05	1.07	0.03	1.14	0.04	1.12	0.05	< 0.001
Women (<i>n</i> 1337)	0.98	0.02	0.004	1.00	0.02	0.002	0.78	0.06	0.93	0.02	1.08	0.03	1.14	0.05	< 0.001
Cholesterol ester															
Men (<i>n</i> 1081)	1.09	0.02		1.12	0.02		0.88	0.05	1.09	0.03	1.21	0.04	1.17	0.04	< 0.001
Women (<i>n</i> 1308)	1.00	0.02	< 0.001	1.02	0.02	< 0.001	0.77	0.05	0.94	0.02	1.12	0.03	1.22	0.06	< 0.001
TAG															
Men (<i>n</i> 1071)	0.27	0.01		0.27	0.01		0.22	0.01	0.28	0.01	0.29	0.01	0.25	0.01	0.036
Women (<i>n</i> 1331)	0.26	0.01	0.224	0.26	0.01	0.205	0.20	0.02	0.26	0.01	0.28	0.01	0.29	0.02	< 0.001
Docosapentaenoic acid (mol%)															
Phospholipid															
Men	0.95	0.01		0.95	0.04		0.93	0.05	0.96	0.01	0.93	0.02	0.97	0.02	0.701
Women	0.85	0.01	< 0.001	0.85	0.09	0.001	0.76	0.03	0.83	0.01	0.90	0.02	0.93	0.02	< 0.001
DHA (mol%)															
Phospholipid															
Men	2.55	0.03		2.57	0.03		2.29	0.09	2.55	0.05	2.69	0.06	2.67	0.07	< 0.001
Women	2.71	0.03	0.001	2.72	0.03	0.002	2.58	0.12	2.67	0.04	2.81	0.07	2.79	0.08	0.035
Cholesterol ester															
Men	0.50	0.01		0.50	0.01		0.44	0.02	0.50	0.01	0.51	0.01	0.52	0.01	< 0.001
Women	0.52	0.01	0.007	0.52	0.01	0.036	0.50	0.03	0.51	0.01	0.54	0.01	0.54	0.02	0.097
TAG															
Men	0.49	0.01		0.49	0.01		0.42	0.03	0.52	0.03	0.51	0.02	0.47	0.03	0.140
Women	0.52	0.02	0.200	0.52	0.02	0.191	0.49	0.07	0.53	0.02	0.52	0.02	0.52	0.03	0.795

Serum *n*-3 fatty acids in New Zealand

* Adjusted for age, BMI, ethnicity and smoking using multiple fractional polynomial regression.
 † Calculated using survey regression.

Table 3. Dietary fat intake by sex and age

	Unadjusted means		Adjusted means*		Adjusted means†						P values for adjusted analysis			
	15 + years		15 + years		20 years		35 years		54 years		73 years		Age	Sex×age
	Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM		
Fish fat (g/d)														
Men	1.7	0.2	1.8	0.2	1.0	0.2	1.8	0.2	2.0	0.3	2.1	0.3	< 0.001	0.152
Women	1.3	0.2	1.3	0.2	0.5	0.2	1.3	0.2	1.6	0.2	1.6	0.2		
Meat fat (g/d)														
Men	16.8	1.0	16.4	0.9	15.7	1.2	17.9	1.1	17.2	1.0	14.7	1.2	0.029	0.146
Women	8.6	0.4	9.1	0.4	7.7	0.9	9.9	0.7	9.2	0.6	6.7	1.0		

* Adjusted for age, BMI, ethnicity and smoking using multiple fractional polynomial regression.

† Adjusted for BMI, ethnicity and smoking using multiple fractional polynomial regression.

Calculated using survey regression.




Fig. 1. *n*-3 Long-chain PUFA composition of serum phospholipid by age for men (○) and women (●). Values are means with their standard errors depicted by vertical bars. There was a positive association between age and EPA ($P < 0.001$) and DHA ($P < 0.001$) but not docosapentaenoic acid ($P = 0.795$) after adjusting for BMI, ethnicity and smoking using multiple fractional polynomial regression. Interactions between sex and age were significant for docosapentaenoic acid ($P = 0.002$) but not EPA ($P = 0.186$) or DHA ($P = 0.188$).

but is not consistent with their lower proportion of DHA if diet was the only determinant of fatty acid status.

Sex differences in *n*-3 long-chain PUFA composition of tissues have been reported previously in most^{15–17} populations.

Higher proportions of DHA in women compared with men have been reported in all but one¹⁸ of four population surveys^{15–17}. Similar differences have been reported in the Norfolk cohort of the European Prospective Investigation into Cancer and Nutrition²⁸ and in another small survey of men and women²⁹, although participant recruitment in both these studies was by convenience sampling rather than population-based. The differences in DHA were not explained by diet because fish consumption in women was not higher than in men^{15,16}. Interestingly, in a study of people living in southern Quebec, women consumed less fish than men yet had a higher proportion of DHA in plasma phospholipids¹⁶. With the exception of one¹⁵, the proportion of EPA in plasma phospholipids of women was not different from men^{16,18}. The present results differ in this regard, because we report a lower proportion of EPA in serum phospholipid of women than men, although the magnitude of difference we report is small.

The higher DHA content of serum phospholipid in women compared with men is consistent with evidence from stable-isotope feeding studies that show women have a higher capacity than men to synthesise DHA from α -linolenic acid^{11–14}. Other mechanisms may also be in play. For example, it was demonstrated in a stable-isotope feeding trial that DHA can be retroconverted to docosapentaenoic acid and EPA³⁰. It is possible that women have a lower capacity than men to retroconvert DHA, which would explain the sex differences we report. There is evidence to suggest that amongst women of postmenopausal age, hormone replacement therapy reduces the retroconversion of DHA to EPA³¹. Information regarding menopausal status or use of hormone replacement therapy was not collected from the participants in the NNS97. However, we found no evidence that the difference in DHA between men and women varied with age. This suggests that at a population level, neither menopausal status nor use of hormone replacement therapy has a major effect on serum DHA.

We found that EPA and DHA in serum phospholipid increased with age. Similar changes have been reported in other populations^{15–19}, and results from an intervention study suggest that older men have a greater capacity to incorporate dietary EPA into plasma phospholipid than younger men³². We extend these previous reports by showing the age-related increases did not differ between men and women. At all ages, women had lower EPA and higher DHA compared with men. Most evidence, to date, suggests the age-related changes in *n*-3 long-chain PUFA reflect changes in fish consumption^{15,16,18}. For example, results from a series of surveys in three ethnic groups in Quebec – southern Quebecers¹⁶, Inuit of Nunavik¹⁵ and the Cree of James Bay¹⁸ – show the age-related increases in *n*-3 long-chain PUFA paralleled increasing fish consumption. However, in the two largest surveys, the Scottish Heart Study (*n* 4114)¹⁷ and the Paris Prospective Study (*n* 3348)¹⁹, diet was not reported.

In the present study population, intake of fish fat increased with age. The difference in fish fat intake between 20- and 73-year-olds was 1.1 g/d, equivalent to a DHA intake of 110 mg/d – assuming each gram of fish fat contains approximately 100 mg DHA²⁶. Based on the results of dietary intervention trials it appears this amount of DHA may account for the age-related changes in serum phospholipid DHA in the present population. Yep *et al.* (2002)³³ found that 54 mg DHA/d

increased the percentage of this fatty acid in plasma phospholipid by 0.34%. This increase is larger than the 0.28 mol% increase we found between participants aged 20 and 73 years. The present results suggest the increase in EPA and DHA with age are related to changes in diet, rather than metabolism, because they paralleled increased fish fat intake.

The increase with age in the proportion of docosapentaenoic acid in serum phospholipid in women follows that of the other *n*-3 long-chain PUFA as well as the intake of fish fat. However, in men docosapentaenoic acid did not increase with age despite the increase in fish fat intake. This suggests that in men, serum phospholipid docosapentaenoic acid may be influenced more by metabolic rather than dietary factors.

The sex and age differences in the *n*-3 long-chain PUFA composition of serum cholesterol ester were similar to phospholipid, however, the sex differences were not significant for TAG. Thus, the metabolic processes that may influence these sex differences appear to be specific to phospholipid and cholesterol ester but not TAG. Alternatively, the absence of sex-related changes in TAG may be the result of collecting blood samples from participants who were not fasting. At a population level, the fat composition of recent meals will reflect habitual intake, although for the individual there may be a sizeable difference; therefore, any effect of postprandial blood sampling on serum TAG *n*-3 long-chain PUFA would be on the variability of the estimate and not on the mean fatty acid composition. This increased variability may have reduced the power to detect a statistically significant sex difference in the *n*-3 long-chain PUFA composition of serum TAG.

The metabolic and physiological effects of EPA and DHA differ. Both fatty acids are important structural components of cell membranes, however, EPA has more pronounced effects on eicosanoid production³⁴ whereas DHA has particular effects on membrane properties and cell signalling³⁵. Thus, it is not surprising that the relation between each of the *n*-3 long-chain PUFA and risk of CVD and some cancers is different^{1,36}. The present results suggest that there are sex differences in the regulation of *n*-3 long-chain serum PUFA and that men and women may need to be considered separately when examining the association between disease risk and biomarkers of *n*-3 long-chain PUFA.

Acknowledgements

This work was supported by a research grant from the National Heart Foundation of New Zealand and a University of Otago Research Committee, by means of a University of Otago Postgraduate Publishing Award (PhD). The New Zealand Ministry of Health funded the 1997 National Nutrition Survey. The Life in New Zealand group was responsible for conducting the 1997 National Nutrition Survey. Christian Thoma, Leanne Hodson, Belinda Hunter and Jody Miller helped with the laboratory analysis. Murray Skeaff conceived the study and obtained the funding for the fatty acid work. Francesca Crowe took overall responsibility for producing the fatty acid dataset. Andrew Gray was the consultant for the statistical analysis. All authors were involved in analysing the dataset, reviewing and interpreting the results and writing the manuscript.

References

- Simon JA, Hodgkins ML, Browner WS, Neuhaus JM, Bernert JT Jr & Hulley SB (1995) Serum fatty acids and the risk of coronary heart disease. *Am J Epidemiol* **142**, 469–476.
- Albert CM, Campos H, Stampfer MJ, Ridker PM, Manson JE, Willett WC & Ma J (2002) Blood levels of long-chain *n*-3 fatty acids and the risk of sudden death. *N Engl J Med* **346**, 1113–1118.
- Lemaitre RN, King IB, Mozaffarian D, Kuller LH, Tracy RP & Siscovick DS (2003) *n*-3 Polyunsaturated fatty acids, fatal ischemic heart disease, and nonfatal myocardial infarction in older adults: the Cardiovascular Health Study. *Am J Clin Nutr* **77**, 319–325.
- MacLean CH, Newberry SJ, Mojica WA, *et al.* (2006) Effects of omega-3 fatty acids on cancer risk: a systematic review. *JAMA* **295**, 403–415.
- Holub BJ, Bakker DJ & Skeaff CM (1987) Alterations in molecular species of cholesterol esters formed via plasma lecithin-cholesterol acyltransferase in human subjects consuming fish oil. *Atherosclerosis* **66**, 11–18.
- Blonk MC, Bilo HJ, Nauta JJ, Popp-Snijders C, Mulder C & Donker AJ (1990) Dose-response effects of fish-oil supplementation in healthy volunteers. *Am J Clin Nutr* **52**, 120–127.
- Leaf DA, Connor WE, Barstad L & Sexton G (1995) Incorporation of dietary *n*-3 fatty acids into the fatty acids of human adipose tissue and plasma lipid classes. *Am J Clin Nutr* **62**, 68–73.
- Katan MB, Deslypere JP, van Birgelen AP, Penders M & Zegwaard M (1997) Kinetics of the incorporation of dietary fatty acids into serum cholesteryl esters, erythrocyte membranes, and adipose tissue: an 18-month controlled study. *J Lipid Res* **38**, 2012–2022.
- Andersen LF, Solvoll K & Drevon CA (1996) Very-long-chain *n*-3 fatty acids as biomarkers for intake of fish and *n*-3 fatty acid concentrates. *Am J Clin Nutr* **64**, 305–311.
- Hjartaker A, Lund E & Bjerve KS (1997) Serum phospholipid fatty acid composition and habitual intake of marine foods registered by a semi-quantitative food frequency questionnaire. *Eur J Clin Nutr* **51**, 736–742.
- Burdge GC, Jones AE & Wootton SA (2002) Eicosapentaenoic and docosapentaenoic acids are the principal products of alpha-linolenic acid metabolism in young men. *Br J Nutr* **88**, 355–363.
- Burdge GC & Wootton SA (2002) Conversion of alpha-linolenic acid to eicosapentaenoic, docosapentaenoic and docosahexaenoic acids in young women. *Br J Nutr* **88**, 411–420.
- Pawlosky R, Hibbeln J, Lin Y & Salem N Jr (2003) *n*-3 Fatty acid metabolism in women. *Br J Nutr* **90**, 993–994.
- Pawlosky RJ, Hibbeln JR, Lin Y, Goodson S, Riggs P, Sebring N, Brown GL & Salem N Jr (2003) Effects of beef- and fish-based diets on the kinetics of *n*-3 fatty acid metabolism in human subjects. *Am J Clin Nutr* **77**, 565–572.
- Dewailly E, Blanchet C, Lemieux S, Sauve L, Gingras S, Ayotte P & Holub BJ (2001) *n*-3 Fatty acids and cardiovascular disease risk factors among the Inuit of Nunavik. *Am J Clin Nutr* **74**, 464–473.
- Dewailly EE, Blanchet C, Gingras S, Lemieux S, Sauve L, Bergeron J & Holub BJ (2001) Relations between *n*-3 fatty acid status and cardiovascular disease risk factors among Quebecers. *Am J Clin Nutr* **74**, 603–611.
- Tavendale R, Lee AJ, Smith WC & Tunstall-Pedoe H (1992) Adipose tissue fatty acids in Scottish men and women: results from the Scottish Heart Health Study. *Atherosclerosis* **94**, 161–169.
- Dewailly E, Blanchet C, Gingras S, Lemieux S & Holub BJ (2002) Cardiovascular disease risk factors and *n*-3 fatty acid status in the adult population of James Bay Cree. *Am J Clin Nutr* **76**, 85–92.
- Cambien F, Warnet JM, Vernier V, Ducimetiere P, Jacqueson A, Flament C, Orssaud G, Richard JL & Claude JR (1988) An epidemiologic appraisal of the associations between the fatty acids esterifying serum cholesterol and some cardiovascular risk factors in middle-aged men. *Am J Epidemiol* **127**, 75–86.
- Quigley R & Watts C (1997) *Food Comes First: Methodologies for National Nutrition Survey of New Zealand*. Wellington: Ministry of Health.
- Russell DG, Parnell WR & Wilson NC, (1997) *NZ Food: NZ People. Key Results of the 1997 National Nutrition Survey*. Wellington: Ministry of Health.
- Parnell WR, Wilson NC & Russell DG (2001) Methodology of the 1997 New Zealand National Nutrition Survey. *N Z Med J* **114**, 123–126.
- Crowe FL, Skeaff CM, Green TJ & Gray AR (2006) Serum fatty acids as biomarkers of fat intake predict serum cholesterol concentrations in a population-based survey of New Zealand adolescents and adults. *Am J Clin Nutr* **83**, 887–894.
- Bligh EG & Dyer WJ (1959) A rapid method of total lipids extractions and purification. *Can J Biochem Physiol* **37**, 911–917.
- Royston P, Ambler G & Sauerbrei W (1999) The use of fractional polynomials to model continuous risk variables in epidemiology. *Int J Epidemiol* **28**, 964–974.
- Quigley RJ, Burlingame BA, Milligan GC & Gibson JJ (1995) *Fats and Fatty Acids in New Zealand Foods*. Palmerston North: New Zealand Institute for Crop & Food Research.
- Howe P, Meyer B, Record S & Baghurst K (2006) Dietary intake of long-chain omega-3 polyunsaturated fatty acids: contribution of meat sources. *Nutrition* **22**, 47–53.
- Welch AA, Bingham SA, Ive J, Friesen MD, Wareham NJ, Riboli E & Khaw KT (2006) Dietary fish intake and plasma phospholipid *n*-3 polyunsaturated fatty acid concentrations in men and women in the European Prospective Investigation into Cancer-Norfolk United Kingdom cohort. *Am J Clin Nutr* **84**, 1330–1339.
- Bakewell L, Burdge GC & Calder PC (2006) Polyunsaturated fatty acid concentrations in young men and women consuming their habitual diets. *Br J Nutr* **96**, 93–99.
- Brossard N, Croset M, Pachioudi C, Riou JP, Tayot JL & Lagarde M (1996) Retroconversion and metabolism of [¹³C]22:6*n*-3 in humans and rats after intake of a single dose of [¹³C]22:6*n*-3-triacylglycerols. *Am J Clin Nutr* **64**, 577–586.
- Stark KD & Holub BJ (2004) Differential eicosapentaenoic acid elevations and altered cardiovascular disease risk factor responses after supplementation with docosahexaenoic acid in postmenopausal women receiving and not receiving hormone replacement therapy. *Am J Clin Nutr* **79**, 765–773.
- Rees D, Miles EA, Banerjee T, Wells SJ, Roynette CE, Wahle KW & Calder PC (2006) Dose-related effects of eicosapentaenoic acid on innate immune function in healthy humans: a comparison of young and older men. *Am J Clin Nutr* **83**, 331–342.
- Yep YL, Li D, Mann NJ, Bode O & Sinclair AJ (2002) Bread enriched with microencapsulated tuna oil increases plasma docosahexaenoic acid and total omega-3 fatty acids in humans. *Asia Pac J Clin Nutr* **11**, 285–291.
- Calder PC (2006) *n*-3 polyunsaturated fatty acids, inflammation, and inflammatory diseases. *Am J Clin Nutr* **83**, 1505S–1519S.
- Salem N Jr, Litman B, Kim HY & Gawrisch K (2001) Mechanisms of action of docosahexaenoic acid in the nervous system. *Lipids* **36**, 945–959.
- Kojima M, Wakai K, Tokudome S, *et al.* (2005) Serum levels of polyunsaturated fatty acids and risk of colorectal cancer: a prospective study. *Am J Epidemiol* **161**, 462–471.