

erage will only be determined through population-based surveys, and success can only be measured by good quality surveillance. However, this campaign does suggest that implementing an extended age range measles vaccination campaign on a nationwide scale during a complex emergency is possible.

Keywords: children; complex emergencies; measles; polio; surveys, population-based; surveillance; vaccination

Prehosp Disast Med 2002;17(s2):s4-5.

Modern Organization of the Work of a Tuberculosis Field Hospital for Rendering Assistance in Refugee Settlements

V.N. Preobrajensky, I.V. Sviridov

All-Russian Centre for Disaster Medicine (ARCDM), Moscow, Russia

In the world, practical experience of the work of tuberculosis (TB) field hospitals in emergencies is absent, and the development of main organizational regulations on rendering TB assistance is very important. In the Ingushetia Republic, the All-Russian Centre for Disaster Medicine (ARCDM), "Zaschita," established a 100-bed TB field hospital for early detection and prevention of TB in refugee settlements in accordance with WHO requirements. Its purpose was to treat patients in specialized TB institutions and to provide temporary hospitalization of seriously ill patients awaiting transport to medical institutions. A special TB team worked in the refugee settlements on early detection and prevention. A mobile fluorography unit was used and TB mycobacteria expectoration was collected from suspicious TB patients.

More than 12,000 patients were examined. Among them, according to complex examination, more than 1,500 different TB forms were revealed. By the structure of clinical TB forms the following was revealed: hematogenic disseminated TB in 23% of cases; infiltrative TB - 31%; fibrocavernous - 13%; extrapulmonary - 5%. Treatment was carried out with the complex agent Myrin-P or more than four tuberculosis agents. Drug resistance to TB mycobacterium was noticed in 5% to 7% of cases; to overcome it mycobutin was used.

An analysis of tuberculosis assistance in refugee settlements showed the possibility of using field hospitals and the necessity of integration with TB institutions of other regions.

Keywords: All-Russian Centre for Disaster Medicine; ARCDM; refugee settlements; TB; tuberculosis

Prehosp Disast Med 2002;17(s2):s5.

Intervention in Measles Epidemics

Rannveig Bremer Fjær, MD

Joint Medical Services, Norwegian Defence Forces, Sessvollmoen, Norway

Introduction: Measles epidemics are dreaded, especially in developing countries where the disease has a far more serious course, and where the case fatality rate (CFR) is estimated at 3-4%. Even though a vaccination coverage of more than 70% has been achieved, the death toll is estimated to be 800,000-900,000 children yearly.

Methods: During an 18-month stay in the Yemen Arab Republic as Chief Medical Officer at Norwegian Save the

Children's Mother and Child Clinic, four rural areas reported ongoing measles epidemics. A team of four members went to the remote and poorly accessible villages to vaccinate the healthy children in the surrounding villages, and to treat the sick.

Results: Approximately 3,000 children were vaccinated during these vaccination campaigns. Nearly none of the children had been vaccinated before. In one village, most of the children were sick, and during two days, 70 children were treated for pneumonia.

Conclusion: The most common cause of measles mortality is pneumonia, but few studies have documented higher mortality in children with malnutrition. Serious complications like precipitating kwashiorkor, deafness, and blindness are frequent. The observation that survival was increased and blindness reduced when giving vitamin A, resulted in the routine administration of vitamin A in the Expanded Programme of Immunisation (EPI+).

The Global Alert Response Network (GOARN), initiated by WHO in 2000, registers and monitors epidemics worldwide. The Network also provides interventions when indicated. Acute interventions with treatment of sick children and vaccination in surrounding villages, if accessible, is needed to reduce mortality and serious complications in measles epidemics.

Keywords: children; epidemics; Global Alert Response Network (GOARN); measles; pneumonia; response; vaccinations

Prehosp Disast Med 2002;17(s2):s5.

The Confluence of HIV/AIDS and Tuberculosis in the Philippines: A Disaster Waiting to Happen

Reynaldo Imperial

Padre Faura, Ermita, Manila, Philippines

The interaction of AIDS and tuberculosis (TB) is a matter of great concern for many developing countries such as the Philippines. Tuberculosis kills some 25,000 Filipinos each year. The Philippine Department of Health reports that 67.3% of Filipinos are infected with the disease. The World Health Organization has determined that the Philippines is one of the 10 countries in the world that has not been able to control the infection.

While the Philippines continue to record a low and slow HIV/AIDS prevalence rate with a total of 1,967 individuals said to be infected, most of the opportunistic infections that have been treated are infected with tuberculosis. At the San Lazaro Hospital (SLH), one of the facilities in which patients with AIDS are housed, 76.5% of the admitted patients also have TB. The UNAIDS, in its 2001 Report, claimed that incidence of TB has been growing steadily, and HIV infection is believed to be the culprit. This study was conducted to determine the ill-effects of the confluence of TB and HIV in the quality of life of people living with HIV/AIDS.

It sought to address the following objectives: (1) How has education and information dissemination among patients confined at the SLH changed their behaviour both to reduce the risk of being infected with or transmitting HIV and/or TB; and (2) Determine the level of patient compliance to Directly Observed Treatment Short Course (DOTS) for the management of TB.