

## Associations between the dietary intake of antioxidant nutrients and the risk of hip fracture in elderly Chinese: a case–control study

Li-li Sun<sup>1</sup>†, Bao-lin Li<sup>2</sup>†, Hai-li Xie<sup>1</sup>, Fan Fan<sup>1</sup>, Wei-zhong Yu<sup>2</sup>, Bao-hua Wu<sup>1,2</sup>, Wen-qiong Xue<sup>1</sup> and Yu-ming Chen<sup>1\*</sup>

<sup>1</sup>Guangdong Provincial Key Laboratory of Food, Nutrition and Health, Department of Medical Statistics and Epidemiology, School of Public Health, Sun Yat-sen University, Guangzhou 510080, People's Republic of China

<sup>2</sup>Guangzhou Orthopaedics Trauma Hospital, Guangzhou 510045, People's Republic of China

(Submitted 26 December 2013 – Final revision received 4 August 2014 – Accepted 6 August 2014 – First published online 7 October 2014)

### Abstract

The role of oxidative stress in skeletal health is unclear. The present study investigated whether a high dietary intake of antioxidant nutrients (vitamins C and E,  $\beta$ -carotene, animal-derived vitamin A, retinol equivalents, Zn and Se) is associated with a reduced risk of hip fracture in elderly Chinese. This 1:1 matched case–control study involved 726 elderly Chinese with hip fracture and 726 control subjects, recruited between June 2009 and May 2013. Face-to-face interviews were conducted to determine habitual dietary intakes of the above-mentioned seven nutrients based on a seventy-nine-item FFQ and information on various covariates, and an antioxidant score was calculated. After adjustment for potential covariates, dose-dependent inverse associations were observed between the dietary intake of vitamin C, vitamin E,  $\beta$ -carotene, and Se and antioxidant score and the risk of hip fracture ( $P$  for trend  $\leq 0.005$ ). The OR of hip fracture for the highest (*v.* lowest) quartile of intake were 0.39 (95% CI 0.28, 0.56) for vitamin C, 0.23 (95% CI 0.16, 0.33) for vitamin E, 0.51 (95% CI 0.36, 0.73) for  $\beta$ -carotene, 0.43 (95% CI 0.26, 0.70) for Se and 0.24 (95% CI 0.17, 0.36) for the antioxidant score. A moderate-to-high dietary intake of retinol equivalents in quartiles 2–4 (*v.* 1) was found to be associated with a lower risk of hip fracture (OR range: 0.51–0.63,  $P < 0.05$ ). No significant association was observed between dietary Zn or animal-derived vitamin A intake and hip fracture risk ( $P$  for trend  $> 0.20$ ). In conclusion, a higher dietary intake of vitamins C and E,  $\beta$ -carotene, and Se and a moderate-to-high dietary intake of retinol equivalents are associated with a lower risk of hip fracture in elderly Chinese.

**Key words:** Antioxidant nutrients: Dietary intakes: Hip fractures: Case–control studies: Chinese

Osteoporotic fractures contribute significantly to the societal disease burden<sup>(1)</sup>. Hip fracture is considered to be the most severe type of osteoporotic fracture due to the high morbidity, mortality and economic cost<sup>(2–4)</sup>. Therefore, prevention strategies for hip fracture are particularly important.

Some studies have suggested that oxidative stress plays an important role in bone resorption. Oxidative stress has been shown in basic studies to increase osteoclastic resorption by inducing the activation of NF- $\kappa$ B<sup>(5,6)</sup>, and 8-iso-PGF<sub>2</sub> $\alpha$  (a biomarker of oxidative stress) concentrations have been reported to be negatively associated with bone mineral density (BMD) in observational studies<sup>(7)</sup>.

Previous studies<sup>(8–10)</sup> have shown that higher consumption of fruit and vegetables is associated with higher BMD and bone mass and a reduced risk of fractures. Fruit and vegetables are major sources of antioxidants, such as vitamin C and  $\beta$ -carotene. Therefore, a high intake of fruit and vegetables may be a proxy for a high intake of antioxidants. Several

epidemiological studies have investigated the relationships between the retinol equivalent of animal-derived vitamin A and plant-derived  $\beta$ -carotene, vitamin C, vitamin E, Zn, and Se and BMD or fracture, but the findings have been inconsistent<sup>(11–19)</sup>. Many studies have found a positive association between antioxidant consumption and bone health<sup>(11,14,15)</sup>. In contrast, some studies have shown retinol-equivalent and animal-derived vitamin A to be associated with a low BMD or risk of fracture<sup>(16,17)</sup>; in the Women's Health Initiative Study, no significant association was found between retinol equivalents, vitamin C, vitamin E or Se and BMD<sup>(12)</sup>, and a longitudinal study also failed to establish a relationship between Zn and BMD<sup>(19)</sup>. Conflicting research findings suggest a potentially complex relationship between serum or plasma antioxidant concentrations and skeletal health<sup>(20–23)</sup>. The different study populations, study designs and sample sizes used may explain the discrepant observations in the studies reported thus far. The majority of previous studies have been conducted in

**Abbreviations:** BMD, bone mineral density; OC, oral contraceptives; USNBH, Utah Study of Nutrition and Bone Health.

\* **Corresponding author:** Dr Y.-m. Chen, fax +86 20 87330446, email chenym@mail.sysu.edu.cn

† These authors contributed equally to the study.

Western populations, and less is known about the association between the intake of antioxidants and skeletal health in Asian populations. The present study investigated the associations between the consumption of antioxidant nutrients, including vitamins C (mg/d) and E (mg/d), retinol equivalents ( $\mu\text{g}$  retinol equivalents/d), animal-derived vitamin A ( $\mu\text{g}$ /d),  $\beta$ -carotene ( $\mu\text{g}$ /d), Zn (mg/d) and Se ( $\mu\text{g}$ /d), and the risk of hip fracture in elderly Chinese.

## Participants and methods

### Study population

The present case-control study was conducted between June 2009 and May 2013 in Guangzhou, Guangdong Province, China. The total study group comprised 1452 recruits from four participating hospitals and communities. A detailed description of the study design has been published previously<sup>(10,24)</sup>.

In brief, case participants were newly diagnosed (within the previous 2 weeks) with hip fracture on the basis of X-ray examination. Patients with any of the following conditions were excluded from the study: (1) a high-energy or pathological fracture; (2) a change in dietary habits within the previous 5 years; (3) a chronic disease such as diabetes, CVD, cancer, cognitive disorder, liver cirrhosis, thyroid disorder, renal failure or chronic diarrhoea; (4) current use of exogenous oestrogen, thiazine, corticosteroid or other medications; (5) poor vision that might affect routine activities. Control subjects were individually matched (1:1) by sex and age ( $\pm 3$  years) from the same hospitals or the local communities in the same cities. The controls were recruited through local advertisements and subjects' referrals and interviewed within 3 months of the enrolment of the corresponding cases. The present study was conducted according to the guidelines laid down in the Declaration of Helsinki, and all procedures involving human subjects were approved by the Ethics Committee of the School of Public Health of Sun Yat-sen University (no. 3 in 2009). Written informed consent was obtained from all study participants.

### Data collection

Trained interviewers with relevant medical knowledge conducted face-to-face interviews with the study participants and also took anthropometric measurements. Participants' sociodemographic characteristics, lifestyle habits, medical history, family history, years since menopause for females, and dietary and supplement intakes were determined by means of structured questionnaires and recorded. Each interviewer completed an equal proportion of interviews between the case group and the control group.

### Calculation of dietary antioxidant intakes

Dietary intake information was collected by means of a modified semi-quantitative FFQ, which was used to determine the frequency ('never', 'per year', 'per month', 'per week' and 'per day') of consumption and amount of seventy-nine food

items consumed over the previous year, with a colour picture booklet used as a guide for portion sizes. The average daily intake of a given food item was multiplied by its nutrient content based on the Chinese Food Composition Table 2002<sup>(25)</sup>. For each study participant, daily intake of energy and nutrients was then calculated by totalling the values across all food items. Retinol equivalents included both animal-derived vitamin A and fruit- and vegetable-derived  $\beta$ -carotene, which can be converted to retinol, and the intake was expressed as retinol equivalents (1  $\mu\text{g}$  retinol equivalent = 1  $\mu\text{g}$  of retinol = 12  $\mu\text{g}$  of  $\beta$ -carotene). The validity and reproducibility of the FFQ have been reported elsewhere<sup>(26)</sup>. Correlation coefficients for energy-adjusted nutrients assessed from the questionnaires and six 3 d dietary records in 12 months were computed in the local population, and the correlation coefficients for vitamins A, C and E, retinol and carotene were found to be 0.32, 0.32, 0.25, 0.31 and 0.32, respectively<sup>(26)</sup>.

### Assessment of covariates

A face-to-face interview was conducted by a trained interviewer using a structured questionnaire to collect information on the following: age (year); sex (male/female); use of oral contraceptives (OC) and oestrogen (yes or no); education level (primary school or below, secondary school, and high school or above); occupation (full non-physical work, main non-physical work, and main physical labour, full physical labour, and others); household income (Yuan/month per person:  $\leq 500$ , 501–2000, 2001–3000, and  $> 3000$ ); family history of fracture (yes/no); smoking (yes/no); passive smoking (yes/no); alcohol drinking (yes/no); Ca supplement use (yes/no); multivitamin supplement use (yes/no); daily energy intake and intake of selected dietary nutrients (energy-adjusted protein and Ca and P). Subjects who smoked at least one cigarette per d or drank alcohol once a week for at least 6 months were defined as smokers or drinkers. Subjects who had been exposed to other people's tobacco smoking for at least 5 min daily in the previous 5 years were defined as passive smokers. Body height (cm) and weight (kg) were measured in the controls dressed in light clothing and without shoes and self-reported by the case participants. BMI ( $\text{kg}/\text{m}^2$ ) was then calculated. Daily physical activity (metabolic equivalent h/d) was estimated using a 24 h physical activity questionnaire containing nineteen items<sup>(27)</sup>.

### Statistical analyses

The characteristics of the case participants and control subjects were compared using the *t* test for continuous variables and Pearson's  $\chi^2$  test for categorical variables. The distribution of energy and nutrient intake data was normalised by log transformation. The dietary intakes of all nutrients were adjusted for total energy intake using a residual method<sup>(28)</sup>. An antioxidant score (ranging between 4 and 16) was calculated by summing the quartile points of each nutrient to evaluate the combined association of vitamins C and E,  $\beta$ -carotene, and Se. The participants were then categorised into quartiles (Q1–Q4) of intake according to the consumption of each energy-adjusted antioxidant or the score in control subjects, and the cut-offs

were applied for the classification of the case participants. The lowest quartile (Q1) was used as the reference.

As the matching of socio-economic factors (education level, household income and occupation) was not successful, both non-conditional and conditional regression methods were used and compared. Non-conditional logistic regression, as in the INTERHEART study<sup>(29)</sup>, was used to estimate the association between the intake of selected antioxidants and the risk of hip fracture as it is a more conservative approach.

Antioxidant nutrient intakes and antioxidant scores were analysed as continuous variables as well as categorical quartile variables to calculate OR and 95% CI. The lowest quartile was considered as the reference quartile in the categorical variable analysis. Trend tests were carried out by modelling the mean values of the intake of each antioxidant nutrient or antioxidant score in the control groups as a continuous variable. Models were adjusted for age, sex + drugs (defined as men and women using OC or oestrogen and women not using OC and oestrogen; model 1). Subsequent models were further adjusted for BMI, education level, occupation, household income, family history of fracture, smoking and alcohol drinking, passive smoking, Ca and multivitamin supplement use, physical activity, daily intakes of energy and energy-adjusted protein and Ca and P (model 2). All covariates were selected using the forward stepwise method.

Interaction analyses were conducted to explore whether the above associations varied in different sexes (male or female) and with the source of the control subjects (community or hospital), and the stratified results by sex (male or female) and the source of the control subjects are reported. Years since menopause, former use of oestrogen and use of OC were further adjusted by female sex in the multivariate analysis. Interaction between sex and the source of the control subjects and the antioxidants studied was tested using the likelihood ratio test. A two-sided *P* value <0.05 was considered significant. Considering type 1 error caused by multiple testing, *P* values were adjusted using Bonferroni correction. *P* adjusted=0.05/number of tests. All analyses were conducted using SPSS version 17.0 (SPSS, Inc.).

## Results

### Study participants

A total of 1402 potential case patients and 1215 potential control subjects from participating hospitals and local communities were screened, and 501 (35.7%) of the former and 355 (29.2%) of the latter were found to not meet the study criteria. An additional 175 case patients and 134 control subjects who did meet the eligibility criteria were excluded for the following reasons: difficulty in communicating (forty-eight case patients and twenty-one control subjects); unreasonable energy intake (nineteen case patients and sixteen control subjects; reasonable range: 3347–16 736 kJ/d (800–4000 kcal/d) for males and 2092–14 644 kJ/d (500–3500 kcal/d) for females); refusal to participate (108 case patients and twenty-four control subjects); a history of fracture (seventy-three control subjects). Thus, 726 case patients and 726 control subjects (542 recruited

from local communities and 184 recruited from hospitals) were included in the final analysis. Among the community-based controls, 6% were attendants or relatives of a patient from a non-hip fracture ward without any diseases related to the studied factors or bone health and 94% were recruited from the local communities of the same cities from where the cases came from.

The cases and controls had similar ages of 70 and 71 years in men and women, respectively. The mean values of dietary intakes of selected antioxidant nutrients were 474 and 476 µg retinol equivalents/d, 102 and 106 mg/d for vitamin C, 11 and 10 mg/d for vitamin E, 12.2 and 11.5 mg/d for Zn, and 48 and 48 µg/d for Se among men and women in the control group (Table 1). The dietary intake levels met 59 and 68% (retinol equivalents), 79 and 71% (vitamin E), and 96–106% (vitamin C, Zn and Se) of the values recommended by the Chinese Nutrition Society in 2000.

The characteristics of the participants stratified by case–control status are also given in Table 1. Overall, patients with hip fracture were more likely than control subjects to have a low BMI, to have low levels of education and household income, to engage in physical work, to be smokers, and to have consumed fewer multivitamin and Ca supplements. A trend of low physical activity and low OC and oestrogen use was observed in the female case patients.

### Associations between the dietary intake of the studied antioxidant nutrients and the risk of hip fracture

In model 1, a significant inverse association was observed between the dietary intake of each studied antioxidant nutrient and antioxidant score and the risk of hip fracture (*P* for trend <0.001–0.035; Table 2). After adjustment for age, sex, BMI, socio-economic factors, family history of fracture, lifestyle habits, Ca and multivitamin supplement use, physical activity and some dietary factors, the dose-dependent inverse associations between the dietary intake of vitamin C, vitamin E, β-carotene, and Se and antioxidant score and the risk of hip fracture were found to remain significant (*P* for trend ≤0.005). The OR for the highest *v.* lowest quartiles of vitamin C, vitamin E, β-carotene, and Se intake and antioxidant score were 0.39 (95% CI 0.28, 0.56), 0.23 (95% CI 0.16, 0.33), 0.51 (95% CI 0.36, 0.73), 0.43 (95% CI 0.26, 0.70) and 0.24 (95% CI 0.17, 0.36), respectively. A moderate-to-high dietary intake of retinol equivalents (animal- and plant-derived retinol combined) in quartiles 2–4 (*v.* quartile 1) was found to be associated with a reduced hip fracture risk (OR range: 0.51–0.63, all *P*<0.05). No significant association was found between the dietary intake of other antioxidant nutrients (animal-derived vitamin A and Zn) and the risk of hip fracture (*P* for trend 0.661 and 0.277, respectively). Similar results were obtained in the conditional logistic regression analyses (Table S1, available online) as well as when exposure variables were analysed as continuous variables (*P* range <0.001–0.040 for vitamin C, vitamin E, Se, and β-carotene and antioxidant score; Table S2, available online).

In the stratified analysis, no significant interactions were found between the dietary intake of the studied antioxidant

**Table 1.** Demographics, lifestyle characteristics and selected hip fracture risk factors of the study population (Mean values and standard deviations; number of participants and percentages)

	Men (n 177 pairs)					Women (n 549 pairs)				
	Controls		Cases		P	Controls		Cases		P
	Mean	SD	Mean	SD		Mean	SD	Mean	SD	
<b>Continuous variables*</b>										
Age (years)	69.8	7.1	70.2	7.1	0.560	71.2	6.8	71.3	6.9	0.822
BMI (kg/m <sup>2</sup> )	23.3	2.4	21.1	2.2	<0.001	23.1	3.2	21.6	3.9	<0.001
Years since menopause (years)	–	–	–	–	–	21.1	9.4	21.8	9.7	0.242
Physical activity† (MET-h/d)	72	43.4	64.3	45.5	0.103	73.2	54.6	62.4	42.7	<0.001
<b>Dietary intake</b>										
Energy (MJ/d)	6.41	1.62	6.37	1.53	0.836	6.03	1.51	5.74	1.60	0.003
Protein (g/d)	62.7	19.1	58.7	20.5	0.060	59.4	17.9	52.6	18.5	<0.001
Ca (mg/d)	468	208	357	140	<0.001	473	199	372	193	<0.001
Retinol equivalents (µg retinol equivalents/d)	474	227	401	219	<0.001	476	211	395	200	0.003
Vitamin A (animal-derived) (µg/d)	161	113	162	141	0.976	166	127	146	114	0.007
β-Carotene (µg/d)	3749	2243	2909	1535	<0.001	3728	1986	2993	1797	<0.001
Vitamin C (mg/d)	102	60	77	40	<0.001	106	57	82	47	<0.001
Vitamin E (mg/d)	11.10	4.87	8.12	4.78	<0.001	10.07	4.46	7.71	4.11	<0.001
Zn (mg/d)	12.17	3.44	11.98	3.64	0.613	11.46	3.08	10.65	3.38	<0.001
Se (µg/d)	48.3	21.2	43.5	18.3	0.024	47.7	19.4	40.8	17.5	<0.001
	<i>n</i>	%	<i>n</i>	%		<i>n</i>	%	<i>n</i>	%	
<b>Categorical variables‡</b>										
Education level					<0.001					<0.001
Primary school or below	54	30.7	93	52.2		247	45.1	359	66.4	
Secondary school	40	22.7	28	15.7		96	17.5	57	10.5	
High school or above	82	46.6	57	32		205	37.4	125	23.1	
Household income (Yuan/month per person)					0.001					<0.001
≤500	4	2.2	15	8.5		15	2.7	69	12.1	
501–2000	41	23.2	67	37.9		165	30.1	225	41.4	
2000–3000	83	46.9	62	35		241	44	178	32.7	
>3000	49	27.7	33	18.6		127	23.2	75	13.8	
Occupation§					0.028					<0.001
Full non-physical work	42	23.7	40	22.5		98	17.9	82	15.1	
Main non-physical work	62	35	37	20.8		137	25	89	16.4	
Main physical labour	26	14.7	35	19.7		63	11.5	74	13.6	
Full physical labour	41	23.2	58	32.6		200	36.5	268	49.3	
Others	6	3.4	8	4.5		50	9.1	31	5.7	
Family history of fractures	14	7.9	27	15.2	0.032	69	12.6	87	15.9	0.114
Smoker	68	38.4	88	49.4	0.036	11	2	28	5.1	0.005
Passive smoking¶	20	11.3	50	28.1	<0.001	100	18.2	129	23.6	0.029
Alcohol drinker**	20	11.3	34	19.1	0.041	24	4.4	15	2.7	0.145
Ca supplement user	50	28.2	25	14	0.001	228	41.6	177	32.4	0.002
Multivitamin user	43	24.3	13	7.3	<0.001	137	25	50	9.1	<0.001
Oral contraceptive user	–	–	–	–	–	92	17	28	5.3	<0.001
Former oestrogen user	–	–	–	–	–	49	9	8	1.6	<0.001

MET, metabolic equivalent.

\* Evaluated by *t* tests.

† Physical activities included daily occupational activities, leisure-time activities and household chores, evaluated by MET-h/d.

‡ Evaluated by  $\chi^2$  tests.

§ Occupation was categorised into five levels on the basis of labour model.

|| Smoker was defined as having smoked at least one cigarette daily for at least 6 consecutive months.

¶ Passive smoking was defined as having been exposed to other people's tobacco smoking for at least 5 min daily in the previous 5 years.

\*\* Alcohol drinker was defined as having had alcoholic beverages (wine, beer or Chinese spirits) at least once a week for at least 6 consecutive months.

nutrients and sex or source of the control subjects (Table 3; *P* interactions > 0.004, 0.05/fourteen tests).

## Discussion

In the present study, a reduced risk of hip fracture was found to be associated with a high dietary intake of vitamins C and E, β-carotene, and Se and with a moderate dietary intake of

retinol equivalents. Similar inverse associations were found when antioxidant scores were analysed.

Many epidemiological studies have found that vitamins C and E have beneficial effects on skeletal health<sup>(11,13,14)</sup>. A nested case–control study of 1120 elderly Swedish women aged 40–76 years included in the Swedish Mammography Cohort showed that a low intake of vitamins C and E increased the risk of hip fracture in current smokers after adjustment for

**Table 2.** Risk of hip fracture for quartiles (Q) of antioxidant intake in the study population (Number of cases and controls; odds ratios and 95 % confidence intervals)

Variables	Quartiles of dietary energy-adjusted intake				P for trend*
	Q1 (lowest)	Q2	Q3	Q4	
<b>Retinol equivalents</b>					
Intake† (µg retinol equivalents/d)					
Male	216	356	459	678	
Female	226	354	465	673	
Cases (n)	324	134	147	120	
Controls (n)	181	182	181	181	
OR I‡	1.00	0.41	0.46	0.37	<0.001
95 % CI		0.31, 0.55	0.34, 0.61	0.28, 0.50	
OR II§	1.00	0.51	0.61	0.63	0.050
95 % CI		0.36, 0.72	0.43, 0.88	0.42, 0.98	
<b>β-Carotene</b>					
Intake (µg/d)					
Male	1882	2931	3900	5954	
Female	1622	2648	3987	6281	
Cases (n)	287	170	145	123	
Controls (n)	181	182	181	181	
OR I	1.00	0.59	0.50	0.43	<0.001
95 % CI		0.44, 0.78	0.38, 0.67	0.32, 0.57	
OR II	1.00	0.68	0.57	0.51	<0.001
95 % CI		0.49, 0.94	0.41, 0.79	0.36, 0.73	
<b>Vitamin A (animal-derived)</b>					
Intake (µg/d)					
Male	64	108	159	308	
Female	64	111	159	303	
Cases (n)	212	204	150	159	
Controls (n)	181	182	181	181	
OR I	1.00	0.95	0.71	0.75	0.035
95 % CI		0.72, 1.26	0.53, 0.96	0.56, 1.00	
OR II	1.00	1.24	1.08	1.14	0.661
95 % CI		0.89, 1.73	0.75, 1.56	0.80, 1.63	
<b>Vitamin C</b>					
Intake (mg/d)					
Male	55	84	110	167	
Female	49	77	110	171	
Cases (n)	311	165	138	111	
Controls (n)	181	182	181	181	
OR I	1.00	0.51	0.45	0.35	<0.001
95 % CI		0.39, 0.68	0.34, 0.59	0.26, 0.47	
OR II	1.00	0.52	0.55	0.39	<0.001
95 % CI		0.38, 0.72	0.39, 0.77	0.28, 0.56	
<b>Vitamin E</b>					
Intake (mg/d)					
Male	5.99	8.23	10.36	14.48	
Female	6.37	9.17	11.58	16.74	
Cases (n)	388	144	118	75	
Controls (n)	181	182	181	181	
OR I	1.00	0.38	0.31	0.19	<0.001
95 % CI		0.29, 0.50	0.23, 0.41	0.14, 0.27	
OR II	1.00	0.46	0.38	0.23	<0.001
95 % CI		0.34, 0.63	0.28, 0.53	0.16, 0.33	
<b>Zn</b>					
Intake (mg/d)					
Male	9.18	10.43	11.52	13.86	
Female	9.74	11.19	12.12	14.55	
Cases (n)	257	160	166	142	
Controls (n)	181	182	181	181	
OR I	1.00	0.62	0.65	0.54	<0.001
95 % CI		0.47, 0.83	0.49, 0.86	0.41, 0.73	
OR II	1.00	0.83	0.99	0.76	0.277
95 % CI		0.60, 1.15	0.70, 1.39	0.53, 1.10	
<b>Se</b>					
Intake (µg/d)					
Male	31.6	40.8	47.9	60.5	
Female	31.3	39.5	47.7	63.9	
Cases (n)	267	196	175	87	
Controls (n)	181	182	181	181	

**Table 2.** *Continued*

Variables	Quartiles of dietary energy-adjusted intake				P for trend*
	Q1 (lowest)	Q2	Q3	Q4	
OR I	1.00	0.72	0.65	0.33	< 0.001
95% CI		0.55, 0.95	0.49, 0.86	0.24, 0.45	
OR II	1.00	0.87	0.95	0.43	0.005
95% CI		0.62, 1.22	0.63, 1.41	0.26, 0.70	
Antioxidant score					
Score	5	8	11	14	
OR I	1.00	0.48	0.32	0.22	< 0.001
95% CI		0.36, 0.64	0.23, 0.43	0.16, 0.31	
OR II	1.00	0.47	0.36	0.24	< 0.001
95% CI		0.34, 0.65	0.26, 0.51	0.17, 0.36	

\* Significant levels:  $P < 0.006$ , adjusted using Bonferroni correction.

† Mean intake of vitamin A in male/female controls, which was adjusted for daily energy intake using the residual method, and the mean of daily energy intake was 5669 kJ (1355 kcal) for males and 5347 kJ (1278 kcal) for females.

‡ OR I: from unconditional logistic model adjusted for age and sex + drugs (defined as men and women using oral contraceptives (OC) or oestrogen and women not using OC and oestrogen).

§ OR II: from unconditional logistic model adjusted for age; sex + drugs; BMI; educational level; occupation; household income; family history of fracture; smoking; passive smoking; alcohol drinking; Ca supplement use; multivitamin supplement use; physical activity; daily energy intake; and dietary intake of selected nutrients (protein and Ca and P; energy-adjusted), and all covariates were selected using the forward stepwise method.

age, weight and other osteoporosis risk factors<sup>(13)</sup>. Similar results were obtained in the Utah Study of Nutrition and Bone Health (USNBH), which examined the risk of hip fracture in 2564 Americans<sup>(11)</sup>, in the Framingham Osteoporosis Study of 4-year bone loss<sup>(14)</sup>, in an interventional study of BMD<sup>(30)</sup> and in a cross-sectional study<sup>(22)</sup>. Consistent with the results of these studies, we found a significant inverse association between the increased intake of vitamins C and E and the risk of hip fracture. However, the protective effect exerted by vitamins C and E against hip fracture and BMD was not observed in the Women's Health Initiative Study<sup>(12)</sup>, in the fracture intervention trial study<sup>(31)</sup>, or in a case-control study of 329 American women after adjustment for important covariates<sup>(32)</sup>. The reasons for the between-study differences remain unclear. Differences in the study designs and in the methods used for dietary intake assessment, as well as the varied populations, might in part explain the discrepancies. Vitamins C and E are important dietary antioxidants. They might improve bone health by scavenging free radicals<sup>(33)</sup>, which have been found to be involved in rodent bone metabolism and to enhance bone resorption<sup>(34)</sup>. In addition, there is much evidence suggesting that vitamins C and E play a role in the formation of collagen matrices<sup>(35,36)</sup>. Therefore, they are needed for normal bone development.

Our finding of an inverse association between the dietary intake of  $\beta$ -carotene and the risk of hip fracture is congruent with previously reported findings<sup>(11,37)</sup>. In the USNBH, Zhang *et al.*<sup>(11)</sup> observed an inverse association between the intake of  $\beta$ -carotene and the risk of osteoporotic hip fracture in 2564 Americans aged  $\geq 50$  years. Similar favourable effects of dietary  $\beta$ -carotene<sup>(37)</sup>, serum  $\beta$ -carotene concentrations<sup>(38)</sup> and plasma  $\beta$ -carotene concentrations<sup>(21)</sup> on BMD or BMD changes have been found. Furthermore, a cross-sectional study showed dietary  $\beta$ -carotene intake to be inversely correlated with the excretion of deoxypyridinoline (a marker of bone resorption)<sup>(39)</sup>. However, neither dietary intake nor the serum concentration of  $\beta$ -carotene was found to be associated with

hip fracture risk or bone loss in the Nurses' Health Study involving 72 337 postmenopausal women<sup>(17)</sup>, in the Aberdeen Prospective Osteoporosis Screening Study<sup>(19)</sup>, in the Uppsala Longitudinal Study of Adult Men<sup>(20)</sup>, in the Framingham Osteoporosis Study<sup>(40)</sup> or in the Swedish Mammography Cohort<sup>(13)</sup>. Vitamin C and many other phytochemicals coexist with  $\beta$ -carotene in foods. Although the positive association between  $\beta$ -carotene and reduced hip fracture risk might be attributed in part to its antioxidant effects<sup>(33)</sup>, many studies have shown phytochemicals such as lycopene to have positive effects on hip fracture risk<sup>(40)</sup> and BMD<sup>(21)</sup>. Further studies are needed to clarify the independent association of  $\beta$ -carotene with hip fracture risk by well adjusting for the coexisting phytochemicals.

Studies on the effect of retinol-equivalent intake on skeletal health have yielded inconsistent results. A population-based cohort study of 1526 American women aged  $\geq 55$  years showed an inverse U-shaped association between animal-derived vitamin A and BMD, particularly at the femoral neck<sup>(18)</sup>. Opatowsky & Bilezikian<sup>(41)</sup> also described a U-shaped association between serum vitamin A concentrations and hip fracture risk. Many studies have shown that retinol-equivalent and animal-derived vitamin A in high doses or high serum vitamin A concentrations accelerate bone loss and increase fracture risk<sup>(16,17,20)</sup>. Consistent with these findings, we found that a moderate intake of retinol equivalents had the strongest positive effect on hip fracture risk. However, we did not find a deleterious effect for the highest quartile, possibly because of a relatively low intake of retinol equivalents in the present study population in comparison with that in other populations<sup>(17,19)</sup>, and the highest intake was observed for the plant-derived retinol equivalents. Nevertheless, several studies have failed to establish a relationship between vitamin A and skeletal health<sup>(21,42)</sup>. Several biological mechanisms might explain the potential U-shaped association. Vitamin A deficiency has been shown to increase both osteoclastic and osteoblastic activities, resulting in abnormal bone growth in animals<sup>(43)</sup>. On the other hand, there is much evidence in rodents showing that excessive vitamin A or synthetic retinoid

**Table 3.** Risk of hip fracture for quartiles (Q) of vitamin and mineral intakes stratified by sex and source of controls in the study population†‡ (Odds ratios and 95 % confidence intervals)

	Quartiles of dietary energy-adjusted intake								<i>P</i> for trend	<i>P</i> for interaction*
	Q1	Q2		Q3		Q4 (highest)				
	OR	OR	95 % CI	OR	95 % CI	OR	95 % CI			
<b>Retinol equivalent</b>										
Sex										0.787
Male	1.00	0.91	0.39, 2.14	2.01	0.87, 4.68	1.11	0.42, 2.89	0.574		
Female	1.00	0.51	0.34, 0.75	0.55	0.36, 0.83	0.66	0.42, 1.04	0.194		
Source of controls										0.690
Hospital	1.00	0.34	0.18, 0.65	0.3	0.15, 0.6	0.35	0.18, 0.72	0.001		
Community	1.00	0.60	0.40, 0.90	0.71	0.47, 1.09	0.76	0.48, 1.23	0.296		
<b>β-Carotene</b>										
Sex										0.941
Male	1.00	1.67	0.74, 3.76	1.85	0.80, 4.27	0.41	0.16, 1.06	0.039		
Female	1.00	0.49	0.33, 0.72	0.36	0.24, 0.53	0.43	0.29, 0.64	<0.001		
Source of controls										0.946
Hospital	1.00	0.66	0.35, 1.24	0.57	0.29, 1.11	0.43	0.21, 0.9	0.016		
Community	1.00	0.58	0.39, 0.86	0.47	0.31, 0.71	0.41	0.27, 0.62	<0.001		
<b>Animal-derived vitamin A</b>										
Sex										0.372
Male	1.00	1.89	0.86, 4.15	3.22	1.32, 7.87	2.23	0.93, 5.30	0.148		
Female	1.00	1.06	0.72, 1.55	0.93	0.62, 1.40	1.06	0.71, 1.58	0.807		
Source of controls										0.200
Hospital	1.00	0.88	0.44, 1.75	0.64	0.34, 1.23	0.46	0.25, 0.85	0.008		
Community	1.00	1.3	0.88, 1.92	1.21	0.78, 1.88	1.48	0.95, 2.31	0.124		
<b>Vitamin C</b>										
Sex										0.605
Male	1.00	0.69	0.31, 1.53	1.01	0.44, 2.32	0.25	0.10, 0.61	0.004		
Female	1.00	0.46	0.32, 0.67	0.42	0.28, 0.62	0.38	0.25, 0.56	<0.001		
Source of controls										0.354
Hospital	1.00	0.55	0.29, 1.03	0.72	0.36, 1.42	0.53	0.24, 1.19	0.076		
Community	1.00	0.49	0.33, 0.73	0.47	0.31, 0.7	0.32	0.21, 0.49	<0.001		
<b>Vitamin E</b>										
Sex										0.213
Male	1.00	0.53	0.26, 1.07	0.13	0.06, 0.31	0.19	0.08, 0.45	<0.001		
Female	1.00	0.50	0.34, 0.73	0.55	0.37, 0.81	0.30	0.19, 0.46	<0.001		
Source of controls										0.029
Hospital	1.00	0.4	0.23, 0.70	0.42	0.19, 0.92	0.52	0.24, 1.11	0.006		
Community	1.00	0.41	0.28, 0.62	0.29	0.2, 0.43	0.15	0.1, 0.23	<0.001		
<b>Zn</b>										
Sex										0.263
Male	1.00	2.09	0.96, 4.54	1.40	0.62, 3.19	1.80	0.74, 4.35	0.303		
Female	1.00	0.73	0.50, 1.07	0.98	0.66, 1.44	0.80	0.53, 1.23	0.443		
Source of controls										0.991
Hospital	1.00	0.53	0.3, 0.94	1.08	0.57, 2.07	0.78	0.34, 1.79	0.265		
Community	1.00	0.88	0.58, 1.32	0.91	0.60, 1.38	0.75	0.48, 1.16	0.240		
<b>Se</b>										
Sex										0.737
Male	1.00	0.57	0.27, 1.21	0.44	0.20, 0.98	0.19	0.07, 0.54	0.002		
Female	1.00	1.01	0.69, 1.47	1.19	0.80, 1.78	0.64	0.41, 0.99	0.001		
Source of controls										0.255
Hospital	1.00	0.73	0.41, 1.32	0.89	0.46, 1.73	0.42	0.2, 0.88	0.052		
Community	1.00	0.87	0.57, 1.31	0.88	0.54, 1.41	0.42	0.23, 0.76	0.013		

\* Significant level:  $P > 0.004$  (= 0.05/forteen tests), adjusted using Bonferroni correction.

† Study size: male, 177 pairs; female, 549 pairs; hospital controls, 184 pairs; community controls, 542 pairs.

‡ OR from multivariate unconditional logistic regression models. The following covariates were adjusted for: age; BMI; educational level; occupation; household income; family history of fracture; smoking; passive smoking; alcohol drinking; Ca supplement use; multivitamin supplement use; physical activity; daily energy intake; dietary intake of selected nutrients (protein and Ca and P; energy-adjusted). For women, years since menopause, oral contraceptive use, and former use of oestrogen were further adjusted for using the stepwise forward method.

is associated with increased osteoclastic bone resorption<sup>(44,45)</sup>. These findings suggest that retinol equivalents are required for skeletal growth, but hypervitaminosis A may have a deleterious effect on skeletal health.

Se supplementation can reinforce endogenous antioxidative systems<sup>(46)</sup>; thus, it may improve bone health by defending

against oxidative stress. We found Se to be inversely associated with the risk of hip fracture, and this association has also been observed in other studies<sup>(11)</sup>. However, no significant association was found in the Swedish Mammography Cohort<sup>(13)</sup>. The null association might be attributed in part to the pronounced errors in the assessment of Se intake because

of the variation in the Se content of foods between different countries and different regions<sup>(47)</sup>.

To assess the combined association of vitamins C and E,  $\beta$ -carotene, and Se, we further examined the association of the antioxidant score and the risk of hip fracture by summing the quartile points of each nutrient. A more significant inverse association was observed when compared with that observed for individual nutrients, except vitamin E, suggesting that the combination of vitamin C,  $\beta$ -carotene and Se might be more efficient than individual nutrients. We also assessed whether the inverse association was modified by sex and the source of the control subjects. No significant interaction was found after adjusting for the number of multiple tests, indicating similar associations across the subgroups by these variables. However, we had insufficient power to detect the interactions between some studied nutrients and sex or the source of the control subjects with regard to the risk of hip fracture. For interactions between antioxidant nutrients and sex, we had a power <50% for retinol equivalents,  $\beta$ -carotene and vitamin C, but >90% for the remaining antioxidant nutrients to detect an OR of 0.7, with an  $\alpha$  error of 0.004, assuming an ordinal trend across quartiles of intake. For interactions between antioxidant nutrients and the source of the control subjects, we had a power <30% for retinol equivalents and  $\beta$ -carotene and >90% for the remaining exposures<sup>(48)</sup>.

The present study has several limitations. In a case–control study, the time period between the exposure and the outcome is unclear. Nevertheless, this factor might have been mitigated in the present study because only new cases were selected; potential case patients and control subjects with a chronic disease that could have altered dietary habits or nutritional factors were excluded, and adults maintain relatively stable long-term dietary habits<sup>(49)</sup>. Moreover, recall bias might affect the results. We attempted to minimise recall bias through face-to-face interviews and by visual aids for the assessment of portion sizes. We also controlled interviewer bias by having each researcher interview a similar proportion of cases and controls. Another limitation was the assumptions made in the calculation of antioxidant values for food items. Antioxidant concentrations varied across foods that were combined in one item. When this occurred, we assigned the mean value of the contributing foods. In addition, limited information was collected about the dietary intake of oil, which is a rich source of vitamin E. Measurement errors could have resulted in the misclassification of the participants. However, such errors might have reduced rather than strengthened the observed association. In addition, the calculation of antioxidant score from the intake of antioxidant nutrients could lead to misclassification, and we assumed all the included antioxidants to have contributed equally to the association. Finally, as the study was hospital-based, selection bias could not be excluded completely, despite the fact that the case patients were recruited from various types of hospitals and controls were mainly recruited from the local communities.

In conclusion, the results of the present study suggest that a high dietary intake of vitamin C, vitamin E,  $\beta$ -carotene, and Se and a moderate-to-high dietary intake of retinol equivalents may protect against hip fracture in elderly Chinese.

## Supplementary material

To view supplementary material for this article, please visit <http://dx.doi.org/10.1017/S0007114514002773>

## Acknowledgements

The authors are grateful to Dr Wei-fu Ouyang and Sulan Tu for helping with data collection and the doctors and nurses in the above-mentioned hospitals for facilitating both the recruitment of participants and the interviews.

The present study was supported by the National Natural Science Foundation of China (Y.-m. C., grant no. 81072299, 81273049 and 30872100). The sponsor had no role in the design and analysis of the study or in the writing of this article.

The authors' contributions are as follows: Y.-m. C. conceived and designed the study and critically revised the manuscript; L.-l. S. analysed the data and wrote the article; B.-l. L., H.-l. X., F. F., W.-z. Y., B.-h. W. and W.-q. X. carried out the study and data cleansing and wrote the article.

None of the authors has any conflicts of interest to declare.

## References

1. Johnell O & Kanis JA (2006) An estimate of the worldwide prevalence and disability associated with osteoporotic fractures. *Osteoporos Int* **17**, 1726–1733.
2. Johnell O & Kanis J (2005) Epidemiology of osteoporotic fractures. *Osteoporos Int* **16**, S3–S7.
3. Johnell O & Kanis JA (2004) An estimate of the worldwide prevalence, mortality and disability associated with hip fracture. *Osteoporos Int* **15**, 897–902.
4. Cooper C, Campion G & Melton LJ III (1992) Hip fractures in the elderly: a world-wide projection. *Osteoporos Int* **2**, 285–289.
5. Hall TJ, Schaeublin M, Jeker H, *et al.* (1995) The role of reactive oxygen intermediates in osteoclastic bone resorption. *Biochem Biophys Res Commun* **207**, 280–287.
6. Franzoso G, Carlson L, Xing L, *et al.* (1997) Requirement for NF-kappa B in osteoclast and B-cell development. *Genes Dev* **11**, 3482–3496.
7. Basu S, Michaelsson K, Olofsson H, *et al.* (2001) Association between oxidative stress and bone mineral density. *Biochem Biophys Res Commun* **288**, 275–279.
8. Tucker KL, Hannan MT, Chen H, *et al.* (1999) Potassium, magnesium, and fruit and vegetable intakes are associated with greater bone mineral density in elderly men and women. *Am J Clin Nutr* **69**, 727–736.
9. Chen YM, Ho SC & Woo JLF (2006) Greater fruit and vegetable intake is associated with increased bone mass among postmenopausal Chinese women. *Br J Nutr* **96**, 745–751.
10. Xie HL, Wu BH, Xue WQ, *et al.* (2013) Greater intake of fruit and vegetables is associated with a lower risk of osteoporotic hip fractures in elderly Chinese: a 1:1 matched case–control study. *Osteoporos Int* **24**, 2827–2836.
11. Zhang JJ, Munger RG, West NA, *et al.* (2006) Antioxidant intake and risk of osteoporotic hip fracture in Utah: an effect modified by smoking status. *Am J Epidemiol* **163**, 9–17.
12. Wolf RL, Cauley JA, Pettinger M, *et al.* (2005) Lack of a relation between vitamin and mineral antioxidants and



- bone mineral density: results from the Women's Health Initiative. *Am J Clin Nutr* **82**, 581–588.
13. Melhus H, Michaelsson K, Holmberg L, *et al.* (1999) Smoking, antioxidant vitamins, and the risk of hip fracture. *J Bone Miner Res* **14**, 129–135.
  14. Sahni S, Hannan MT, Gagnon D, *et al.* (2008) High vitamin C intake is associated with lower 4-year bone loss in elderly men. *J Nutr* **138**, 1931–1938.
  15. Hall SL & Greendale GA (1998) The relation of dietary vitamin C intake to bone mineral density: results from the PEPI study. *Calcif Tissue Int* **63**, 183–189.
  16. Melhus H, Michaelsson K, Kindmark A, *et al.* (1998) Excessive dietary intake of vitamin A is associated with reduced bone mineral density and increased risk for hip fracture. *Ann Intern Med* **129**, 770–778.
  17. Feskanich D, Singh V, Willett WC, *et al.* (2002) Vitamin A intake and hip fractures among postmenopausal women. *JAMA* **287**, 47–54.
  18. Promislow JHE, Goodman-Gruen D, Slymen DJ, *et al.* (2002) Retinol intake and bone mineral density in the elderly: the Rancho Bernardo Study. *J Bone Miner Res* **17**, 1349–1358.
  19. Macdonald HM, New SA, Golden MH, *et al.* (2004) Nutritional associations with bone loss during the menopausal transition: evidence of a beneficial effect of calcium, alcohol, and fruit and vegetable nutrients and of a detrimental effect of fatty acids. *Am J Clin Nutr* **79**, 155–165.
  20. Michaelsson K, Lithell H, Vessby B, *et al.* (2003) Serum retinol levels and the risk of fracture. *N Engl J Med* **348**, 287–294.
  21. Maggio D, Polidori MC, Barabani M, *et al.* (2006) Low levels of carotenoids and retinol in involuntarily osteoporosis. *Bone* **38**, 244–248.
  22. Maggio D, Barabani M, Pierandrei M, *et al.* (2003) Marked decrease in plasma antioxidants in aged osteoporotic women: results of a cross-sectional study. *J Clin Endocrinol Metab* **88**, 1523–1527.
  23. Simon JA & Hudes ES (2001) Relation of ascorbic acid to bone mineral density and self-reported fractures among US adults. *Am J Epidemiol* **154**, 427–433.
  24. Fan F, Xue WQ, Wu BH, *et al.* (2013) Higher fish intake is associated with a lower risk of hip fractures in Chinese men and women: a matched case–control study. *PLOS ONE* **8**, e56849.
  25. Yang YX, Wang GY & Pan XC (2002) *China Food Composition Table*. Beijing: Peking University Medical Press.
  26. Zhang C-X & Ho SC (2009) Validity and reproducibility of a food frequency questionnaire among Chinese women in Guangdong province. *Asia Pac J Clin Nutr* **18**, 240–250.
  27. Wang P, Chen YM, He LP, *et al.* (2012) Association of natural intake of dietary plant sterols with carotid intima–media thickness and blood lipids in Chinese adults: a cross-section study. *PLOS ONE* **7**, e32736.
  28. Willett WC (1998) Implications of total energy intake for epidemiologic analysis. In *Nutritional Epidemiology*, 2nd ed., pp. 273–301 [WC Willett, editor]. New York, NY: Oxford University Press.
  29. Yusuf S, Hawken S, Ounpuu S, *et al.* (2004) Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case–control study. *Lancet* **364**, 937–952.
  30. Chuin A, Labonte M, Tessier D, *et al.* (2009) Effect of antioxidants combined to resistance training on BMD in elderly women: a pilot study. *Osteoporos Int* **20**, 1253–1258.
  31. Leveille SG, LaCroix AZ, Koepsell TD, *et al.* (1997) Dietary vitamin C and bone mineral density in postmenopausal women in Washington State, USA. *J Epidemiol Community Health* **51**, 479–485.
  32. Nieves JW, Grisso JA & Kelsey JL (1992) A case–control study of hip fracture: evaluation of selected dietary variables and teenage physical activity. *Osteoporos Int* **2**, 122–127.
  33. Fairfield KM & Fletcher RH (2002) Vitamins for chronic disease prevention in adults: scientific review. *JAMA* **287**, 3116–3126.
  34. Garrett IR, Boyce BF, Oreffo RO, *et al.* (1990) Oxygen-derived free radicals stimulate osteoclastic bone resorption in rodent bone *in vitro* and *in vivo*. *J Clin Invest* **85**, 632–639.
  35. Schwartz ER (1979) Effect of vitamins C and E on sulfated proteoglycan metabolism and sulfatase and phosphatase activities in organ cultures of human cartilage. *Calcif Tissue Int* **28**, 201–208.
  36. Kipp DE, McElvain M, Kimmel DB, *et al.* (1996) Scurvy results in decreased collagen synthesis and bone density in the guinea pig animal model. *Bone* **18**, 281–288.
  37. Sahni S, Hannan MT, Blumberg J, *et al.* (2009) Inverse association of carotenoid intakes with 4-y change in bone mineral density in elderly men and women: the Framingham Osteoporosis Study. *Am J Clin Nutr* **89**, 416–424.
  38. Sugiura M, Nakamura M, Ogawa K, *et al.* (2008) Bone mineral density in post-menopausal female subjects is associated with serum antioxidant carotenoids. *Osteoporos Int* **19**, 211–219.
  39. New SA, Robins SP, Campbell MK, *et al.* (2000) Dietary influences on bone mass and bone metabolism: further evidence of a positive link between fruit and vegetable consumption and bone health? *Am J Clin Nutr* **71**, 142–151.
  40. Sahni S, Hannan MT, Blumberg J, *et al.* (2009) Protective effect of total carotenoid and lycopene intake on the risk of hip fracture: a 17-year follow-up from the Framingham Osteoporosis Study. *J Bone Miner Res* **24**, 1086–1094.
  41. Opatowsky AR & Bilezikian JP (2004) Serum vitamin A concentration and the risk of hip fracture among women 50 to 74 years old in the United States: a prospective analysis of the NHANES I follow-up study. *Am J Med* **117**, 169–174.
  42. Barker ME, McCloskey E, Saha S, *et al.* (2005) Serum retinoids and beta-carotene as predictors of hip and other fractures in elderly women. *J Bone Miner Res* **20**, 913–920.
  43. Mellanby E (1941) Skeletal changes affecting the nervous system produced in young dogs by diets deficient in vitamin A. *J Physiol* **99**, 467–486.
  44. Frankel TL, Seshadri MS, McDowall DB, *et al.* (1986) Hypervitaminosis A and calcium-regulating hormones in the rat. *J Nutr* **116**, 578–587.
  45. Kneissel M, Studer A, Cortesi R, *et al.* (2005) Retinoid-induced bone thinning is caused by subperiosteal osteoclast activity in adult rodents. *Bone* **36**, 202–214.
  46. Tapiero H, Townsend DM & Tew KD (2003) The antioxidant role of selenium and seleno-compounds. *Biomed Pharmacother* **57**, 134–144.
  47. Monsen ER (2000) Dietary reference intakes for the antioxidant nutrients: vitamin C, vitamin E, selenium, and carotenoids. *J Am Diet Assoc* **100**, 637–640.
  48. Foppa I & Spiegelman D (1997) Power and sample size calculations for case–control studies of gene–environment interactions with a polytomous exposure variable. *Am J Epidemiol* **146**, 596–604.
  49. MacDonald HM, New SA & Reid DM (2005) Longitudinal changes in dietary intake in Scottish women around the menopause: changes in dietary pattern result in minor changes in nutrient intake. *Public Health Nutr* **8**, 409–416.