

Breakfast habits affect overall nutrient profiles in adolescents

C Matthys^{1,*}, S De Henauw^{1,2}, M Bellemans¹, M De Maeyer¹ and G De Backer¹

¹Department of Public Health, De Pintelaan 185, Ghent University, B-9000 Ghent, Belgium: ²Department of Health Sciences, Vesalius – Hogeschool Gent, Ghent, Belgium

Submitted 11 July 2005; Accepted 4 July 2006

Abstract

Objective: To describe breakfast consumption patterns, on a nutrient and food item level, in Belgian adolescents.

Design: A 7-day estimated food record was administered in a cross-sectional survey.

Setting: Secondary schools in Ghent, Belgium.

Subjects: A total of 341 adolescents (13–18 years old), multistage clustered sampling.

Results: The energy contribution of breakfast to daily energy intake was on average 15.7% in boys and 14.9% in girls. Significantly more overweight girls and significantly more girls following vocational training were categorised as eating a low-quality breakfast. In boys, the energy contribution of polysaccharides was significantly higher in consumers of good-quality breakfasts. The intake of all selected micronutrients was significantly higher in consumers of good-quality breakfasts. In girls, the total energy intake and the proportional intake of proteins and polysaccharides were significantly higher in consumers of good-quality breakfasts, while the proportional contribution of total fat, monounsaturated and polyunsaturated fatty acids was significantly lower in these girls. The intake of all micronutrients was significantly higher in girls consuming a good-quality breakfast. In all adolescents, consumers of a good-quality breakfast had significantly higher intakes of bread, fruit, vegetables, milk and milk products, and fruit juice, while intake of soft drinks was significantly lower than in consumers of low-quality breakfasts.

Conclusions: Consumers of a good-quality breakfast had a better overall dietary pattern – on a nutrient and food group level – than consumers of a low-quality breakfast. A daily breakfast, including whole-grain products, fruit and (semi-) skimmed milk products or an alternative source of calcium, is recommended.

Keywords
Adolescents
Breakfast habits
Dietary intake
Nutritional survey

Breakfast is widely being promoted as essential for the nutritional well-being of children. Skipping breakfast is associated with health-compromising behaviours in adults and adolescents¹. Breakfast consumers tend to have higher intakes of micronutrients and lower intake of fat, and are more likely to have a better overall diet quality^{2,3}. The consumption of breakfast has been positively associated with enhanced cognitive and academic performance, psychosocial function and school attendance^{4,5}. However, other studies indicate that cognitive performance is relatively robust to short-term fasting in relatively well-nourished children⁶. Aranceta and co-workers stated that it is not clear to what extent breakfast contributes to a better cognitive performance in school. Learning is a complex process resulting from multiple interactions⁷. However, a recent review stated that breakfast may possibly benefit cognitive function, but the interpretation of the results can be complicated by confounding factors such as social and educational variables³.

It has been reported that routinely eating breakfast may lead to more regular eating habits and exercise patterns, healthy food choices and consistent energy intake, which when taken together contribute to a reduced body mass index (BMI)⁸. Eating a healthy breakfast is important to adolescents' health needs in general. Breakfast provides an ideal opportunity for adolescents to begin the day by eating bread, other cereals and fruit, which are all important elements of a healthy and balanced diet. The importance of breakfast consumption in relation to nutritional balance has been shown in different sub-populations^{7,9,10}. Despite the benefits of consumption of breakfast, it is the meal most often skipped by adolescents¹¹.

In the present report, breakfast consumption patterns, on the nutrient and food item level, in Belgian adolescents are described. The analyses reported herein must be situated within a broad context of a search for elements that may be helpful for developing strategies to implement the translation of nutrient dietary guidelines into food- and meal-based dietary guidelines.

*Corresponding author: Email Christophe.Matthys@UGent.be

Materials and methods

The data presented here were obtained from a cross-sectional dietary survey carried out in an adolescent population (males and females aged 13–18 years) in the region of Ghent (Belgium) in the spring of 1997. The sampling design and the methodology of the field work have been described in detail elsewhere¹². In brief, a random sample of 341 adolescents (129 boys and 212 girls) – selected from all educational levels in the Belgian secondary school system – completed a 7-day food record (consecutive) under rigorous conditions of quality control carried out by experienced dietitians.

A 7-day estimated food record method (semi-structured diary) was used to quantify food and nutrient intake. Information on the type (including brand names) and amount of food consumed was collected through an open entry format. Instructions for the completion of the diary and regular checks for quality and completeness of the diaries were carried out by experienced dietitians. In the diaries, days were truncated into six eating moments, namely breakfast, lunch, dinner and snacks, the latter divided into morning, afternoon and late-evening snacks.

Breakfast was defined as the first eating occasion involving a solid food or a beverage that occurred after waking.

Nutrient composition data used in the current study are those from the Belgian and the Dutch food composition tables^{13–15}. Calculation of nutrients was done by means of a nutritional software package developed by the Unilever company in The Netherlands¹⁶. Average nutrient and food intakes were calculated as the mean of the 7-day intake period. Food items were classified into different food groups according to the Dutch food composition table¹³.

For the sake of the present analyses, a concise characterisation of the breakfast habits at the level of the individual was undertaken. This characterisation included

both qualitative and quantitative aspects of breakfast. Qualitative aspects of breakfast were related to the presence or absence (in relevant amounts) of food items of three specific food groups (hereafter called ‘target food groups’): cereal products, dairy products and fruit/vegetables. Quantitative aspects of breakfast included both the frequency of having breakfast and the relative contribution of the target food groups to the total daily energy from breakfast.

Qualitative and quantitative aspects of breakfast were combined into a so-called ‘individual breakfast score’ on the basis of the following consecutive steps.

In a first step, every single breakfast from all diaries was characterised in a qualitative way. For that purpose, 14 different labels were developed on the basis of different combinations of food groups present in the breakfast and the amount of energy they represented. These 14 labels were subsequently recoded into five specific ‘individual breakfast labels’ (see Table 1).

In a second step, the ‘individual breakfast labels’ were summed up for all days included in the food diary, resulting in a potential overall range of sum-labels from a minimum of seven (individual breakfast label of one for all days in the food diary) up to a maximum of 35 (individual breakfast label of five for each day in the food diary). From this ‘sum-score’, a final ‘individual breakfast score’ was assigned to all 341 subjects in the study (see Table 1).

For this study, the five individual breakfast score categories for breakfast habits were dichotomised into a group who ‘never eats breakfast or eats a low-quality breakfast, i.e. not enough calories and/or not enough variation in foods’ (group 1, score 1–3) and a group who ‘eats a good-excellent-quality breakfast (nearly) every day’ (group 2, score 4–5).

The ratio of energy intake to basal metabolic rate (EI/BMR) was calculated to give an indication of the quality of reporting¹⁷.

Table 1 Individual breakfast label and individual breakfast score

Individual breakfast label		Individual breakfast score	
Number	Explanation	Number	Explanation
Label 1	No breakfast or very limited breakfast (< 400 kJ)	Score 1	Subjects who never have any breakfast of any significance (sum-score 7–10)
Label 2	Breakfast including only energy-containing beverages	Score 2	Subjects who usually do not have breakfast of any significance (sum-score 11–14)
Label 3	Breakfast including food items from one of the target food groups, potentially in combination with a non-dairy fat or protein source	Score 3	Subjects who either usually have breakfast of low nutritional value or only occasionally have breakfast of higher nutritional value (sum-score 15–21)
Label 4	Breakfast including food items from two of the target food groups, potentially in combination with a non-dairy fat or protein source	Score 4	Subjects who usually have breakfast of good/excellent nutritional value (sum-score 22–29)
Label 5	Breakfast including food items from the three target food groups, potentially in combination with a non-dairy fat or protein source	Score 5	Subjects who eat good-excellent-quality breakfast practically every day (sum-score 30–35)

Statistical analysis was done with the SPSS software version 12 (SPSS Inc.). Descriptive statistics used means and standard deviations for continuous data. Tests for normality were performed using a Kolmogorov–Smirnov test. Student's *t*-tests or Mann–Whitney *U*-tests were used to compare the means of the different groups. In order to search for potential confounders, the number of adolescents in different categories of BMI¹⁸ and education – ‘general’ education (mainly theoretical courses) and vocational training (based on practical skills) – in the different breakfast classifications were compared by use of a Fisher's exact test. A value of <0.05 was taken as the threshold for significance.

The study was approved by the Ethical Committee of the Ghent University Hospital.

Results

The energy contribution of breakfast to daily energy intake was on average 15.7% in boys and 14.9% in girls. A restricted number of boys and girls had a breakfast energy contribution of >25%, 9.7% in boys and 5.7% in girls, respectively. Figure 1 shows the distribution of the energy contribution of breakfast to the total daily energy intake in both boys and girls.

Overall, the individual breakfast score was <3 (never eat or usually do not eat breakfast). In boys, 13.2% had an individual breakfast score <3, in girls 16.9% had an individual breakfast score <3. In boys, ~18% ate a low-quality breakfast (score 3), ~56% ate a ‘moderate- to good-quality breakfast’ (score 4), while only 13% ate a full value breakfast (score 5). In girls, ~27% had an individual breakfast score of 3, ~45% had a score of 4 and only 10% had an individual breakfast score of 5.

Table 2 presents the number of adolescents in different categories of BMI and education, and mean age and mean EI/BMR according to the different breakfast categories. In both boys and girls, a higher number of subjects were identified as good-quality breakfast eaters. In boys, the age of low-quality breakfast eaters was significantly higher than that of the good-quality breakfast eaters. In girls, the mean EI/BMR ratio was significantly lower in low-quality breakfast consumers than in the subjects identified as good-quality breakfast consumers. In boys, no differences were found between breakfast categories according to BMI and educational training. Significantly more overweight girls and significantly more girls following a vocational training were categorised as low-quality breakfast consumers.

Table 3 shows the energy intake at breakfast, the energy contribution of macronutrients to the total energy supplied by breakfast and micronutrient intake at breakfast according to breakfast habits. In both boys and girls, the energy intake and the proportional contribution of proteins were significantly higher in subjects having a good-quality breakfast. Girls who consumed a good-quality breakfast had a significantly higher proportional intake of polysaccharides than the low-quality breakfast consumers. In both boys and girls, the intake at breakfast of the selected micronutrients was significantly higher in subjects consuming a good-quality breakfast.

In Table 4 total energy intake, the energy contribution of macronutrients to the total energy intake and micronutrient intake according to breakfast habits are presented. In boys, there was no significant difference in the total energy intake between the two kinds of breakfast consumers.




Fig. 1 Histograms showing the percentage energy from breakfast in boys and girls

Table 2 The age, ratio of EI/BMR and number of adolescents in different categories of BMI and education according to breakfast habit (low- or good-quality)

	Boys (n = 129)		P-value	Girls (n = 212)		P-value
	Low	Good		Low	Good	
n	40	89		94	118	
Age (years)*	16 (1)	15 (1)	0.013	16 (1)	16 (1)	0.482
EI/BMR*	1.49 (0.33)	1.59 (0.29)	0.130	1.27 (0.37)	1.43 (0.28)	<0.001
BMI†,‡,§			>0.999			0.029
Normal	38	83		73	105	
Overweight	2	6		19	11	
Education§			0.400			<0.001
General	33	79		25	76	
Vocational	7	10		69	42	

EI – energy intake; BMR – basal metabolic rate; BMI – body mass index.

Data for age and EI/BMR presented as mean (standard deviation).

* Mann–Whitney *U*-test.

† BMI categories according to Cole *et al.*¹⁸.

‡ Data of four girls are missing.

§ Fisher's exact test.

The energy contribution of polysaccharides was significantly higher in good-quality breakfast consumers. The intake of all selected micronutrients was significantly higher in good-quality breakfast consumers.

In girls, the total energy intake was significantly higher in good-quality breakfast consumers. The proportional intake of proteins and polysaccharides was significantly higher in female good-quality breakfast consumers, while the proportional contribution of total fat, monounsaturated and polyunsaturated fatty acids was significantly lower in these girls. The intake of all micronutrients was significantly higher in girls consuming good-quality breakfast. In girls, the relative micronutrient intake (expressed as mg 1000 kcal⁻¹) was significantly higher in good-quality breakfast consumers (data not shown).

The intake of different food groups is presented in Table 5. Both male and female adolescents who consumed

a good-quality breakfast had significantly higher intakes of bread, fruit, vegetables, milk and milk products, and fruit juice, while their intake of soft drinks was significantly lower than those who consumed a low-quality breakfast. In girls, a larger number of differences were found. Female good-quality breakfast consumers also had significantly higher intakes of cereal products, cheese and water.

Discussion

The Iowa Breakfast Study carried out in the USA at the beginning of the 1960s was one of the first studies that addressed the importance of the role of breakfast in dietary balance, physical and cognitive performance¹⁹. To our knowledge, the present study is the first Belgian research on the relationship between breakfast parameters and the overall diet of adolescents. The current study found that

Table 3 Energy intake (kcal) at breakfast, contribution of macronutrients (as a percentage of energy) to the total energy supplied at breakfast and micronutrient intake (mg) at breakfast according to breakfast habit (low- or good-quality)

	Boys (n = 129)		P-value*	Girls (n = 212)		P-value*
	Low (n = 40)	Good (n = 89)		Low (n = 94)	Good (n = 118)	
Energy	274.8 (161.91)	478.3 (164.38)	<0.001	196.3 (113.89)	371.8 (119.34)	<0.001
Protein	10.7 (4.56)	12.4 (2.81)	0.022	12.1 (6.42)	14.1 (3.47)	<0.001
Carbohydrates, total	59.4 (15.17)	61.2 (8.46)	0.643	57.9 (18.34)	60.1 (9.20)	0.335
Mono- and disaccharides	32.3 (16.43)	30.6 (8.57)	0.699	34.3 (20.08)	30.3 (9.76)	0.389
Polysaccharides	26.8 (12.51)	30.1 (8.18)	0.278	23.4 (15.36)	29.7 (9.62)	0.002
Fat, total	27.3 (12.98)	26.3 (8.07)	0.492	25.6 (13.06)	25.7 (7.71)	0.519
SFA	12.4 (6.96)	11.2 (4.50)	0.562	11.1 (6.26)	11.3 (4.44)	0.941
MUFA	8.7 (5.08)	8.3 (3.20)	0.533	8.4 (4.98)	8.0 (3.05)	0.334
PUFA	3.8 (3.62)	3.9 (2.83)	0.418	3.9 (3.30)	3.5 (1.90)	0.775
Calcium	112.7 (98.86)	261.5 (145.54)	<0.001	108.1 (105.60)	248.5 (147.35)	<0.001
Phosphorus	154.8 (111.99)	323.1 (144.62)	<0.001	125.6 (107.44)	291.9 (157.57)	<0.001
Iron	1.6 (1.11)	2.7 (1.40)	<0.001	0.9 (1.03)	2.1 (1.16)	<0.001
Magnesium	33.1 (22.64)	66.8 (30.25)	<0.001	25.7 (17.71)	56.8 (27.54)	<0.001
Thiamin	0.2 (0.11)	0.3 (0.19)	<0.001	0.1 (0.12)	0.3 (0.16)	<0.001
Riboflavin	0.2 (0.19)	0.5 (0.31)	<0.001	0.2 (0.21)	0.5 (0.29)	<0.001
Vitamin C	3.2 (8.61)	17.1 (25.95)	<0.001	7.9 (14.28)	18.1 (21.27)	<0.001

SFA – saturated fatty acids; MUFA – monounsaturated fatty acids; PUFA – polyunsaturated fatty acids.

Data presented as mean (standard deviation).

* Mann–Whitney *U* tests.

Table 4 Total energy intake (kcal), contribution of macronutrients (as a percentage of energy) to the total energy and micronutrient intake (mg) according to breakfast habit (low- or good-quality)

	Boys (<i>n</i> = 129)			Girls (<i>n</i> = 212)		
	Low (<i>n</i> = 40)	Good (<i>n</i> = 89)	<i>P</i> -value*	Low (<i>n</i> = 94)	Good (<i>n</i> = 118)	<i>P</i> -value*
Energy	2603.9 (577.33)	2674.8 (488.20)	0.502	1841.2 (494.96)	2072.9 (384.72)	<0.001
Protein	13.9 (1.84)	14.6 (2.08)	0.058	14.1 (2.71)	15.5 (2.37)	<0.001
Carbohydrates, total	47.8 (5.28)	49.2 (4.81)	0.140	48.7 (6.06)	49.6 (4.92)	0.266
Mono- and disaccharides	22.0 (6.74)	22.7 (5.04)	0.596	22.6 (7.02)	22.7 (4.86)	0.831
Polysaccharides	22.0 (4.06)	24.0 (4.36)	0.016	22.9 (5.17)	24.4 (4.22)	0.025
Fat, total	36.8 (4.64)	35.9 (4.49)	0.327	36.5 (5.44)	34.6 (4.39)	0.005
SFA	14.6 (2.54)	14.6 (2.33)	0.990	14.6 (2.42)	14.3 (2.22)	0.386
MUFA	14.3 (2.30)	13.6 (2.17)	0.120	14.1 (2.81)	12.7 (2.17)	<0.001
PUFA	6.0 (1.48)	5.7 (1.59)	0.288	5.9 (1.62)	5.4 (1.25)	0.005
Calcium	739.9 (277.44)	991.4 (351.52)	<0.001	638.3 (278.96)	937.8 (347.02)	<0.001
Phosphorus	1315.6 (291.61)	1543.3 (363.79)	0.001	979.9 (283.37)	1304.5 (373.22)	<0.001*
Iron	12.6 (2.64)	13.8 (2.97)	0.035	8.9 (2.45)	10.9 (2.75)	<0.001
Magnesium	233.9 (52.30)	288.1 (70.28)	<0.001	179.2 (48.49)	242.3 (65.30)	<0.001
Thiamin	1.3 (0.37)	1.6 (0.87)	0.004*	0.9 (0.74)	1.3 (1.00)	<0.001 ^a
Riboflavin	1.4 (0.39)	1.8 (0.58)	<0.001	1.1 (0.53)	1.5 (0.56)	<0.001
Vitamin C	56.5 (27.80)	94.5 (51.88)	<0.001*	64.1 (42.64)	89.5 (46.35)	<0.001 ^a

SFA – saturated fatty acids; MUFA – monounsaturated fatty acids; PUFA – polyunsaturated fatty acids.

Data presented as mean (standard deviation).

*Mann–Whitney *U*-test.

consumers of a good-quality breakfast had a better overall dietary pattern – on the nutrient and food group level – than consumers of low-quality breakfast. However, the current differences in dietary intake could not be related to nutrient status due to the lack of biochemical assays of cholesterol, vitamin and mineral status. Nevertheless, the associations between breakfast patterns and biomarkers of nutrient status are not widely reported and are inconsistent. Preziosi and colleagues only found a significantly higher blood thiamin concentration in adolescents consuming high-energy breakfasts²⁰.

In the current study, some methodological considerations have to be taken into account. In this study, individual breakfast labels and scores were computed. These breakfast scores were based on both nutrient and food item recommendations. However, there is no general scientific agreement as to which foods have to be consumed and in what amounts. Therefore, the current rationale for the development of the scores is two-fold. For food items, the rationale is based on the guidelines for a healthy breakfast in schoolchildren^{21,22} and for energy the rationale is based on the assumption that an ideal breakfast should provide at least 25% of the recommended daily energy^{9,23}. This recommendation is based on the above-mentioned Iowa Breakfast Study¹⁹. In addition, by introducing a breakfast score, the authors are aware that a limitation could be introduced due to the lack of information about the heterogeneity and variability of the breakfast consumption pattern of all adolescents, both within and between the good- and low-quality breakfast consumers. As the classification method can group subjects with widely differing intakes into one category and subjects with very similar intakes into different categories if they are close to the cut-off point, this could

introduce difficulties for the interpretation of the results. Another possible disadvantage of the breakfast score algorithm used is that a subject receives a lower score when he/she does not consume one of the selected foods. A methodological issue that also has to be taken into account is the sample size. The total sample size contains more girls than boys. The limited number of boys could cause a lack of power in the analyses. The originally selected sample contained fewer boys than girls. This could be explained by the study design where male-only schools were less well represented. It is not clear in what way this may have affected the results of the study but it does not affect the value of the within-gender analyses. However, the strength of the current study is the use of the 7-day dietary record, which reflects the usual dietary intake and reflects in that way the usual breakfast consumption of the different participants. It is not clear how the possible impact of the possible limitations may have affected the results of the study.

The importance of breakfast in ensuring adequate nutrient intake in children and adolescents has been documented in different studies^{7,20,24–26}. The current findings, i.e. that consumers of a good-quality breakfast have a better overall dietary pattern than those who consume a low-quality breakfast, are in line with these European studies. However, an inherent problem with comparing different breakfast studies are the methodological differences regarding dietary intake data collection and how breakfast is defined (types, amounts, categories). These inconsistencies should be kept in mind when comparing different studies. In the current study, female good quality breakfast consumers have a higher energy intake, while no difference was found in boys. In the Göteborg Adolescence Study (15–16 years), a similar trend was found, i.e. female adolescents eating breakfast on

Table 5 Intake of some food groups (g day⁻¹) in adolescents according to breakfast habit (low- or good-quality)

	Boys (n = 129)					Girls (n = 212)				
	Low (n = 40)		Good (n = 89)		P-value*	Low (n = 94)		Good (n = 118)		P-value*
	Mean (SD)	Median	Mean (SD)	Median		Mean (SD)	Median	Mean (SD)	Median	
Potatoes	159.2 (74.42)	149.29	140.1 (84.33)	127.86	0.094	106.8 (58.46)	98.14	95.8 (53.11)	90.11	0.152
Bread, rusk and breakfast rolls	172.2 (69.97)	170.43	206.3 (77.27)	193.57	0.028	123.1 (51.08)	124.64	165.7 (50.81)	162.64	<0.001
Egg	12.3 (12.64)	12.11	7.6 (8.86)	4.29	0.116	8.9 (10.26)	4.50	9.7 (12.77)	5.89	0.838
Fruit	61.6 (74.31)	33.21	95.2 (71.30)	71.86	0.002	84.1 (87.94)	68.00	138.6 (101.61)	117.14	<0.001
Cakes and biscuits	32.6 (33.46)	20.57	39.6 (29.34)	33.43	0.070	29.2 (23.73)	24.39	32.1 (26.47)	24.86	0.515
Poultry	28.2 (23.63)	21.43	29.6 (31.59)	22.86	0.693	22.9 (24.26)	17.86	23.2 (22.81)	18.79	0.782
Cereal products and binding agents	50.8 (40.33)	46.68	66.7 (59.76)	58.14	0.245	37.8 (43.02)	25.11	56.9 (43.99)	47.64	<0.001
Vegetables	111.5 (60.54)	100.46	131.3 (64.72)	120.86	0.048	90.5 (55.43)	75.39	116.0 (57.26)	107.71	0.001
Cheese	25.8 (24.49)	17.14	26.5 (23.59)	19.00	0.766	22.1 (21.14)	15.07	31.1 (21.65)	30.14	0.001
Milk and milk products	165.6 (127.77)	120.07	318.1 (229.66)	292.00	<0.001	152.7 (167.21)	110.86	307.8 (239.60)	251.96	<0.001
Nuts, seeds and snacks	20.2 (21.85)	15.18	15.3 (19.15)	8.00	0.238	13.6 (17.07)	8.57	10.8 (12.65)	7.93	0.438
Sugar, confectionery, sweet fillings and sweet sauces	42.6 (37.07)	32.00	48.9 (33.75)	39.43	0.164	32.7 (23.46)	25.14	32.9 (23.15)	27.79	0.646
Fats, oils and savoury sauces	52.8 (23.77)	53.07	49.9 (27.33)	46.36	0.288	37.2 (20.62)	33.00	33.3 (15.73)	31.54	0.277
Fish and fish products	16.8 (21.77)	10.00	18.6 (22.22)	14.29	0.772	13.6 (15.64)	8.57	15.5 (17.55)	9.29	0.581
Meat and meat products	152.3 (45.88)	144.25	141.8 (60.92)	140.00	0.335	88.9 (48.82)	88.57	94.9 (49.63)	86.57	0.313
Water	380.1 (424.41)	208.96	457.5 (370.08)	406.79	0.078	411.9 (345.91)	326.50	559.9 (367.16)	517.57	0.002
Soft drinks	636.3 (413.14)	608.57	361.7 (291.16)	322.86	<0.001	317.7 (310.26)	215.00	170.6 (192.89)	104.29	<0.001
Fruit juice	68.1 (107.45)	28.57	171.4 (187.73)	130.00	0.001	106.1 (146.45)	57.14	166.0 (148.31)	142.86	<0.001

SD – standard deviation.

* Mann–Whitney *U*-test.

a regular basis had significantly higher energy intake compared with those with irregular breakfast intake¹¹. In the current study, female good-quality breakfast consumers had a relative higher intake of total protein and polysaccharides, and a relative lower intake of total fat, monounsaturated fatty acids and polyunsaturated fatty acids. In boys, only a relative higher intake of polysaccharides in good-quality breakfast consumers was found. In a French study, adolescents (10–18 years) consuming a high-energy breakfast (>25% of energy from breakfast) had higher total daily intakes (expressed as percentage energy) of carbohydrates but lower intake of total fat and saturated fat than their counterparts²⁰. Swedish regular breakfast consumers had a higher relative intake of total protein and a lower relative intake of sucrose¹¹. However, in the literature, conflicting results can be found. De Graaf and co-workers showed that neither energy content nor macronutrient composition of breakfast had any effect on energy and macronutrient intake consumed over the rest of the day²⁷.

Good-quality breakfast consumers have a higher intake of micronutrients. In a French, Spanish and Swedish adolescent population, the intake of vitamin C, thiamin, riboflavin, calcium, iron and zinc was higher in regular breakfast consumers (Sweden) or high-energy breakfast consumers (France, Spain)^{11,20,26}. Adolescents eating a low-quality breakfast seem not to be able to make up the low micronutrient intake at other meals during the day. In a recent review, all the above findings are reflected and it is clear that there is evidence that breakfast consumption significantly contributes to the overall nutrient adequacy of the diet³.

In the current study, in girls, the group who consume a good-quality breakfast is associated with a lower proportion of overweight adolescents, despite the higher daily energy intake. Skipping breakfast is a popular method of losing weight among adolescents²⁸ and it is shown that adolescents who have the perception of a too high body weight may be more likely to skip breakfast¹¹. Skipping breakfast may lead to hunger in the morning and result in increased snacking. From a public health point of view, this phenomenon of increased snacking should be associated with a higher consumption of whole-grain products, fruit, vegetables and water. Unfortunately, snack foods commonly consumed by adolescents tend to be high in added sugars and fat, and low in minerals and vitamins. This could lead to a less healthy dietary pattern. In addition, the food availability at schools could play a role in this increased consumption of 'empty calories'. A recent study showed that the majority of the food items available in schools, supplied by school shops or vending machines, could be categorised as 'empty calories'²⁹. Due to these factors, i.e. skipping breakfast, increased snacking on 'empty calories', the availability of poor-quality food items and weight gain, adolescents could get caught in a 'vicious circle'. However, based on several cross-sectional studies, no uniform association between skipping breakfast and higher weight

or BMI is found in adolescents³. Some studies, however, state that breakfast consumption is part of a weight-reduction programme³⁰ and it is one of the factors affecting weight-loss maintenance in adults³¹. However, to date, to our knowledge, there have been no randomised controlled trials investigating breakfast consumption and weight control in adolescents. Breakfast consumption, a regular meal rhythm and an increase in feeding frequencies seem to have a positive impact on weight management³². However, there is no scientific consensus about specific recommendations concerning meal frequency and health outcomes^{33,34}.

Good-quality breakfast consumers tend to make more healthy food choices during the day, such as the consumption of more vegetables, fruit, milk and milk products, and water, and a lower consumption of soft drinks. The significantly higher intake of milk and milk products and cheese in female good-quality breakfast consumers is possibly the main contributors of the higher saturated fatty acid intake. A similar pattern was found in Swedish adolescents; girls with irregular breakfast and lunch intake make less healthy food choices and consume more soft drinks and less milk, vegetables and fruit¹¹.

Several studies have mentioned the importance of ready-to-eat breakfast cereals; the nutritional benefits of these cereals are associated with improved compliance with dietary recommendations and improved nutrient status based on biochemical measures^{35,36}. In the current study, <50% of the adolescents consumed ready-to-eat breakfast cereals and the average portion size was <20 g. As the intake of these food items is rather small, the influence of these cereals could be neglected in this study.

Several authors emphasise the importance of breakfast and that it should be included in educational programmes promoting healthy diets and lifestyles³⁷ or even included in school-based nutrition education programmes, involving families, teachers and others in achieving healthier patterns⁷. In the UK, the Department of Health initiated breakfast club schemes in schools. Breakfast clubs are a form of before-school provision serving food to children who arrive early. The main aims of the scheme were to provide breakfast to children who might otherwise not have eaten, to establish a positive relationship at the start of the school day and to offer children a choice of healthy food, which may help to encourage healthier eating habits. However, a recent study indicates that children who attend breakfast clubs have a poorer nutrient intake than other children at the same schools³⁸. It has been shown that parents eating breakfast is associated with adolescents eating breakfast. This suggests that breakfast programmes that address the whole family or just parents may be more effective¹.

The present results and the literature show that the nutritional profile of Belgian adolescents could be substantially improved by the consumption of a healthy breakfast in a family setting on a daily basis, consisting of a

variety of foods, namely whole-grain products, fruit and (semi-) skimmed milk products or an alternative source of calcium.

Acknowledgements

The original food consumption survey was financially supported by the National Fund for Scientific Research (fund no. 31557898), the Kellogg's Benelux Company, Unilever Belgium, the Belgian Nutrition Information Center and the 'Vlaams Wetenschappelijk Fonds'.

References

- Keski-Rahkonen A, Kaprio J, Rissanen A, Virkkunen M, Rose RJ. Breakfast skipping and health-compromising behaviors in adolescents and adults. *European Journal of Clinical Nutrition* 2003; **57**: 842–53.
- Ruxton CH, Kirk TR. Breakfast: a review of associations with measures of dietary intake, physiology and biochemistry. *British Journal of Nutrition* 1997; **78**: 199–213.
- Rampersaud GC, Pereira MA, Girard BL, Adams J, Metz J. Breakfast habits, nutritional status, body weight, and academic performance in children and adolescents. *Journal of American Dietetic Association* 2005; **105**: 743–60.
- Pollitt E. Does breakfast make a difference in school? *Journal of the American Dietetic Association* 1995; **95**: 1134–9.
- Pollitt E, Mathews R. Breakfast and cognition: an integrative summary. *American Journal of Clinical Nutrition* 1998; **67**: 804S–13S.
- Rogers PJ. How important is breakfast? *British Journal of Nutrition* 1997; **78**: 197–8.
- Aranceta J, Serra-Majem L, Ribas L, Perez-Rodrigo C. Breakfast consumption in Spanish children and young people. *Public Health Nutrition* 2001; **4**: 1439–44.
- Affenito SG, Thompson DR, Barton BA, Franko DL, Daniels SR, Obarzanek E, *et al.* Breakfast consumption by African-American and white adolescent girls correlates positively with calcium and fiber intake and negatively with body mass index. *Journal of the American Dietetic Association* 2005; **105**: 938–45.
- Morgan KJ, Zabik ME, Stampley GL. The role of breakfast in diet adequacy of the U.S. adult population. *Journal of American College of Nutrition* 1986; **5**: 551–63.
- Navia B, Requejo AM, Ortega RM, Lopez Sobaler AM, Quintas ME, Andres P, *et al.* The relationship between breakfast and whole diet energy profiles in a group of preschool children. *Annals of Nutrition & Metabolism* 1997; **41**: 299–306.
- Sjoberg A, Hallberg L, Hoglund D, Hulthen L. Meal pattern, food choice, nutrient intake and lifestyle factors in The Goteborg Adolescence Study. *European Journal of Clinical Nutrition* 2003; **57**: 1569–78.
- Matthys C, De Henauw S, Devos C, De Backer G. Estimated energy intake, macronutrient intake and meal pattern of Flemish adolescents. *European Journal of Clinical Nutrition* 2003; **57**: 366–75.
- NEVO. *NEVO Tabel, Nederlands Voedingsstoffenbestand*. Zeist: NEVO, 1993.
- NUBEL. *Belgische Voedingsmiddelentabel*. Brussels: Ministerie van Volksgezondheid, 1992.
- NUBEL. *Belgische Voedingsmiddelentabel, Tweede Uitgave*. Brussels: Ministerie van Volksgezondheid, 1995.
- Unilever. *Becel Voedingsprogramma*. Rotterdam: Nederlandse Unilever Bedrijven BV, 1992.
- Goldberg GR, Black AE, Jebb SA, Cole TJ, Murgatroyd PR, Coward WA, *et al.* Critical evaluation of energy intake data using fundamental principles of energy physiology: 1. Derivation of cut-off limits to identify under-recording. *European Journal of Clinical Nutrition* 1991; **45**: 569–81.
- Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *British Medical Journal* 2000; **320**: 1240–3.
- Cereal Institute Inc. *A Complete Summary of the Iowa Breakfast Studies*. Chicago, IL: Cereal Institute Inc., 1962.
- Preziosi P, Galan P, Deheeger M, Yacoub N, Drewnowski A, Hercberg S. Breakfast type, daily nutrient intakes and vitamin and mineral status of French children, adolescents, and adults. *Journal of American College of Nutrition* 1999; **18**: 171–8.
- Pérez-Rodrigo C, Ribas Barba L, Serra Majem L, Aranceta Bartina J. Recomendaciones para un desayuno saludable. In: Serra Majem L, Aranceta Bartina J, eds. *Desayuno y Equilibrio Alimentario*. Barcelona: Masson, 2000; 91–7.
- Vlaams Instituut voor Gezondheidspromotie. *De Voedingsdriehoek: Een Praktische Voedingsgids [The Food Triangle: A Practical Guide]*. Brussels: Vlaams Instituut voor Gezondheidspromotie, 2003.
- Morgan KJ, Zabik ME, Leveille GA. The role of breakfast in nutrient intake of 5- to 12-year-old children. *American Journal of Clinical Nutrition* 1981; **34**: 1418–27.
- Hercberg S, Preziosi P, Galan P, Yacoub N, Kara G, Deheeger M. La consommation du petit-déjeuner dans l'étude du Val-De-Marne. 3. La valeur nutritionnelle du petit-déjeuner et ses relations avec l'équilibre nutritionnel global et le statut minéral et vitaminique. *Cahier Nutritionnelle et Diététique* 1996; **31**(Suppl. 1): 18–24.
- Baric IC, Satalic Z. Breakfast quality differences among children and adolescents in Croatia. *International Journal of Food Science and Nutrition* 2002; **53**: 79–87.
- Ortega RM, Requejo AM, Lopez-Sobaler AM, Andres P, Quintas ME, Navia B, *et al.* The importance of breakfast in meeting daily recommended calcium intake in a group of schoolchildren. *Journal of American College of Nutrition* 1998; **17**: 19–24.
- De Graaf C, Hulshof T, Weststrate JA, Jas P. Short-term effects of different amounts of protein, fats, and carbohydrates on satiety. *American Journal of Clinical Nutrition* 1992; **55**: 33–8.
- Lattimore PJ, Halford JC. Adolescence and the diet-dieting disparity: healthy food choice or risky health behaviour? *British Journal of Health Psychology* 2003; **8**: 451–63.
- Vereecken CA, Bobelijn K, Maes L. School food policy at primary and secondary schools in Belgium-Flanders: does it influence young people's food habits? *European Journal of Clinical Nutrition* 2005; **59**: 271–7.
- Schlundt DG, Hill JO, Sbrocco T, Pope-Cordle J, Sharp T. The role of breakfast in the treatment of obesity: a randomized clinical trial. *American Journal of Clinical Nutrition* 1992; **55**: 645–51.
- Elfhag K, Rossner S. Who succeeds in maintaining weight loss? A conceptual review of factors associated with weight loss maintenance and weight regain. *Obesity Reviews* 2005; **6**: 67–85.
- Louis-Sylvestre J, Lluch A, Neant F, Blundell JE. Highlighting the positive impact of increasing feeding frequency on metabolism and weight management. *Forum Nutrition* 2003; **56**: 126–8.
- Mattson MP. The need for controlled studies of the effects of meal frequency on health. *Lancet* 2005; **365**: 1978–80.
- Gibney MJ, Wolever TM. Periodicity of eating and human health: present perspective and future directions. *British Journal of Nutrition* 1997; **77**(Suppl 1): S3–5.
- Gibson S. Micronutrient intakes, micronutrient status and lipid profiles among young people consuming different

- amounts of breakfast cereals: further analysis of data from the National Diet and Nutrition Survey of Young People aged 4 to 18 years. *Public Health Nutrition* 2003; **6**: 815–20.
- 36 Galvin MA, Kiely M, Flynn A. Impact of ready-to-eat breakfast cereal (RTEBC) consumption on adequacy of micronutrient intakes and compliance with dietary recommendations in Irish adults. *Public Health Nutrition* 2003; **6**: 351–63.
- 37 Gassin AL. Helping to promote healthy diets and lifestyles: the role of the food industry. *Public Health Nutrition* 2001; **4**: 1445–50.
- 38 Belderson P, Harvey I, Kimbell R, O'Neill J, Russell J, Barker ME. Does breakfast-club attendance affect schoolchildren's nutrient intake? A study of dietary intake at three schools. *British Journal of Nutrition* 2003; **90**: 1003–6.