


## Concise Communication

# Sensitivity of midturbinate versus nasopharyngeal swabs for the detection of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)

Alainna J. Jamal PhD<sup>1,2,3</sup> , Mohammad Mozafarihashjin MD<sup>2</sup>, Eric Coomes MD<sup>3</sup>, Sofia Anceva-Sami MD<sup>2</sup>, Shiva Barati MD<sup>2</sup>, Gloria Crowl MSc<sup>2</sup>, Amna Faheem MD<sup>2</sup>, Lubna Farooqi MD<sup>2</sup>, Christopher E. Kandel MD<sup>1,2,3</sup>, Saman Khan MD<sup>2</sup>, Angel X. Li MSc<sup>2</sup>, Henna Mistry BSc<sup>4</sup>, Aimee Paterson MSc<sup>2</sup>, Simon Plenderleith BSc<sup>4</sup>, Karren Prost DVM<sup>4</sup>, Susan Poutanen MD<sup>2,3</sup>, Jeff Powis MD<sup>5</sup>, Renée Schryer BSc<sup>4</sup>, Maureen Taylor PA<sup>5</sup>, Lily Yip MSc<sup>4</sup>, Xi Zoe Zhong PhD<sup>2</sup>, Allison J. McGeer MD<sup>1,2,3</sup>, Samira Mubareka MD<sup>3,4</sup> and for the Toronto Invasive Bacterial Diseases Network COVID-19 Investigators<sup>a</sup>

<sup>1</sup>Institute of Health Policy, Management and Evaluation, University of Toronto, Toronto, Canada, <sup>2</sup>Sinai Health System, Toronto, Canada, <sup>3</sup>Faculty of Medicine, University of Toronto, Toronto, Canada, <sup>4</sup>Sunnybrook Health Sciences Centre, Toronto, Canada and <sup>5</sup>Michael Garron Hospital, Toronto, Canada

## Abstract

To compare sensitivity of specimens for COVID-19 diagnosis, we tested 151 nasopharyngeal/midturbinate swab pairs from 117 COVID-19 inpatients using reverse-transcriptase polymerase chain reaction (RT-PCR). Sensitivity was 94% for nasopharyngeal and 75% for midturbinate swabs ( $P = .0001$ ). In 88 nasopharyngeal/midturbinate pairs with matched saliva, sensitivity was 86% for nasopharyngeal swabs and 88% for combined midturbinate swabs/saliva.

(Received 20 August 2020; accepted 9 November 2020)

Nasopharyngeal (NP) swabs are currently the preferred specimen collection method for coronavirus disease 2019 (COVID-19) diagnosis. However, there is increasing interest in midturbinate (MT) swabs because they cause less discomfort to patients, may be safer for healthcare workers to collect, and can be self-collected by adolescent and adult patients.<sup>1,2</sup>

Two small studies have compared the sensitivity of NP and MT swabs for detecting severe acute respiratory coronavirus virus 2 (SARS-CoV-2) and found comparable sensitivity early in illness.<sup>2,3</sup> In our prospective cohort of inpatients with COVID-19 in Ontario,

Canada, we compared the sensitivities of NP and saliva versus MT swabs.

## Methods

### Study population

The Toronto Invasive Bacterial Diseases Network (TIBDN) performs population-based surveillance for infectious diseases in Toronto and Peel (Ontario, Canada). The TIBDN clinical microbiology laboratories report specimens yielding SARS-CoV-2 to our study office. In this study, we enrolled consecutive inpatients with COVID-19 at 4 TIBDN hospitals from March 23 through May 22, 2020. The research ethics boards of all TIBDN hospitals approved this study.

### Data collection

Demographic and clinical data were collected from charts and patient interviews. Study staff collected NP and MT swabs and saliva from consenting patients at enrollment and 6 days later if the patient was still hospitalized. Swabs were obtained using standard procedures<sup>4,5</sup> and placed in universal transport medium (UTM, COPAN Diagnostics, Murrietta, CA). Patients were asked to spit 1 teaspoon of saliva into a sterile container, then 2.5 mL phosphate-buffered saline was added. From April 14 through May 4, in a pilot test of alternative collection methods, saliva was provided by gargle and/or salivette; these specimens were excluded.

**Author for correspondence:** Allison J. McGeer, E-mail: [allison.mcgeer@sinaihealth.ca](mailto:allison.mcgeer@sinaihealth.ca)

<sup>a</sup>Toronto Invasive Bacterial Diseases Network COVID-19 Investigators: Brenda L. Coleman (Sinai Health System, Toronto, Canada), Danny Chen (Mackenzie Health, Toronto, Canada), Nataly Farshait (Humber River Regional Hospital, Toronto, Canada), Wayne L. Gold (University Health Network, Toronto, Canada), Kevin Katz (North York General Hospital, Toronto, Canada), Robert Kozak (Sunnybrook Health Sciences Centre, Toronto, Canada), Tony Mazzulli (Sinai Health System, Toronto, Canada), Matthew Muller (St. Michael's Hospital, Toronto, Canada), Anne Opavsky (Joseph Brant Hospital, Toronto, Canada), Mario Ostrowski (St. Michael's Hospital, Toronto, Canada), Agron Plevneshi (Sinai Health System, Toronto, Canada), Neil Rau (Halton Healthcare, Oakville, Canada), Daniel Ricciuto (Lakeridge Health, Oshawa, Canada), David Richardson (William Osler Health System, Brampton, Canada), David Rose (Scarborough Health Network, Scarborough, Canada), Valerie Sales (Markham Stouffville Hospital, Toronto, Canada), Sharon Walmsley (University Health Network, Toronto, Canada).

**Cite this article:** Jamal AJ, *et al.* (2020). Sensitivity of midturbinate versus nasopharyngeal swabs for the detection of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). *Infection Control & Hospital Epidemiology*, <https://doi.org/10.1017/ice.2020.1326>

© The Author(s), 2020. Published by Cambridge University Press on behalf of The Society for Healthcare Epidemiology of America. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Laboratory methods

Specimens were aliquoted and frozen at  $-80^{\circ}\text{C}$  on the day of collection until processing at the Sunnybrook Health Sciences research microbiology laboratory (Toronto, Canada). RNA extractions were performed using the QIAamp Viral RNA Mini Kit (Qiagen, Venlo, The Netherlands). Reverse transcriptase polymerase chain reaction (RT-PCR) using the Luna Universal Probe One-Step RT-qPCR Kit (New England BioLabs, Ipswich, MA) was performed to detect the 5' untranslated region and envelope (E) gene of SARS-CoV-2, with RNaseP as internal control. Cycle thresholds (Cts) were determined using Rotor-Gene Q software (Qiagen, Venlo, The Netherlands); results where all targets had Cts  $<40$  were reported as positive.

### Statistical methods

Sensitivities of specimen types were calculated using as the denominator samples with any 1 specimen in the pair or triplet testing positive for SARS-CoV-2. The McNemar test was used to assess differences in sensitivity between specimen types. The Wilcoxon rank-sum test was used to compare the Cts of NP swabs when the MT tested positive versus negative. Two-sided  $P$  values  $<.05$  were considered statistically significant. Analyses were performed using SAS version 9.4M6 software (SAS Institute, Cary, NC).

### Results

The 117 included inpatients were confirmed to have COVID-19 with a nasal, MT, or NP swab at or prior to admission. At admission, 88 of these patients (75%) had fever and 87 (74%) had cough. Their median age was 65 years (range, 23–106); 51 (44%) were female; 91 (78%) had  $\geq 1$  comorbidity; 14 (12%) were immunocompromised; 47 (40%) required intensive care; and 18 (15%) died. The median time from illness onset to specimen collection was 11 days (interquartile range [IQR], 8–16 days).

#### NP and MT swab pairs

Overall, 151 NP/MT swab pairs were collected from these 117 patients, with at least 1 swab positive in 122 pairs (Table 1). Both NP and MT swabs were positive in 84 patients (69%), only the NP swab was positive in 31 patients (25%), and only the MT swab was positive in 7 patients (6%). The overall sensitivities of NP and MT swabs were 115 of 122 (94%; 95% CI, 90%–98%) and 91 of 122 (75%; 95% CI, 67%–83%), respectively ( $P = .0001$ , Table 1).

The difference in sensitivity between NP and MT swabs increased with time from illness onset to specimen collection, but it was statistically significant even in the first week of illness: 33 of 33 (100%; 95% CI, 90%–100%) versus 28 of 33 (85%; 95% CI, 67%–94%), respectively ( $P = .03$ ) (Table 1).

The median E gene Ct of NP swabs was 25 (IQR, 22–29) for positive ( $n = 84$ ) and 32 (IQR, 29–35) for negative ( $n = 31$ ) MT swabs ( $P < .0001$ ). The E gene Cts of NP and MT swabs increased with time from illness onset (Spearman's  $\rho = 0.4$  [ $P < .0001$ ] and  $\rho = 0.5$  [ $P < .0001$ ], respectively).

#### NP swab, MT swab, and saliva triplets

A corresponding saliva specimen was available for 88 of the 151 NP/MT swab pairs. These 88 triplets were from 75 patients; at least 1 specimen was positive in 74 triplets. NP swabs detected SARS-CoV-2 in 64 of the 74 (86%; 95% CI 77–92%) triplets with at least one positive specimen, saliva detected 55 (74%; 95% CI 63–83%),

**Table 1.** Results of Testing of 151 NP and MT Swab Pairs for SARS-CoV-2 Among Hospitalized Patients With COVID-19, by Time From Illness Onset to Collection of Swab Pair

Time From Illness Onset to Collection of Swab Pair	No. (%) Swab Pairs			
	MT and NP Swabs Positive	Only NP Swab Positive	Only MT Swab Positive	MT and NP Swabs Negative
Day 0–7 (n=35)	28 (80)	5 (14)	0	2 (6)
Day 8+ (n=116)	56 (48)	26 (22)	7 (6)	27 (23)
All (n=151)	84 (56)	31 (21)	7 (5)	29 (19)

Note. COVID-19, coronavirus disease 2019; SARS-CoV-2, severe acute respiratory syndrome virus 2; MT, midturbinate; NP, nasopharyngeal.

MT swabs detected 49 (66%; 95% CI 54–77%), and MT and saliva in combination detected 65 (88%; 95% CI 78–94%) (Table 2).

For all analyses, results were similar when only the first or last pair or triplet for each patient was analyzed (Supplementary Material online).

### Discussion

In this prospective cohort study of hospitalized patients with COVID-19, NP swabs were significantly more sensitive for SARS-CoV-2 detection than MT swabs. The difference in sensitivity between NP and MT swabs was greater later in illness. Saliva demonstrated intermediate sensitivity between MT and NP swabs.

We are aware of 2 studies comparing sensitivity of MT and NP swabs for SARS-CoV-2 detection.<sup>2,3</sup> Pinninti *et al*,<sup>3</sup> in inpatients similar to ours, reported NP swabs as more sensitive than MT swabs (80% vs 64% overall), with the difference in sensitivity increasing with time from illness onset. However, our difference in sensitivity between NP and MT swabs was statistically significant even in the first week of illness, whereas that of Pinninti *et al* was not. In an outpatient setting, Tu *et al*<sup>2</sup> found that NP swabs detected all 52 patients with either swab positive, while MT swabs detected 50.

Our previous analysis comparing saliva to NP swabs in this cohort using a different testing platform indicated that saliva was less sensitive than NP swabs, particularly later in illness.<sup>6</sup> The difference was not statistically significant in this sample, in which specimens were collected less frequently but over a longer period, likely due to sampling variability. More importantly, however, other studies have not found differences in sensitivities between NP swabs and saliva, particularly early in illness.<sup>7,8</sup> Saliva is an appealing alternative to NP swabs, especially during swab shortages or in settings requiring repeated COVID-19 screening.

The fact that MT swabs and saliva in combination have a similar sensitivity to NP swabs may simply be because a second specimen adds sensitivity; repeated testing is known to improve sensitivity among patients with COVID-19.<sup>9</sup> These data reinforce the concept that 1 specimen is not 100% sensitive and that patients with a high index of clinical suspicion for COVID-19 who test negative require repeated testing.

This study had several limitations. It included only hospitalized patients, with first study specimen at 1–27 days after diagnosis. Differences in sensitivity may be smaller at presentation. We only tested specimens on a single platform; results may vary between platforms, especially when Ct values are near the limit of detection. We did not perform sequencing or other validation of results in samples in which only 1 specimen

**Table 2.** Results of Testing of 88 NP Swab/MT Swab/Saliva Triplets for SARS-CoV-2 Among Hospitalized Patients With COVID-19, by Time From Illness Onset to Collection of Triplet

Time From Illness Onset to Collection of Triplet	No. (%) Triplets							
	NP, MT, and Saliva Positive	Only NP and MT Positive	Only NP and Saliva Positive	Only MT and Saliva Positive	Only NP Positive	Only MT Positive	Only Saliva Positive	NP, MT, and Saliva Negative
Day 0–7 (n=19)	14 (74)	1 (5)	2 (11)	0	1 (5)	0	0	1 (5)
Day 8+ (n=69)	24 (35)	8 (12)	6 (9)	1 (1)	8 (12)	1 (1)	8 (12)	13 (19)
All (n=88)	38 (43)	9 (10)	8 (9)	1 (1)	9 (10)	1 (1)	8 (9)	14 (16)

Note. COVID-19, coronavirus disease 2019; SARS-CoV-2, severe acute respiratory syndrome virus 2; MT, mid-turbinate; NP, nasopharyngeal.

yielded positive results, so some results may have been false positives.

In conclusion, this study of hospitalized patients found NP swabs to be 15% more sensitive than MT swabs for SARS-CoV-2 detection in the first week of COVID-19. Larger studies are needed, as are studies of specimens obtained for initial diagnosis and in milder illness. In the interim, we believe that the use of MT swabs for COVID-19 diagnosis should be approached with caution.

**Acknowledgments.** We thank the patients with COVID-19 who agreed, in difficult circumstances, to help with research. We are grateful to the many laboratory, infection prevention and control, public health, and research ethics staff who make the work of the Toronto Invasive Bacterial Diseases Network possible.

**Financial support.** This work was supported by the Canadian Institutes of Health Research (Canadian 2019 Novel Coronavirus [COVID-19] Rapid Research Application no. 440359). A.J.J. is supported by the Vanier Canada Graduate Scholarship.

**Conflicts of interest.** All authors report no conflicts of interest relevant to this article.

**Supplementary material.** To view supplementary material for this article, please visit <https://doi.org/10.1017/ice.2020.1326>

## References

- Wehrhahn MC, Robson J, Brown S, *et al.* Self-collection: an appropriate alternative during the SARS-CoV-2 pandemic. *J Clin Virol* 2020;128:104417.
- Tu Y-P, Jennings R, Hart B, *et al.* Swabs collected by patients or healthcare workers for SARS-CoV-2 testing. *N Engl J Med* 2020;383:494–496.
- Pinninti S, Trieu C, Pati S, *et al.* Comparing nasopharyngeal and mid-turbinate nasal swab testing for the identification of SARS-CoV-2. *Clin Infect Dis* 2020. doi: [10.1093/cid/ciaa882](https://doi.org/10.1093/cid/ciaa882).
- Marty F, Chen K, Verrill K. How to obtain a nasopharyngeal swab specimen. *N Engl J Med* 2020;382. doi: [10.1056/NEJMvcm2010260](https://doi.org/10.1056/NEJMvcm2010260).
- Interim guidelines for collecting, handling, and testing clinical specimens for COVID-19. Centers for Disease Control and Prevention website. <https://www.cdc.gov/coronavirus/2019-ncov/lab/guidelines-clinical-specimens.html>. Published 2020. Accessed November 16, 2020.
- Jamal AJ, Mozafarihashjin M, Coomes E, *et al.* Sensitivity of nasopharyngeal swabs and saliva for the detection of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). *Clin Infect Dis* 2020. doi: [10.1093/cid/ciaa848](https://doi.org/10.1093/cid/ciaa848).
- Cheuk S, Wong Y, Tse H, *et al.* Posterior oropharyngeal saliva for the detection of SARS-CoV-2. *Clin Infect Dis* 2020. doi: [10.1093/cid/ciaa797](https://doi.org/10.1093/cid/ciaa797).
- Wyllie A, Fournier J, Casanovas-Massana A, *et al.* Saliva is more sensitive for SARS-CoV-2 detection in COVID-19 patients than nasopharyngeal swabs. *N Engl J Med* 2020;383:1283–1286.
- Arevalo-Rodriguez I, Buitrago-Garcia D, Simancas-Racines D, *et al.* False-negative results of initial RT-PCR assays for COVID-19: a systematic review. *medRxiv* 2020. doi: [10.1101/2020.04.16.20066787](https://doi.org/10.1101/2020.04.16.20066787).