

# Suicidal risk factors in major affective disorders

Ross J. Baldessarini, Leonardo Tondo, Marco Pinna, Nicholas Nuñez and Gustavo H. Vázquez

## Background

Rates and risk factors for suicidal behaviour require updating and comparisons among mood disorders.

## Aims

To identify factors associated with suicidal risk in major mood disorders.

## Method

We considered risk factors before, during and after intake assessments of 3284 adults with/without suicidal acts, overall and with bipolar disorder (BD) versus major depressive disorder (MDD), using bivariate comparisons, multivariable regression modelling and receiver operating characteristic (ROC) analysis.

## Results

Suicidal prevalence was greater in BD versus MDD: ideation, 29.2 versus 17.3%; attempts, 18.8 versus 4.78%; suicide, 1.73 versus 0.48%; attempts/suicide ratio indicated similar lethality, 10.9 versus 9.96. Suicidal acts were associated with familial BD or suicide, being divorced/unmarried, fewer children, early abuse/trauma, unemployment, younger onset, longer illness, more dysthymic or cyclothymic temperament, attention-deficit hyperactivity disorder (ADHD), substance misuse, mixed features, hospital admission, percentage time unwell, less antidepressants and more antipsychotics and mood stabilisers. Logistic regression found five independent factors: hospital

admission, more depression at intake, BD diagnosis, onset age  $\leq 25$  years and mixed features. These factors were more associated with suicidal acts in BD than MDD: percentage time depressed/ill, alcohol misuse, >4 pre-intake depressions, more dysthymic/cyclothymic temperament and prior abuse/trauma. ADHD and total years ill were similar in BD and MDD; other factors were more associated with MDD. By ROC analysis, area under the curve was 71.3%, with optimal sensitivity (76%) and specificity (55%) with any two factors.

## Conclusions

Suicidal risks were high in mood disorders: ideation was highest with BD type II, attempts and suicides (especially violent) with BD type I. Several risk factors for suicidal acts differed between BD versus MDD patients.

## Declaration of interest

No author or immediate family member has financial relationships with commercial entities that might appear to represent potential conflicts of interest with the information presented.

## Keywords

Bipolar disorders; major depression; risk factors; suicide; suicide attempt.

## Copyright and usage

© The Royal College of Psychiatrists 2019.

Suicide and attempts are prevalent among psychiatric patients and constitute a major global, clinical and public health concern. The risks are particularly high with major mood disorders, especially with bipolar disorder (BD) and severe major depressive disorder (MDD).<sup>1–5</sup> Suicide has been associated with various ‘risk factors’, as reviewed in the Supplementary Information available at <https://doi.org/10.1192/bjp.2019.167>. Despite the substantial literature on risk factors for suicide, there are surprisingly few direct comparisons of factors across patients with mood disorders evaluated under the same conditions. Clarifying risk factors for individual types of major affective disorders should enhance earlier identification of suicidal risk, support preventive interventions and improve the treatment and prognosis of patients at risk. Accordingly, this study aimed to compare demographic and clinical risk factors in large samples of patients diagnosed with MDD or type I or II BD. We hypothesised that both rates and associated risk factors would differ by diagnosis, especially between patients diagnosed with BD versus MDD.

## Method

### Clinical assessments

The study sample included patients with diagnoses (updated to meet DSM-5 [2013] criteria) of BD type I (BD-I), BD type II (BD-II) or major depressive disorder (MDD) at the Lucio Bini Mood Disorders Center in Cagliari, Sardinia. Included participants had repeated assessments of clinical history and of suicidal behaviour (attempts or suicides) or reported suicidal ideation over several years of follow-up. Suicide attempt included any self-injurious act, with or without evidence of intent to die; violent acts included

severe self-injury with medical intervention or death, or the involvement of gunshot, hanging, drowning or asphyxiation or jumping from height. All participants underwent systematic initial and repeated diagnostic and follow-up assessments by the same mood disorders expert (L.T.). Clinical information concerning demographic, descriptive and clinical characteristics – including prospectively evaluated morbidity over time, characteristics of suicidal behaviours and treatments given – was derived from semi-structured interviews and life charts constructed at intake and updated during prospective, clinical follow-up (weekly through an index episode of illness at intake and at 1- to 6-monthly intervals thereafter, as indicated clinically). Clinical data were recorded systematically, converted to a digitised summary form from 2000 and data in computerised records were coded to protect patient identity – all as detailed previously.<sup>6–9</sup> Participants provided voluntary, written informed consent at clinic entry for the collection and analysis of data to be presented anonymously in aggregate form. This is in accordance with the requirements of Italian law, which states that a specific authorisation from local ethical committees is not required when data are anonymous, aggregated and identified by individual codes.<sup>10</sup> Data management also complied with the US Health Insurance Portability and Accountability Act which pertains to confidentiality of patient records.

### Data analysis

Data are presented as means with 95% confidence intervals or proportions (percentage) and rates (percentage per year). Preliminary tests of association used standard bivariate comparisons based on analysis of variance methods (*t*-score) for continuous data and contingency tables ( $\chi^2$ ) for categorical data. We evaluated relative risk

ratios for prevalence or magnitude of each tested factor for association with suicidal status, overall and for BD versus MDD. The resulting *P*-values were not adjusted for multiple comparisons to guide the selection of factors for multivariable logistic modelling which generated odds ratios and their confidence intervals and  $\chi^2$  values. Factors supported by logistic regression modelling were included in receiver operating characteristic (ROC) analyses of Bayesian sensitivity (true positive rate) versus (100 – specificity) (false positive rate) to compute the area under the curve (AUC) as a percentage. Analyses used the following commercial software: Statview version 5 for Mac (SAS Institute, Cary, NC) for spreadsheets and Stata version 13 for Mac (StataCorp, College Station, TX) for analyses.

## Results

### Sample and exposures times

The study sample included 3284 patients with one of the following DSM-5 major mood disorders: BD-I (*n* = 714), BD-II (*n* = 497), all BD (*n* = 1211) or unipolar MDD (*n* = 2073). The duration of illness averaged 17.0 years with BD and 11.7 years with MDD (Table 1); 64.4% were women. Participants were followed prospectively at the study site for an overall average of 2.95 (95% CI 2.74–3.16) years.

### Suicidal risks

The lifetime risk (percentage of patients) of identified suicidal ideation was significantly greater among patients with BD-II (35.0%) than those with BD-I (25.2%), and among patients with BD (29.2%) than those with MDD (17.3%), as were the respective annualised rates (1.92 *v.* 1.55%/year and 1.72 *v.* 1.47%/year; Table 1). The lifetime risk of suicide attempts was non-significantly higher among participants with BD-I (20.9%) than those with BD-II (15.9%); but this risk was significantly 3.93-times greater among all patients with BD (18.8%) than those with MDD (4.78%). The exposure time-adjusted attempt rates (percentage per year) were higher in those with BD-I versus BD-II (1.28 for BD-I *v.* 0.88 for BD-II, 1.45-fold higher; and 1.11 for BD-I *v.* 0.41 for BD-II, 2.72-fold higher) and for those with BD versus MDD (Table 1). The risks and rates of suicide were similar between those with BD-I (1.82%; 0.11%/year) and BD-II (1.61%; 0.09%/year), but significantly greater with those with BD overall (1.73%; 0.10%/year) than those with MDD (0.48%; 0.04%/year). In terms of the risks and rates for all suicidal acts (attempts + suicides), patients with BD-I (22.7%; 1.39%/year) had the highest rates, followed by those with BD-II (17.5%; 0.97%/year) and then by those with MDD (5.26%; 0.45%/year); these rates were high for patients with BD overall (20.5%; 1.21%/year).

The attempts/suicides ratio of rates (%/year), a proposed measure of lethality (greater lethality with lower ratio), indicated similar lethality among all diagnostic groups: BD-II (9.78), MDD (10.2), BD-I (11.6) and all BD (11.1; Table 1). The proportion of violent attempts or suicides (involving gunshot, jumping, hanging, drowning or asphyxiation) of all suicidal acts was greater among patients with BD-I than those with BD-II (38.2 *v.* 23.0%), and not significantly greater in patients with BD versus those with MDD (33.1 *v.* 23.9%). In addition, as expected, violent acts were more prevalent among men by 1.66-fold (40.8%, 95% CI 32.2–49.7) versus women (24.6%, 19.2–30.6,  $\chi^2=10.4$ , *P* = 0.001) in both BD and MDD (not shown).

### Factors associated with suicidal acts overall

Among all patients with mood disorders, the factors that were present before intake at the study site and that were significantly associated with lifetime suicidal acts (Table 2A) included: family

**Table 1** Types of suicidal risks in 3284 patients with bipolar disorder type I, bipolar disorder type II or unipolar major depressive disorder

Ideation or behaviour	Prevalence (%) (95% CI)			Relative risk ratio (95% CI)	P-value ( $\chi^2$ or <i>t</i> )	Prevalence (%) (95% CI)			Relative risk ratio (95% CI)	P-value ( $\chi^2$ or <i>t</i> )
	BD-I ( <i>n</i> = 714)	BD-II ( <i>n</i> = 497)	All BD ( <i>n</i> = 1211)			MDD ( <i>n</i> = 2073)				
Ideation	25.2 (22.1–28.6)	35.0 (30.8–39.4)	29.2 (26.7–31.9)	0.72 (0.34–0.85)	0.007 (7.36)	17.3 (15.7–19.0)	1.69 (1.54–1.89)	<0.0001 (40.5)		
Attempts	20.9 (17.9–24.0)	15.9 (12.8–19.4)	18.8 (16.7–21.1)	1.31 (1.09–1.89)	0.07 (3.27)	4.78 (3.90–5.78)	3.93 (3.49–4.39)	<0.0001 (134)		
Suicides	1.82 (0.97–3.09)	1.61 (0.70–3.15)	1.73 (1.08–2.64)	1.13 (1.08–1.20)	0.07 (0.79)	0.48 (0.23–0.89)	3.60 (2.86–4.75)	0.004 (12.5)		
All acts (attempts + suicides)	22.7 (19.7–25.9)	17.5 (14.3–21.1)	20.6 (18.3–22.9)	1.30 (1.22–1.41)	0.07 (3.20)	5.26 (4.34–6.31)	3.92 (3.64–4.22)	<0.0001 (144)		
Percentage violent acts <sup>a</sup>	38.2 (30.9–46.0)	23.0 (14.6–33.2)	33.1 (27.4–39.2)	1.66 (1.53–1.81)	0.01 (6.04)	23.9 (16.2–33.0)	1.38 (1.30–1.49)	0.08 (3.08)		
Exposure (years)	16.3 (15.3–17.3)	18.0 (16.8–19.2)	17.0 (16.2–17.7)	0.91 (0.85–0.94)	0.03 (2.19)	11.7 (11.2–12.3)	1.45 (1.32–1.63)	<0.0001 (11.5)		
Attempts/suicides prevalence ratio	11.5	9.88	10.9	—	—	9.96	—	—		

Ideation or behaviour	Rates (%/year) (95% CI)			IRR (95% CI)	P-value (exact)
	BD-I	BD-II	All BD		
Ideation	1.55 (1.36–1.75)	1.92 (1.71–2.19)	1.72 (1.57–1.88)	0.80 (0.64–0.99)	0.02
Attempts	1.28 (1.10–1.47)	0.88 (0.71–1.08)	1.11 (0.98–1.24)	1.45 (1.10–1.93)	0.003
Suicides	0.11 (0.06–0.19)	0.09 (0.04–0.18)	0.10 (0.06–0.16)	1.25 (0.48–3.48)	0.32
All acts (attempts + suicides)	1.39 (1.21–1.59)	0.97 (0.79–1.17)	1.21 (1.08–1.35)	1.43 (1.10–1.88)	0.003
Attempts/suicides rate ratio	11.6	9.78	11.1	—	—

Ideation or behaviour	Rates (%/year) (95% CI)			IRR (95% CI)	P-value (exact)
	BD-I	MDD	All BD		
Ideation	1.55 (1.36–1.75)	1.47 (1.34–1.62)	1.72 (1.57–1.88)	1.17 (1.01–1.36)	0.02
Attempts	1.28 (1.10–1.47)	0.41 (0.33–0.49)	1.11 (0.98–1.24)	2.72 (2.14–3.48)	<0.0001
Suicides	0.11 (0.06–0.19)	0.04 (0.02–0.08)	0.10 (0.06–0.16)	2.48 (1.19–5.90)	0.008
All acts (attempts + suicides)	1.39 (1.21–1.59)	0.45 (0.37–0.54)	1.21 (1.08–1.35)	2.70 (2.15–3.41)	<0.0001
Attempts/suicides rate ratio	11.6	10.2	11.1	—	—

Lifetime prevalence in patients with bipolar disorder (BD, type I or II) or major depressive disorder (MDD) compared as relative risk ratios (with  $\chi^2$ ). Exposure years are compared by analysis of variance (*F*-score) and exposure-adjusted rates (percentage/year) are compared as incidence rate ratios (IRRs, with exact *P*-values). BD-I, BD type I; BD-II, BD type II.  
<sup>a</sup> Prevalence of violent acts is expressed as proportion of all suicidal acts (attempts + suicides).

**Table 2** Factors associated with suicidal acts in 3284 patients with mood disorders

Factors	Proportion or value (95% CI)		Risk ratio	Statistic ( $\chi^2$ or <i>t</i> )	P-value
	Suicidal acts ( <i>n</i> = 371; 11.3%)	No suicidal acts ( <i>n</i> = 2913; 88.7%)			
<b>A: Before intake</b>					
Gender (%)			1.03 <sup>a</sup>	0.05	0.82
Women	11.4 (10.1–12.8)	88.6 (87.2–89.9)			
Men	11.1 (9.38–13.1)	88.9 (86.9–90.6)			
Family history (%)					
Major affective	64.9 (59.0–70.4)	57.7 (55.4–60.0)	1.12	5.20	0.02
Bipolar disorder	32.3 (26.9–38.0)	19.6 (17.8–21.5)	1.65	23.7	<0.0001
Suicide	11.5 (8.12–15.6)	5.48 (4.54–6.55)	2.10	16.3	<0.0001
Marital status					
Married/widowed (%) <sup>b</sup>	51.0 (45.7–56.2)	63.1 (61.3–64.8)	0.81	19.9	<0.0001
Separated/divorced (%) <sup>c</sup>	17.9 (12.6–24.3)	8.21 (6.99–9.58)	2.18	18.4	<0.0001
Children	0.91 (0.75–1.07)	1.25 (1.19–1.31)	0.73	3.85	0.0001
Socioeconomic status					
Unemployed <sup>d</sup>	17.5 (12.7–23.3)	10.9 (9.49–12.5)	1.61	8.06	0.004
Socioeconomic rank <sup>e</sup>				7.20	0.03
High	12.9 (8.86–17.9)	87.1 (82.1–91.1)	1.00		
Medium	18.3 (15.3–21.6)	81.7 (78.4–84.7)	1.42		
Low	24.7 (16.5–34.5)	75.3 (65.5–83.5)	1.91		
Prior stressors					
Early abuse/trauma (%)	50.5 (40.2–60.8)	37.1 (32.5–41.9)	1.36	5.96	0.01
Age at parental loss (years)	25.1 (22.7–27.6)	29.2 (28.2–30.1)	0.86	3.04	0.002
Onset age $\leq$ 25 years (%)	59.0 (53.8–54.1)	38.4 (36.6–40.2)	1.54	57.9	<0.0001
Years from onset	12.8 (11.6–14.0)	10.5 (10.1–10.9)	1.22	3.71	0.0002
More than four depressions (%)	31.0 (25.8–36.6)	21.6 (19.6–23.7)	1.44	12.4	0.0004
Admission to hospital (%)	58.5 (51.1–65.6)	21.0 (18.4–23.7)	2.79	112	<0.0001
<b>B: At intake</b>					
Age $\leq$ 40 years (%)	58.8 (53.6–63.8)	48.1 (46.3–50.0)	1.22	14.9	0.0001
Diagnosis (%)					
MDD	5.40 (4.47–6.46)	94.6 (93.5–95.5)	3.96 <sup>f</sup>	195	<0.0001
BD	21.4 (19.1–23.8)	78.6 (76.2–80.9)			
BD-I	23.8 (20.7–27.1)	76.2 (72.9–79.3)	1.33 <sup>g</sup>	5.93	0.01
BD-II	17.9 (14.7–21.6)	82.1 (78.4–85.3)			
Comorbidity (%)					
ADHD					
Present (%)	41.2 (34.0–48.6)	30.9 (27.9–34.0)	1.33	7.47	0.006
ASRS-A score	33.9 (31.3–36.5)	29.6 (28.5–30.7)	1.15	3.12	0.002
Anxiety disorder	57.3 (48.1–66.1)	71.9 (68.3–75.4)	0.59	10.6	0.001
Any substance misuse	21.8 (17.4–26.3)	9.47 (8.42–10.6)	2.28	47.9	<0.0001
Alcohol misuse	20.4 (13.9–28.3)	10.5 (8.18–13.1)	1.94	10.2	0.001
Smoking	50.0 (42.8–57.2)	34.4 (31.5–37.5)	1.45	16.9	<0.0001
Mixed depression in MDD (%)	29.5 (20.3–40.2)	17.4 (15.7–19.2)	1.70	8.37	0.004
Manic switching in BD-I (%)	22.2 (16.8–28.4)	13.6 (11.3–16.2)	1.63	6.59	0.01
Temperament (TEMPS-A)					
Dysthymic	4.36 (3.83–4.89)	3.31 (3.11–3.52)	1.32	3.91	0.0001
Cyclothymic	6.65 (5.89–7.41)	5.17 (4.85–5.49)	1.29	3.66	0.0003
Irritable	2.07 (1.56–2.58)	1.52 (1.34–1.70)	1.36	2.32	0.02
Ratings at intake					
Depression (HRSD total $\geq$ 20; %)	46.5 (40.8–52.3)	39.9 (38.0–41.8)	1.17	4.83	0.03
Mood history (MDQ)	11.2 (10.1–12.2)	7.91 (7.36–8.46)	1.42	5.02	<0.0001
<b>C: During follow-up</b>					
Years at risk	4.79 (4.51–5.07)	2.72 (2.51–2.93)	1.76	6.26	<0.0001
Prospective percentage time spent ill					
Total morbidity	29.3 (25.2–33.5)	19.8 (17.9–21.6)	1.48	4.17	<0.0001
Depression	20.1 (17.0–23.2)	14.1 (12.9–15.4)	1.43	3.83	0.0001
(Hypo)mania	9.23 (7.35–11.1)	5.60 (4.56–6.64)	1.65	2.93	0.0003
Treatments given (%)					
Antidepressants	59.3 (52.3–66.0)	78.2 (76.1–80.2)	0.76	35.9	<0.0001
Mood stabilisers	23.0 (17.4–29.3)	8.25 (6.94–9.74)	2.79	44.4	<0.0001
Antipsychotics	45.0 (38.1–52.0)	22.3 (20.2–24.4)	2.03	50.8	<0.0001
Admission to hospital (%)	18.2 (12.9–24.6)	6.93 (5.38–8.77)	2.63	24.1	<0.0001

Comparisons are of proportions or values of measures among patients with versus those without suicidal behaviour (attempts or suicides); those without suicidal acts include patients with suicidal ideation only. Factors not significantly associated with suicidal behaviour included: education beyond high school, and having an anxious and hyperthymic temperament. MDD, major depressive disorder; BD, bipolar disorder; BD-I, BD type I; BD-II, BD type II; ASRS-A, Adult ADHD Self-Report Scale Part A; TEMPS-A, Temperament Evaluation of Memphis, Pisa, Paris and San Diego-Autoquestionnaire; HRSD, Hamilton Rating Scale for Depression; MDQ, Mood Disorder Questionnaire.

a. In suicidal women versus men.

b. Versus single or in long-term relationship.


c. Versus married/widowed.

d. Excludes students, homemakers, people who are disabled.

e. In low or medium versus high socioeconomic status.

f. In BD versus MDD.

g. In BD-I versus BD-II.



**Fig. 1** Relative risk factors associated with suicidal acts in bipolar disorder (BD) and major depressive disorder (MDD). (a) Relative risk of 31 factors highly significantly ( $P \leq 0.01$ ) associated with 371 patients with major mood disorders who were suicidal versus 2913 patients with major mood disorders who were non-suicidal. Factors are shown in descending rank order by risk ratio. (b) Comparisons of strength of association of these risk factors as the ratio of their prevalence or magnitude among patients who were suicidal/non-suicidal, separated by diagnosis (BD,  $n = 1211$ , light bars; versus MDD,  $n = 2073$ , dark bars), indicating which factors associated more with BD than MDD (upper cluster), which were similarly associated in both (middle cluster) and which were more associated with MDD than BD (lower cluster).

ADHD, attention-deficit hyperactivity disorder; BD-I, BD type I; BD-II, BD type II; MDQ, Mood Disorder Questionnaire; TEMPS, Temperament Evaluation of Memphis, Pisa, Paris and San Diego.

history of major affective disorder, BD or suicide; being unmarried or divorced and having fewer children; being unemployed and having a low socioeconomic status; having experienced early abuse or trauma, and relatively early loss of a parent; being younger at illness onset and with more years at risk; having had more than four prior depressions; and being admitted to hospital for psychiatric illness.

At intake assessment, the factors associated with lifetime risk of suicidal acts (Table 2B) included: older current age; BD (BD-I > BD-II) versus MDD diagnosis; co-occurring attention-deficit hyperactivity disorder (ADHD), especially with inattention (higher Adult ADHD Self-Report Scale Part A score); misuse of drugs or alcohol or smoking; lack of an anxiety disorder; mixed (hypomanic) symptoms in depressive episodes (in MDD or BD); switching into (hypo)mania in BD-I; higher ratings for dysthymic, cyclothymic or irritable temperament (Temperament Evaluation of Memphis, Pisa, Paris and San Diego-Autoquestionnaire ratings); somewhat higher Hamilton Rating Scale for Depression (HRSD) ratings; and higher ratings of lifetime BD-like symptoms (Mood Disorder Questionnaire score).

The factors associated with suicidal behaviour that were evaluated prospectively during follow-up (Table 2C) included: longer exposure time (years at risk), greater morbidity (as percentage time spent ill, depressed, or [hypo]manic) and psychiatric admission to hospital. In addition, although treatments were assigned clinically, it is noteworthy that antidepressant monotherapy was much less often given to patients who were suicidal, whereas use of mood stabilisers or antipsychotic drugs alone or with

antidepressants was much more prevalent among patients who were suicidal (Table 2C).

The risk factors that were most strongly associated with suicidal behaviour among the 3284 patients with major mood disorders (based on findings in Table 2) are shown in descending rank order in Fig. 1a. Additionally, associations (risk ratio) with suicidal behaviour among patients with BD and those with MDD are shown separately in Fig. 1b. Risk factors fell into three clusters: (a) those more associated with suicidal acts in patients with BD than those with MDD, (b) those similar in patients with BD and those with MDD, and (c) those more strongly associated among patients with MDD than those with BD.

### Multivariable logistic regression modelling

Factors preliminarily associated with suicide attempts or suicides among all patients with mood disorders (Table 2) were further tested by stepwise, multivariable logistic regression modelling. In order of strength of independent association with suicidal behaviour, the five significantly and independently associated factors ranked as follows: psychiatric admission to hospital, intake HRSD depression score  $\geq 20$ , BD diagnosis, onset age  $\leq 25$  years, and presence of mixed features in any illness episode (Supplementary Table A1).

### ROC analysis

We used factors sustained for association with suicidal acts in multivariable regression modelling for ROC analysis, based on regressing

**Table 3** Numbers of factors selectively associated with suicidal acts in 3284 patients with a major mood disorder

Factors ( <i>n</i> )	Distribution of risk factors						Bayesian analysis		
	Total ( <i>n</i> , %)	Suicidal ( <i>n</i> , %)	Non-suicidal ( <i>n</i> , %)	Odds ratio (95% CI)	$\chi^2$	<i>P</i> -value	Sensitivity (%)	Specificity (%)	Correctly classified (%)
0	559 (17.0)	19 (5.12)	540 (18.5)	1.00	—	—	100	0.00	11.3
1	1145 (34.9)	70 (18.9)	1075 (36.9)	1.85 (1.10–3.11)	5.44	0.02	94.9	18.5	27.2
2	966 (29.4)	115 (31.0)	851 (29.2)	3.84 (2.34–6.31)	28.1	<0.0001	76.0	55.4	57.8
3	451 (13.7)	103 (27.8)	348 (11.9)	8.41 (5.06–14.0)	67.6	<0.0001	45.0	84.6	80.2
4	140 (4.26)	54 (14.5)	86 (2.95)	17.8 (10.1–31.6)	98.1	<0.0001	17.3	96.6	87.6
5	23 (0.70)	10 (2.70)	13 (0.45)	21.9 (8.52–56.1)	41.1	<0.0001	2.70	95.6	88.6

Factors tested were: (a) diagnosis of bipolar disorder versus major depressive disorder, (b) intake age  $\leq 40$  years, (c) previous admission to hospital, (d) presence of mixed features, and (e) intake Hamilton Rating Scale for Depression total depression score  $> 18$ .

Bayesian sensitivity versus (100 – specificity) or true positive versus false positive rate (Supplementary Fig. A1). The computed AUC was 71.3% (95% CI 68.6–74.0), well above a chance association (50.0%). Optimal differentiation of patients with versus those without suicidal acts was associated with the presence of two factors, optimally yielding sensitivity of 76.0% and specificity of 55.4%, and correct classification of 57.8% of patients as suicidal or not (Table 3).

## Discussion

This study provides risks and annualised rates for patients with suicidal acts (attempts or suicides), relative risks of violent and non-violent suicidal acts, as well as estimated rates of suicidal ideation among 3284 similarly and consistently evaluated patients with major affective disorders at a single study site, with comparisons among the diagnostic types (Table 1). Overall, we found a lifetime risk for suicide ideation of about 68%, for suicide attempts of various severity of 30%, for suicide of 2.8%, and for all acts of 33% during an average exposure time of 14.3 years. The attempts/suicides ratio was similar in patients with BD (11.1) and those with MDD (10.2), and several times lower (greater lethality) than in the general population (at least 30).<sup>11</sup> The study also quantified associations of many demographic and clinical factors with suicidal acts (Table 2, Fig. 1a), as well as their relative likelihood among those diagnosed with BD or MDD (Fig. 1b).

As expected, indications of less conventionally successful social functioning were associated with greater risk of suicidal behaviour.<sup>12–14</sup> Such risk factors included being unmarried or divorced and consequently having fewer children, as well as lower socio-economic status and more unemployment. These factors are also independently associated with the presence of a mood disorder. Also associated with suicide and attempts were younger illness onset, longer time from onset to intake and older current age, which are all indicative of greater exposure or more years at risk.

Greater suicidal risk was more associated with BD than with MDD, and risk was similar among patients with BD-I and those with BD-II (Table 1). These findings accord with previous observations involving cases of MDD being unselected for illness severity.<sup>6,15</sup> Co-occurring morbidities associated with suicidal acts included ADHD and substance misuse as well as smoking. However, contrary to expectation,<sup>2,16,17</sup> co-occurring anxiety disorders were associated with lower risk of suicidal acts (Table 2B), perhaps by association with a lower degree of impulsivity.

In addition to a BD diagnosis, risk factors associated with suicidal behaviour included indices of greater illness severity: notably, time spent ill or depressed, multiple early recurrences of depression, psychiatric admission to hospital and co-occurring substance misuse (Table 2). Also strongly associated with suicidal behaviour was a history of BD or of suicide among first-degree family members. We also found higher ratings for certain types of

affective temperaments, particularly dysthymic and cyclothymic, among patients who were suicidal, as reported previously.<sup>18,19</sup> Particularly of note, the present findings also support a well-known, strong association of suicidal risk with mixed features, including patients diagnosed with either BD or MDD (Table 2).<sup>9,20–23</sup> Elements that may connect BD, mixed states and suicidal risk include aggressiveness and other mania-associated characteristics in both juveniles and adults.<sup>24,25</sup>

Violent methods of suicide or attempts were identified in 28.4% (95% CI 7.21–49.6) of all suicidal acts (attempts + suicides), more with a diagnosis of BD-I than BD-II or MDD (Table 1) and, as expected,<sup>26–28</sup> more with suicides than attempts, and more among men than women. The overall rate of violent acts is somewhat lower than recently reported rates that averaged 45.1% (95% CI 14.8–75.4; 36% for attempts *v.* 54.6% for suicides, 38.2% among women *v.* 55.3% in men), probably due to regional and cultural differences.<sup>26–28</sup>

Another interesting observation is that use of antidepressants alone was much less prevalent among patients who were suicidal than those who were non-suicidal, in contrast to greater use of antipsychotics and mood stabilisers alone or combined with antidepressants (Table 2). This clinical, nonrandomised choice of treatments reflects clinical practice at the study site, in which antidepressants were avoided in favour of antipsychotics or mood stabilisers in patients who are agitated or show mixed features, so as to avoid potentially increasing suicidal risk.

Five factors significantly and independently differentiated patients with versus those without suicidal acts based on multivariable logistic regression modelling: psychiatric admission to hospital, higher depression rating at intake, BD diagnosis, younger at onset and presence of mixed features in one or more illness episodes (Supplementary Table A1). Some of these factors may reflect greater illness severity. ROC analysis found that these risk factors differentiated patients who were suicidal from those who were non-suicidal with an AUC of 71.3%, and that any two of these independently associated factors optimally differentiated those who were suicidal from those who were non-suicidal with maximal sensitivity and specificity (Table 3; Supplementary Fig. A1). Risk factors may be associated with suicidal risk but would be of limited value in predicting suicidal behaviour for particular patients and times. Notably, they must be combined with attention to clinical conditions and behaviours that may be associated with planning for suicide, including verbal statements, organising personal and business affairs or giving away valued possessions.

## Limitations

These findings include factors derived from clinical histories and so may include errors of ascertainment. However, these should be similarly likely across the diagnostic groups compared. Observations made in a mood disorder centre might introduce biases, such as bias related to illness severity, although most study patients were

referred by primary care clinicians or were self-referred. In addition, the bivariate testing of large numbers of risk factors for association with suicidal status risks finding chance associations. Rather than making Bonferroni-type adjustments of significance, factors with strong preliminary associations with suicidal behaviour were further tested with multivariable regression modelling, as well as by comparing the magnitude of observed bivariate associations overall and between diagnostic groups.

The present findings test and extend extensive previous research on risk factors for suicidal behavior among subjects with a major mood disorder, as reviewed in the Appendix, and add quantitative assessments of the strength and independence of the association of risk factors with suicidal status, comparing them among specific diagnostic groups. Factors with especially strong associations with suicidal acts ( $P < 0.0001$ ; Table 2) include: BD diagnosis, familial BD or suicide, being unmarried, early illness-onset, being admitted to hospital, substance abuse or smoking, years of illness, and prospective proportion of time ill or depressed, as well as less use of antidepressant treatment alone and greater use of mood-stabilising or antipsychotic medicines. The findings should contribute to improving earlier identification of suicidal risk, supporting preventive interventions, and improving the treatment and prognosis of patients with mood disorders at particularly high suicidal risk.

**Ross J. Baldessarini**, Director, International Consortium for Research on Mood and Psychotic Disorders, McLean Hospital; and Professor, Department of Psychiatry, Harvard Medical School, USA; **Leonardo Tondo**, Investigator, International Consortium for Research on Mood and Psychotic Disorders, McLean Hospital; Research Associate, Department of Psychiatry, Harvard Medical School, USA; Director, Lucio Bini Mood Disorders Centers, Italy; and Investigator, Centre for Affective Disorders, Department of Psychological Medicine, Institute of Psychiatry, King's College, London, UK; **Marco Pinna**, Investigator, Lucio Bini Mood Disorders Centers, Italy; **Nicholas Nuñez**, Investigator, Department of Psychiatry, Queen's University School of Medicine, Canada; **Gustavo H. Vázquez**, Investigator, International Consortium for Research on Mood and Psychotic Disorders, McLean Hospital, USA; and Professor, Department of Psychiatry, Queen's University School of Medicine, Canada

**Correspondence:** R. J. Baldessarini, Mailman Research Center 3, McLean Hospital, 115 Mill Street, Belmont, MA 02478-9106, USA. Email: [rbaldessarini@mclean.harvard.edu](mailto:rbaldessarini@mclean.harvard.edu)

First received 5 Mar 2019, final revision 11 May 2019, accepted 3 Jun 2019

## Supplementary material

Supplementary material is available online at <https://doi.org/10.1192/bjp.2019.167>

## Funding

This work was supported by a grant from the Bruce J. Anderson Foundation and by the McLean Private Donors Psychiatric Research Fund (to R.J.B.), as well as by a grant from the Aretaeus Foundation of Rome (to L.T.).

## References

- Harris EC, Barraclough B. Suicide as an outcome for mental disorders: a meta-analysis. *Br J Psychiatry* 1997; **170**: 205–28.
- Hawton K, Sutton L, Haw C, Sinclair J, Harriss L. Suicide and attempted suicide in bipolar disorder: systematic review of risk factors. *J Clin Psychiatry* 2005; **66**: 693–704.
- Schaffer A, Isometsä ET, Tondo L, Moreno D, Turecki G, Reis C, et al. International Society for Bipolar Disorders Task Force on Suicide: meta-analyses and meta-regression of correlates of suicide attempts and suicide deaths in bipolar disorder. *Bipolar Disord* 2015; **17**: 1–16.
- Tondo L, Baldessarini RJ. Suicide in bipolar disorder. In *The Bipolar Book: History, Neurobiology, and Treatment* (eds Yildiz A, Nemeroff C, Ruiz P): 509–28. Oxford University Press, 2015.
- Tondo L, Pompili M, Forte A, Baldessarini RJ. Suicide attempts in bipolar disorders: comprehensive review of 101 reports. *Acta Psychiatr Scand* 2016; **133**: 174–86.
- Tondo L, Lepri B, Baldessarini RJ. Risks of suicidal ideation, attempts and suicides among 2826 men and women with types I and II bipolar, and recurrent major depressive disorders. *Acta Psychiatr Scand* 2007; **116**: 419–28.
- Tondo L, Baldessarini RJ, Vázquez G, Lepri B, Visioli C. Clinical responses to antidepressants among 1036 acutely depressed patients with bipolar or unipolar major affective disorders. *Acta Psychiatr Scand* 2013; **127**: 355–64.
- Tondo L, Visioli C, Preti A, Baldessarini RJ. Bipolar disorders following initial depression: Modeling predictive clinical factors. *J Affect Disord* 2014; **167**: 44–9.
- Tondo L, Vázquez GH, Pinna M, Vaccotto PA, Baldessarini RJ. Characteristics of depressive and bipolar patients with mixed features. *Acta Psychiatr Scand* 2018; **138**: 243–52.
- Italian Ministry of Economics and Finances. Autorizzazione generale al trattamento dei dati personali effettuato per scopi di ricerca scientifica (Autorizzazione n. 9/2014) [General authorization to treatment of personal data for scientific research (Authorisation no. 9/2014)]. *Gazzetta Ufficiale Serie Generale* n. 301, 30 December 2014 ([http://www.gazzettaufficiale.it/atto/serie\\_generale/caricaDettaglioAtto/originario;jsessionid=k0I0R58vW5a9RX4mnszi2A\\_.ntc-as5-guri2b?atto.dataPubblicazioneGazzetta=2014-12-30&atto.codiceRedazionale=14A09919&elenco30giorni=false](http://www.gazzettaufficiale.it/atto/serie_generale/caricaDettaglioAtto/originario;jsessionid=k0I0R58vW5a9RX4mnszi2A_.ntc-as5-guri2b?atto.dataPubblicazioneGazzetta=2014-12-30&atto.codiceRedazionale=14A09919&elenco30giorni=false)).
- American Foundation for Suicide Prevention (AFSP). Suicide statistics. AFSP, 2017 ([www.afsp.org/AboutSuicide/Suicide-Statistics](http://www.afsp.org/AboutSuicide/Suicide-Statistics)).
- Hansson C, Joas E, Pålsson E, Hawton K, Runeson B, Landén M. Risk factors for suicide in bipolar disorder: a cohort study of 12,850 patients. *Acta Psychiatr Scand* 2018; **138**: 456–63.
- Chaudhury SR, Grunebaum MF, Galfalvy HC, Burke AK, Sher L, Parsey RV, et al. Does first episode polarity predict risk for suicide attempt in bipolar disorder? *J Affect Disord* 2007; **104**: 245–50.
- Bellivier F, Yon L, Luquiens A, Azorin JM, Bertsch J, Gerard S, et al. Suicidal attempts in bipolar disorder: results from an observational study (EMBLEM). *Bipolar Disord* 2011; **13**: 377–86.
- Bottlender R, Jäger M, Strauss A, Möller HJ. Suicidality in bipolar compared to unipolar depressed inpatients. *Eur Arch Psychiatry Clin Neurosci* 2000; **250**: 257–61.
- Isometsä E. Suicidal behavior in mood disorders—who, when, and why? *Can J Psychiatry* 2014; **59**: 120–30.
- Sánchez-Gistau V, Colom F, Mane A, Romero S, Suñer G, Vieta E. Atypical depression is associated with suicide attempt in bipolar disorder. *Acta Psychiatr Scand* 2009; **120**: 30–6.
- Baldessarini RJ, Innamorati M, Erbutto D, Serafini G, Fiorillo A, Amore M, et al. Differential associations of affective temperaments and diagnosis of major affective disorders with suicidal behavior. *J Affect Disord* 2017; **210**: 19–21.
- Vázquez GH, Gonda X, Löllich M, Tondo L, Baldessarini RJ. Suicidal risk and affective temperaments evaluated with TEMPS-A. *Harv Rev Psychiatry* 2018; **26**: 8–18.
- Saunders KE, Hawton K. Clinical assessment and crisis intervention for the suicidal bipolar disorder patient. *Bipolar Disord* 2013; **15**: 575–83.
- Saunders KE, Hawton K. Suicidal behaviour in bipolar disorder: understanding the role of affective states. *Bipolar Disord* 2015; **17**: 24–6.
- Akiskal HS, Benazzi F. Psychopathologic correlates of suicidal ideation in major depressive outpatients: is it all due to unrecognized (bipolar) depressive mixed states? *Psychopathology* 2005; **38**: 273–80.
- Vázquez GH, Löllich M, Cabrera C, Jokic R, Kolar D, Tondo L, et al. Mixed symptoms in mood disorders: a systematic review. *J Affect Disord* 2018; **250**: 756–60.
- Diler RS, Goldstein TR, Hafeman D, Merranko J, Liao F, Goldstein BI, et al. Distinguishing bipolar depression from unipolar depression in youth: preliminary findings. *J Child Adolesc Psychopharmacol* 2017; **27**: 310–9.
- Perugi G, Angst J, Azorin JM, Bowden CL, Mosolov S, Reis J, et al. Mixed features in patients with a major depressive episode: the BRIDGE-II-MIX study. *J Clin Psychiatry* 2015; **76**: e351–8.
- Bradvik L. Violent and nonviolent methods of suicide: different patterns may be found in men and women with severe depression. *Arch Suicide Res* 2007; **11**: 255–64.
- Misson H, Mathieu F, Jollant F, Yon L, Guillaume S, Parmentier C, et al. Factor analyses of the Suicidal Intent Scale (SIS) and the Risk-Rescue Rating Scale (RRRS): toward the identification of homogeneous subgroups of suicidal behaviors. *J Affect Disord* 2010; **121**: 80–7.
- Forsman J, Masterman T, Ahlner J, Isacson G, Hedstrom AK. Selective serotonin re-uptake inhibitors and the risk of violent suicide: a nationwide post-mortem study. *Eur J Clin Pharmacol* 2019; **75**: 393–400.

