

P01.18

Antidepressants of the different chemical groups: clinical and neurophysiological correlations in treating depression

E. Grigorieva*, A. Dyakonov. *Yaroslavl State Medical Academy, Department of Psychiatry, Russia*

Developing options of the psychopharmacology require specialists to conduct goal-oriented and adequate therapy.

The objective of the research was a comparative clinical and neurophysiological study of the action of amitriptyline, tyaneptine, fluoxetine and moclobemide, treating depressive disorders with the dominant affect of anguish.

Totally, 107 depressed patients, aged 20–40, with depressive episode (38,3%), recurrent (53,3%) and bipolar (8,4%) affective disorders were clinically examined. Depression severity was assessed by clinical examination as well as by the administration of HAM-D Rating Scale and CGI Scale prior to and on the 10th, 20th and 40th days of treatment. Brain-wave mapping was recorded before the therapy and on its 20th day. All patients were divided onto 4 groups: those, who were treated with amitriptyline (n=25), tyaneptine (n=28), fluoxetine (n=32) and moclobemide (n=22). Daily dosages of amitriptyline (150–250 mg), tyaneptine (37.5–50 mg), fluoxetine (40–60 mg) and moclobemide (450–600 mg) depended on the depression severity (moderate or severe levels). Control group consisted of 25 healthy people.

Results of the study showed: when registering brain bio-electric activity of depressed patients prior to the treatment in comparison with the healthy group we have observed zones of "increased" activity in the right temporal fields and zones of "decreased" activity in the left temporal fields. Changes, occurring in these zones in response to the treatment, were more stable. Affecting the brain electric impulses all used agents demonstrated a similarity (core pattern) as well as the differences. All antidepressants reduced the activation of the right temporal zone and increased the activation of the left temporal zone. As a specific feature of the action, fluoxetine and moclobemide produced more significant increasing of the left temporal zone's activation. Amitriptyline caused an expanding of the activation zone from the left temporal fields to parietal and occipital fields. Tyaneptine produced migrating zones of activation.

P02. Antipsychotics**P02.01**

Coadministration of clozapine and amisulpride in psychotic patients

M. Ziegenbein*, O. Rosenthal, P. Garlipp. *Hannover Medical School, Department of Psychiatry, Germany*

Clozapine was the first atypical antipsychotic drug to be introduced into clinical use in several European countries. Clozapine treatment is associated with wide side-effects. The mechanisms underlying these side effects are still unknown. The wide side-effects are very often responsible for substantial compliance problems. Previous case reports have shown before that the coadministration of amisulpride and clozapine might be useful to reduce side-effects. We performed an open clinical study to gather more experience in the clinical efficiency of the coadministration of the two drugs. Nine psychotic inpatients (6 female, 3 male; mean age 42,8 [plusminus] 12,3) participated and were diagnosed according to DSM-IV criteria. The clozapine treatment was associated with side-effects and persisting psychotic symptoms. Under addition of amisulpride we found very low dosages of clozapine to be sufficient to obtain effective clozapine concentrations. Secondly,

we found a low incidence of side effects. All patients were in clinical remission. Our results indicate that amisulpride addition to clozapine is highly effective in reducing psychotic symptomatology and side-effects. This might be due to additive effects of the two drugs and/or metabolic interaction.

P02.02

Characteristics of acute schizophrenic patients on atypical antipsychotics

V.P. Kontaxakis*, B.J. Havaki-Kontaxaki, C.T. Kollias, M.M. Margariti, S.S. Stamouli, G.N. Christodoulou. *Department of Psychiatry, University of Athens, Eginition Hospital, Athens, Greece*

Objective: The aim of this study is to reveal the differential characteristics of acute schizophrenic inpatients on atypical antipsychotics.

Methods: The subjects of this study were sixty-three schizophrenic patients consecutively admitted at Eginition Hospital, Athens. All patients were assessed on admission using the Positive and Negative Syndrome Scale (PANSS). Patients' case notes analysis was performed surveying antipsychotic drugs' prescribing on the first week after their admission. Twenty-two patients were on atypical antipsychotics and forty-one patients were on conventional antipsychotics. Patients on atypical antipsychotics (AA) were compared with those on conventional antipsychotics (CA) in many social-demographic, clinical and psychopathological parameters. **Results:** There were no statistically significant differences between schizophrenic patients on AA and those on CA regarding age, sex, family status, employment status, duration of illness, PANSS-total score, PANSS- positive subscale score, PANSS-negative subscale score, PANSS-general psychopathology subscale score. Schizophrenic patients on AA scored lower on the item G14 of the PANSS (poor impulse control, 1.5 vs 2.2, U=168, p=0.08, statistically significant trend).

Conclusion: Acute schizophrenic inpatients on atypical antipsychotics were differentiated from those on conventional antipsychotics in that they had a better impulse control.

P02.03

Cognitive function in stable outpatients switched to ziprasidone

P.D. Harvey¹, H.Y. Meltzer², S.J. Romano³. ¹Mount Sinai School of Medicine, New York; ²Psychiatric Hospital at Vanderbilt, Vanderbilt University Medical Center; ³Pfizer, Inc., New York, USA

Objectives: To assess cognitive function were assessed in stable outpatients switched to ziprasidone from conventional or other novel antipsychotic agents.

Methods: 3 separate, 6-week multicenter trials assessed the effects of switching patients from conventional antipsychotics (n=108), olanzapine (n=104), or risperidone (n=58) to ziprasidone. Patients were randomly switched to ziprasidone 40–160 mg/day by 1 of 3 schedules for discontinuing their previous medication. Just before switching, and weekly during ziprasidone treatment, they were evaluated using a battery of assessments of working and secondary memory, vigilance, visuo-motor speed, verbal fluency, and executive functioning.

Results: Patients switched to ziprasidone manifested wide-ranging improvements in cognitive function. Significant (P<0.05 by ANCOVA) improvements were seen at endpoint (LOCF) in memory, vigilance (in patients switched from conventional antipsychotics or risperidone), executive function (in patients switched from conventional antipsychotics or risperidone), and verbal fluency.