

A SHORT PROOF OF THE FACTOR THEOREM FOR FINITE GRAPHS

W. T. TUTTE

We define a *graph* as a set V of objects called *vertices* together with a set E of objects called *edges*, the two sets having no common element. With each edge there are associated just two vertices, called its *ends*. We say that an edge *joins* its ends. Two vertices may be joined by more than one edge.

A *subgraph* G' of a graph G is a graph whose edges and vertices are edges and vertices respectively of G and in which each edge has the same ends as in G . If S is any set of vertices of G we denote by G_S the subgraph of G whose vertices are the vertices of G not in S and whose edges are the edges of G not having an element of S as an end.

A graph is *finite* if V and E are both finite and *infinite* otherwise. In this paper we consider only finite graphs.

Suppose given a finite graph G . For $a \in V$ and $A \in E$ we write $e(A, a) = 1$ if a is an end of A and $e(A, a) = 0$ otherwise. Let f be a function which associates with each vertex a of G a unique positive integer $f(a)$. We say that G is *f-soluble* for a given f if to each $A \in E$ we can assign a non-negative integer $g(A)$ such that

$$(1) \quad \sum_A e(A, a) g(A) = f(a)$$

for each $a \in V$. If E is null but V is not null, we consider that G is not f -soluble for any f . We ignore the case in which V and E are both null (when G is the *null graph*).

It may be possible to solve (1) so that $g(A) = 0$ or 1 for each A . Then we call the subgraph of G whose vertices are the vertices of G and whose edges are those edges A of G for which $g(A) = 1$ an *f-factor* of G . Thus an f -factor of G is a subgraph of G such that each $a \in V$ is a vertex of the subgraph and an end of just $f(a)$ edges of the subgraph.

If n is any positive integer we define an *n-factor* of G as an f -factor such that $f(a) = n$ for each a .

Necessary and sufficient conditions are known for f -solubility, for the existence of an f -factor and for the existence of a 1-factor. We state these as Theorems *A*, *B*, and *C* after a few preliminary definitions.

The *degree* $d(a)$ of a vertex a of G is the number of edges of G having a as an end. If $S \subseteq V$ and $a \in V - S$ we denote the degree of a in G_S by $d_S(a)$.

Suppose $S \subseteq V$. We write $\alpha(S)$ for the number of vertices of S . The graph G_S is uniquely decomposable into disjoint connected parts which we call *components*. (Hassler Whitney **(6)** uses the term *connected pieces*, and König

Received February 26, 1953; in revised form December 3, 1953.

(2) *zusammenhängende Bestandteile.*) We write $h_u(S)$ for the number of components of G_S for which the number of vertices is odd. We write $T(S)$ for the set of vertices of $V - S$ which are joined only to vertices of S .

We denote by $q(S)$ the number of components C of G_S for which there is more than one vertex and

$$(2) \quad \sum_{a \in C} f(a) \equiv 1 \pmod{2}.$$

Here we write $a \in C$ to denote that a is a vertex of C .

Now suppose $T \subseteq V - S$. If C is a component of $G_{S \cup T}$ we denote by $v(C)$ the number of edges of G having one end a vertex of C and the other an element of T . We denote by $q(S, T)$ the number of components C of $G_{S \cup T}$ such that

$$(3) \quad v(C) + \sum_{a \in C} f(a) \equiv 1 \pmod{2}.$$

THEOREM A. *G is without a 1-factor if and only if there is a subset S of V such that*

$$(4) \quad h_u(S) > \alpha(S).$$

THEOREM B. *G is not f -soluble if and only if there is a subset S of V such that*

$$(5) \quad \sum_{a \in S} f(a) < q(S) + \sum_{c \in T(S)} f(c).$$

THEOREM C. *G is without an f -factor if and only if there is a subset S of V and a subset T of $V - S$ such that*

$$(6) \quad \sum_{a \in S} f(a) < q(S, T) + \sum_{c \in T} (f(c) - d_s(c)).$$

A short proof of Theorem A has been given by the author (4). Maunsell (3) has improved it by substituting a piece of elementary graph theory for an appeal to the theory of determinants. Theorem B is readily deducible from Theorem C; details are given in (5). However, proofs of Theorem C, even in the special case dealing with n -factors, have hitherto been long and complicated (1; 5). In this paper we present a comparatively short argument whereby Theorem C is deduced as a consequence of Theorem A.

Deduction of Theorem C from Theorem A. Suppose first that G has a vertex a such that $d(a) < f(a)$. Then G can have no f -factor. Moreover (6) is satisfied with $S = 0$ and $T = \{a\}$. Thus Theorem C is trivially true in this case.

In the remaining case we have $d(a) \geq f(a)$ for each $a \in V$. We write $s(a) = d(a) - f(a)$.


Given any sufficiently large set Q we define a graph G' whose vertices are elements of Q in the following way. With each $c \in V$ we associate $d(c)$ distinct elements c_A of Q , one for each edge A of G such that $e(A, c) = 1$, and $s(c)$ other distinct elements $c(1), c(2), \dots, c(s(c))$ of Q . We denote the sets of the $d(c)$ elements c_A and the $s(c)$ elements $c(i)$ by $X(c)$ and $Y(c)$ respectively. We

postulate that the two sets $X(c) \cup Y(c)$ defined for two distinct elements c of V shall have no common element. The set V' of vertices of G' is given by

$$(7) \quad V' = \bigcup_{c \in V} (X(c) \cup Y(c)).$$

For any edge A of G , with ends x and y say, we postulate that G' has just one edge joining x_A and y_A . We denote this also by the symbol A . We further postulate that for each $c \in V$ each element of $X(c)$ is joined to each member of $Y(c)$ by just one edge of G' , and that G' has no edges other than those required by these two rules.

For each $c \in V$, the elements of $X(c) \cup Y(c)$ and the edges of G' joining them constitute a subgraph, $St(c)$, of G' , which we call the *star-graph of c in G'* . $St(c)$ is connected if $s(c) > 0$, and in the case $s(c) = 0$ only if $d(c) = f(c) = 1$. The diagram shows a star-graph $St(c)$ for the case $d(c) = 4$ and $f(c) = 2$. (The edges $A B C$ and D in this diagram do not belong to $St(c)$.)



LEMMA. G has an f -factor if and only if G' has a 1-factor.

Proof. If G has an f -factor let F be its set of edges and let F' be the set of edges of G' denoted by the same letters. For each $c \in V$ we adjoin to F' exactly $s(c)$ edges joining the $s(c)$ vertices of $Y(c)$ to the $s(c)$ vertices of $X(c)$ which are not ends of edges of F' . By the definition of G' we can do this without introducing into F' two edges with a common end. We thus construct a 1-factor of G' .

Conversely suppose G' has a 1-factor whose set of edges is H . Let H_0 be the set of edges of H whose two ends are vertices of distinct star-graphs $St(c)$. For each $c \in V$ just $s(c)$ elements of H have an end in $Y(c)$ and therefore just $d(c) - s(c) = f(c)$ elements of H_0 have an end in $X(c)$. It follows that the edges of G corresponding to the members of H_0 define an f -factor of G .

A subset W of V' will be called *simple* if it satisfies the following conditions for each $a \in V$:

- (i) If $X(a) \cap W \neq 0$ then $X(a) \subseteq W$,
- (ii) If $Y(a) \cap W \neq 0$ then $Y(a) \subseteq W$,
- (iii) At most one of $X(a)$ and $Y(a)$ is a subset of W .

Condition (iii) implies that $X(a)$ cannot be a subset of W when $Y(a)$ is the null set, i.e., when $d(a) = f(a)$.

Consider any simple subset W of V' . We write S and T for the sets of vertices c of G such that $X(c) \subseteq W$ and $Y(c) \subseteq W$ respectively. The sets S and T are disjoint. We have

$$(8) \quad \alpha(W) = \sum_{c \in T} (d(c) - f(c)) + \sum_{a \in S} d(a).$$

Let H be any component of G'_W .

It may happen that H has just one vertex, which is of the form c_A . Then $c \in T$ and the end of A in G other than c belongs to S . The number of such components H is the number of edges A of G having one end in S and the other in T , that is

$$\sum_{c \in T} (d(c) - d_S(c)).$$

Another possibility is that H has just one vertex, which is of the form $c(i)$. The number of such components is

$$\sum_{a \in S} (d(a) - f(a)).$$

In the remaining case, H has at least one edge. If H has no edge in common with one of the star-graphs $St(a)$ it must consist of a single edge with its two ends. Then the number of vertices of H is even. If H has an edge in common with $St(a)$ then $Y(a) \neq 0$ and so $St(a)$ is connected. Moreover $St(a)$ is then a subgraph of H . A component of G'_W having a connected star-graph $St(a)$ with at least one edge as a subgraph will be called *large*.

Suppose H is large. Let M be the set of all vertices a of G such that $St(a)$ is a connected subgraph of H with at least one edge. Then $M \subseteq V - (S \cup T)$. H is made up of these star-graphs $St(a)$, a set N of edges which link them to form a connected graph H_0 and a set P of edges having one end a vertex of H_0 and one end a vertex c_A such that $c \in T$. Clearly M is the set of vertices of a component $K(H)$ of $G_{S \cup T}$. We may think of $K(H)$ as derived from H_0 by shrinking each of the star-graphs $St(a)$, $a \in M$, to a single vertex. Conversely suppose K is any component of $G_{S \cup T}$. If c is a vertex of K then $Y(c) \neq 0$ since $c \notin T$ and therefore $St(c)$ is connected and has at least one edge. This star-graph is a subgraph of a large component H of G'_W and we must have $K = K(H)$.

For a large component H of G'_W having just n vertices

$$\begin{aligned} n &= \sum_{a \in K(H)} \{d(a) + (d(a) - f(a))\} + v(K(H)) \\ &\equiv \sum_{a \in K(H)} f(a) + v(K(H)) \pmod{2}. \end{aligned}$$

Hence the number of large components of G'_w for which the number of vertices is odd is $q(S, T)$.

Using (8) we obtain the formulae

$$(9) \quad h_u(W) = q(S, T) + \sum_{a \in S} (d(a) - f(a)) + \sum_{c \in T} (d(c) - d_S(c)),$$

$$(10) \quad \begin{aligned} h_u(W) - \alpha(W) \\ = q(S, T) - \sum_{a \in S} f(a) + \sum_{c \in T} (f(c) - d_S(c)). \end{aligned}$$

The quantities on the left in these equations are defined in terms of G' , those on the right in terms of G .

Suppose there are disjoint subsets S and T of V satisfying (6). Select two such subsets so that $\alpha(S)$ has the least possible value. Assume that $f(b) = d(b)$ for some $b \in S$. If we replace S by $S - \{b\}$ and T by $T \cup \{b\}$ inequality (6) will remain valid, for with at most $d(b)$ exceptions the numbers $v(C)$ associated with the components of $G_{S \cup T}$ are unaltered. This contradicts the definition of S . Hence $f(b) < d(b)$ for each $b \in S$. Let W be the union of the sets $X(a)$ such that $a \in S$ and the sets $Y(c)$ such that $c \in T$. Then W is simple since $Y(c)$ is non-null when $X(c) \subseteq W$. It follows from (10) that $h_u(W) > \alpha(W)$ in G' . Hence G' has no 1-factor, by Theorem A. Hence G has no f -factor, by the Lemma.

Conversely suppose G has no f -factor. Then by Theorem A and the Lemma there is a set W of vertices of G' such that $h_u(W) > \alpha(W)$. Choose such a W so that $\alpha(W)$ has the least possible value.

Suppose there exists $a \in V$ such that $Y(a) \cap W \neq 0$ and $Y(a) \cap (V' - W) \neq 0$. Write $Z = W - (Y(a) \cap W)$. Then G'_w and G'_z differ in one component only, provided that $X(a)$ is not a subset of W , since the members of $Y(a)$ are all joined to the same vertices of G' . If $X(a) \subseteq W$ then each component of G'_w is a component of G'_z . In either case we have $h_u(Z) \geq h_u(W) - 1$ and $\alpha(Z) \leq \alpha(W) - 1$. Hence $h_u(Z) - \alpha(Z) \geq h_u(W) - \alpha(W)$, contrary to the definition of W . We deduce that $Y(a) \subseteq W$ if $Y(a) \cap W \neq 0$.

Suppose next that $X(a) \cap W \neq 0$. Choose $b \in X(a) \cap W$. Write $Z = W - \{b\}$. There is at most one component of G'_w which has a vertex not a member of $Y(a)$ joined to b in G' . Hence if $Y(a)$ is contained in W the numbers $h_u(Z)$ and $h_u(W)$ can differ by at most one. Then $h_u(Z) \geq h_u(W) - 1$, $\alpha(Z) = \alpha(W) - 1$ and therefore $h_u(Z) - \alpha(Z) \geq h_u(W) - \alpha(W)$. This contradicts the definition of W . We deduce that, for the case $X(a) \cap W \neq 0$, $Y(a)$ is not a subset of W and therefore $Y(a) \cap W = 0$ by the result of the preceding paragraph. This proves that $X(a)$ and $Y(a)$ cannot both be subsets of W , since $X(a)$ is never null. ($d(a) \geq f(a) > 0$.)

Suppose both $X(a) \cap W$ and $X(a) \cap (V' - W)$ are non-null. We choose $b \in X(a) \cap W$ and write $Z = W - \{b\}$ as before. Since $Y(a) \cap W = 0$ all the vertices of $Y(a)$ belong to one component of G'_w , for each is joined in G' to each vertex of $X(a) \cap (V' - W)$. But there is at most one component of

G'_W which has a vertex not a member of $Y(a)$ joined to b in G' . Hence with at most two exceptions the components of G'_W are components of G'_Z . Accordingly

$$\begin{aligned} h_u(Z) &\geq h_u(W) - 2, \\ h_u(Z) - \alpha(Z) &\geq h_u(W) - \alpha(W) - 1. \end{aligned}$$

But $h_u(Z)$ is by definition the number of components of G'_Z having an odd number of vertices. Hence

$$h_u(Z) + \alpha(Z) \equiv \alpha(V') \pmod{2}$$

and similarly

$$h_u(W) + \alpha(W) \equiv \alpha(V') \pmod{2}.$$

We may write these results as

$$h_u(Z) - \alpha(Z) \equiv \alpha(V') \equiv h_u(W) - \alpha(W) \pmod{2}.$$

Hence $h_u(Z) - \alpha(Z) \geq h_u(W) - \alpha(W)$ and so the definition of W is contradicted. We deduce that $X(a) \subseteq W$ if $X(a) \cap W \neq \emptyset$.

We have now proved that W is simple. We define S and T in terms of W as before. Using (10) we find that S and T satisfy (6).

This completes the proof of Theorem C.

REFERENCES

1. H. B. Belck, *Reguläre Faktoren von Graphen*, J. Reine Angew. Math., 188 (1950), 228–252.
2. D. König, *Theorie der endlichen und unendlichen Graphen* (Leipzig, 1936).
3. F. G. Maunsell, *A note on Tutte's paper "The factorization of linear graphs,"* J. London Math. Soc., 27 (1952), 127–128.
4. W. T. Tutte, *The factorization of linear graphs*, J. London Math. Soc., 22 (1947), 107–111.
5. ———, *The factors of graphs*, Can. J. Math., 4 (1952), 314–328.
6. Hassler Whitney, *Non-separable and planar graphs*, Trans. Amer. Math. Soc., 34 (1932), 339–362.

University of Toronto