

Relevant associations of the glucokinase regulatory protein/glucokinase gene variation with TAG concentrations in a high-cardiovascular risk population: modulation by the Mediterranean diet

Mercedes Sotos-Prieto^{1,2,3}, Marisa Guillén¹, Jose Vicente Sorli^{1,2}, Olga Portolés¹, Patricia Guillem-Saiz^{1,2}, Jose Ignacio Gonzalez^{1,2}, Lu Qi³ and Dolores Corella^{1,2*}

¹Genetic and Molecular Epidemiology Unit, Department of Preventive Medicine and Public Health, School of Medicine, University of Valencia, Blasco Ibañez, 15. 46010- Valencia, Spain

²CIBER Fisiopatología de la Obesidad y Nutrición, Valencia, Spain

³Department of Nutrition and Epidemiology, Harvard School of Public Health, Boston, MA, USA

(Submitted 13 June 2011 – Final revision received 8 February 2012 – Accepted 8 February 2012 – First published online 13 April 2012)

Abstract

The SNP rs1260326 (P446L) and rs1799884 (–30G > A) for the glucokinase regulatory protein (*GCKR*) and glucokinase (*GCK*) genes, respectively, have been associated with opposing effects on TAG and glucose concentrations. However, their genetic modulation by diet (dietary patterns or foods) remains to be investigated. We studied 945 high-cardiovascular risk subjects aged 67 (SD 6) years who participated in the PREvención con DIeta MEDiterránea-Valencia Study. Demographic, clinical, biochemical and genetic data were obtained. Adherence to the Mediterranean diet (MD) and food intake were measured by validated questionnaires. Carriers of the L allele of *GCKR* had significantly higher TAG concentrations (PP: 1.34 (SD 0.05) mmol/l *v.* PL + LL: 1.54 (SD 0.03) mmol/l; *P*=0.014) and LL carriers had lower glucose concentrations (PL + PP: 6.85 (SD 0.08) mmol/l *v.* LL: 6.40 (SD 0.16) mmol/l; *P*=0.032) after multivariate adjustment. Conversely, homozygous subjects for the variant allele (A) in the *GCK* gene had significantly lower TAG (GG + GA: 1.48 (SD 0.03) mmol/l *v.* AA: 1.17 (SD 0.18) mmol/l; *P*=0.033) and a higher risk of diabetes (OR 3.3, 95% CI 1.2, 9.2). Combined effects for both SNP increased TAG concentrations by 37% (*P*=0.033). Adherence to the MD modulated the effects of *GCKR* polymorphism on TAG: subjects with genetic risk had lower TAG (L-allele carriers; PP: 1.48 (SD 0.14) mmol/l *v.* PL + LL: 1.51 (SD 0.08) mmol/l; *P*=0.917) compared with those with a higher adherence. Analysis of the joint effects of the *GCKR* and individual food items identified significant associations (olive oil (*P*=0.035), vegetables (*P*=0.012), red meat (*P*=0.017), butter (*P*=0.039), sweetened carbonated beverages (*P*=0.036) and nuts (*P*=0.038)). In conclusion, we found that rs1260326 (*GCKR*) is significantly associated with higher TAG concentrations, but is modulated by adherence to the MD.

Key words: Glucokinase regulatory protein: TAG concentration: Nutrigenetics: Mediterranean diet

The glucokinase (*GCK*) gene is a regulator of glucose storage and disposal in the liver, and its activity is modulated by binding to the glucokinase regulatory protein (GCKR). Mice deficient in GCKR show reduced expression of GCK and defects in the clearance of glucose, but maintain a normal activity of GCK⁽¹⁾. Overexpression by the adenoviral-mediated hepatic overproduction of GCKR results in increased GCK activity but lower levels of fasting glucose, while overexpression of GCK in the liver leads to a decrease in glucose concentrations and an increase in TAG concentrations^(2,3). Taken together, these data strongly support a role for these genes in determining plasma TAG and glucose concentrations in human subjects.

Initial studies^(4,5) have reported an intronic polymorphism of *GCKR* (rs780094) to be associated with TAG concentrations in populations of European origin. Subsequent research has found similar associations for TAG and opposing effects on glucose concentrations^(6–8). Moreover, Orho-Melander *et al.*⁽⁵⁾ suggested rs1260326 (P446L) as the functional SNP for the observed effects of the intronic rs780094.

The common variant –30G > A (rs1799884) in the *GCK* gene promoter region has been associated with an increased risk of diabetes⁽⁹⁾, hyperglycaemia^(10–12), impaired β -cell function^(10–13) and lower TAG concentrations. An initial study on the combined effects of these polymorphisms has shown the additive effect of *GCKR* (446L) and *GCK* (–30G)

Abbreviations: GCK, glucokinase; GCKR, glucokinase regulatory protein; MD, Mediterranean diet.

* **Corresponding author:** D. Corella, fax +34 963864166, email dolores.corella@uv.es

alleles in conferring lower fasting glucose and insulin levels, but higher levels of TAG⁽⁹⁾. However, the effects of genetic variants in both genes in determining fasting TAG and glucose concentrations in Mediterranean populations, as well as the possible gene–diet interactions regarding adherence to the Mediterranean diet (MD) and its specific food items, remain to be investigated. We have shown that the MD has a beneficial impact on TAG concentrations and on other biochemical parameters related to a higher risk of CVD⁽¹⁴⁾. Since diet is one of the most important environmental factors in determining plasma TAG and glucose concentrations, we hypothesise that adherence to the MD may be directly involved in the genetic modulation of different intermediate phenotypes of the *GCK* and *GCKR* genes. Although two recent studies have analysed gene–diet interactions for these genes^(15,16), neither of them has focused on examining the effects of the MD in such interactions.

The aims of the present study are as follows: first, to analyse the effects of *GCKR* P446L (rs1260326) and *GCK* –30G > A (rs1799884) on fasting TAG and glucose concentrations in a high-cardiovascular risk Mediterranean population; second, to study the potential modulation of the effect of *GCKR* SNP on TAG concentrations by adherence to the MD; third, to analyse the joint association and the potential interaction of each food item of the MD in relation to TAG concentrations.

Subjects and methods

We studied 945 unrelated consecutive subjects (340 men and 605 women), aged 55–80 years, who participated in the PRE-venición con DIeta MEDiterránea Study. These subjects had been recruited in the Valencia Region (Spain) from October 2003 to September 2008, and details of this study have been reported elsewhere⁽¹⁴⁾. Briefly, high-risk participants were selected by physicians in primary care centres. Eligible subjects were elderly community-dwelling persons who fulfilled at least one of two criteria: type 2 diabetes or three or more CVD risk factors (current smoking, hypertension, dyslipidaemia, overweight or a family history of premature CVD). Exclusion criteria were based on a personal history of CVD, any severe chronic illness, and drug or alcohol addiction. Here, we included data from the baseline cross-sectional examination. The present study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects were approved by the Institutional Review Board of the Valencia University. Written informed consent was obtained from all subjects.

Anthropometric, biochemical, clinical and lifestyle determinations

Anthropometric variables, such as height, weight and waist circumference, were measured by standard techniques⁽¹⁴⁾. BMI was calculated as weight (kg) divided by height (m²). Fasting blood samples were obtained for each participant, and stored at –80°C until biochemical analyses. Fasting glucose, total cholesterol, TAG, HDL-cholesterol and LDL-cholesterol were determined as reported previously⁽¹⁴⁾.

Baseline examination included assessment of standard cardiovascular risk factors, medication use, sociodemographic factors and lifestyle variables, as detailed previously⁽¹⁴⁾. Physical activity was estimated by the Minnesota Leisure-Time Physical Activity Questionnaire⁽¹⁷⁾.

Dietary intake

We determined food consumption by a previously validated semi-quantitative FFQ⁽¹⁸⁾. Energy and nutrient intake was calculated from Spanish food composition tables⁽¹⁹⁾. The baseline examination also included the administration of a validated fourteen-item questionnaire that indicated the degree of adherence to the traditional MD⁽²⁰⁾. Values of 0 or 1 were assigned to each of the fourteen dietary components. For each of the following categories, one point was given: (1) the use of olive oil as the principal source of fat for cooking; (2) ≥ 4 tablespoons (one tablespoon = 13.5 g) of olive oil/d (including that used in frying, salads, meals eaten away from home, etc.); (3) ≥ 2 servings of vegetables/d; (4) ≥ 3 pieces of fruit/d; (5) < 1 serving of red meat or sausages/d; (6) < 1 serving of butter, margarine or cream/d; (7) < 1 cup (one cup = 100 ml) of sugar-sweetened beverages/d; (8) ≥ 7 servings of red wine/week; (9) ≥ 3 servings of pulses/week; (10) ≥ 3 servings of fish/week; (11) < 2 commercial pastries/week; (12) ≥ 3 servings of nuts/week; (13) preferring white meat over red meat; or for consuming (14) ≥ 2 servings/week of a dish with a traditional sauce of tomatoes, garlic, onion or leeks sautéed in olive oil. If the condition was not met, 0 points were recorded for that category. The greater the score obtained from the questionnaire, the greater the adherence to the MD. Further, we created four groups of adherence to the MD according to the points obtained in this questionnaire: group 1 (3–6 points, 15.6%), group 2 (7–8 points, 35.2%), group 3 (9–10 points, 32.6%) and group 4 (11–14 points, 16.6%), such that group 1 had the lowest adherence to the MD and group 4 had the highest adherence.

SNP analyses

DNA was extracted from the buffy coat fraction of centrifuged blood using the MagNa Pure LC DNA Isolation Kit (Roche Diagnostics). *GCKR* P446L (rs1260326) and *GCK* –30A > G (rs1799884) were genotyped using Taqman SNP allelic discrimination by means of an ABI 7900 HT (Applied Biosystems). Both genes were in Hardy–Weinberg equilibrium in the study samples ($P=0.38$ and $P=0.144$, respectively). A concordance rate of 100% was seen in 200 blinded quality-control samples for all SNP assayed.

Statistical analysis

Categorical variables were compared using χ^2 tests. Student's *t* test and ANOVA were applied to compare crude means. Dominant, co-dominant and recessive models of inheritance were first tested to determine the effects of the variant allele. TAG concentrations were log-transformed for statistical testing.

A dominant model for the P446L SNP and a recessive model for the $-30G > A$ SNP were the most significant, and these categories (PP *v.* PL + LL and GG + GA *v.* AA, respectively) were considered for further analyses. We further adjusted for potential confounders using ANCOVA. Joint analysis was performed considering the genotype (PP or PL + LL) and each category of the food item using ANCOVA.

In multivariate analysis, the models were adjusted for covariates including age (continuous), BMI (kg/m^2 ; continuous), sex, diabetes (yes or no), smoking (current smokers or no smokers) and lipid medication (yes or no). Further adjustment of the models for physical activity and total energy intake did not change the statistical significance of the results. Logistic regression methods were also used to estimate the OR of hypertriacylglycerolaemia and diabetes associated with the polymorphisms. Univariate and multivariate logistic regression models including the potential confounders were fitted. Statistical analyses were performed using the SPSS package, version 15.0 (SPSS). Statistical significance was set at the 0.05 level, and all tests were two-tailed.

Results

Characteristics of the population

Clinical, biochemical, dietetic and genetic characteristics are shown in Table 1. Participants were high-cardiovascular risk subjects and then, prevalence of hypertriacylglycerolaemia and diabetes was high. The mean of the adherence to the MD was 8.5 points. The percentages of positive points for the most representative food items of the MD are listed in Table 1.

Association between the glucokinase regulatory protein gene (rs1260326) (P446L) and glucokinase gene (rs1799884) ($-30G > A$) polymorphisms with fasting TAG and glucose concentrations

The *GCKR* (P446L) SNP showed a significant association with TAG concentrations (Table 2). Subjects who were heterozygous and homozygous for the minor allele (L) showed higher TAG concentrations (*P* crude model=0.006). The best model identified for this genetic variant was dominant (*P*=0.002). Multivariate adjustment for age, sex, BMI, smoking, diabetes and lipid medication did not significantly change the results (*P*=0.014). This association was homogeneous in males and females. No associations were found for other lipids parameters and the *GCKR* polymorphism (data not shown).

No significant association was found between the $-30G > A$ SNP in the *GCK* gene and TAG concentrations in the crude analysis. However, after multivariate adjustment, homozygous subjects for the minor allele (AA) presented statistically lower TAG concentrations than the GG + GA subjects (*P*=0.033). No statistical association was found between either of the SNP analysed and glucose concentrations.

We also examined the combined effects for each genotype and found an additive effect on TAG concentrations. After full multivariate adjustment, we found that subjects with the

highest genetic risk for TAG, namely PL/LL (*GCKR*) + GG/GA (*GCK*), had a 37% higher TAG concentration compared with those with a genetic-free risk (i.e. PP (*GCKR*) + AA (*GCK*)) (*P*=0.033).

No combined effects were found for glucose concentrations. We also studied the gene \times gene interaction, but no statistically significant interactive effects were found.

We also estimated the association with the risk of hypertriacylglycerolaemia and diabetes. We found a significantly higher risk for hypertriacylglycerolaemia in the PL + LL subjects (OR 1.86, 95% CI 1.33, 2.61) than in the PP subjects.

Table 1. Anthropometric, clinical, biochemical, lifestyle and genetic characteristics of the high-cardiovascular risk Mediterranean population (Mean values, standard deviations, number of subjects and percentages)

Variables	Mean	SD
Continuous variables		
Sex (n)		
Male		340
Female		605
Age (years)	67.3	6.2
Weight (kg)	77.4	12.4
BMI (kg/m^2)	30.9	5.1
Waist (cm)	104.3	11.8
TC (mmol/l)	5.38	1.03
HDL-C (mmol/l)	1.37	0.36
LDL-C (mmol/l)	3.34	0.93
TAG (mmol/l)	1.48	0.91
Fasting glucose (mmol/l)	6.83	2.22
SBP (mmHg)	147	22
DBP (mmHg)	82	11
Score for MD adherence*	8.5	2.0
Energy intake		
kJ	9228	2454
kcal	2204	586
	<i>n</i>	%
Categorical variables		
Use of olive oil as principal source of fat	754	86.3
Olive oil ≥ 4 tbs/d	708	81.2
Vegetables ≥ 2 servings/d	527	46.7
Fruit consumption ≥ 3 servings/d	400	45.9
Red meat < 1 serving/d	418	47.9
Butter < 1 serving/d	793	90.7
Sugar-sweetened beverages < 1 serving/d	758	86.8
Wine ≥ 7 cups/week	179	20.6
Legumes ≥ 3 servings/week	220	25.2
Fish ≥ 3 servings/week	703	80.4
Commercial pastries < 2 servings/week	602	68.9
Nuts ≥ 3 servings/week	229	26.3
Preferring white meat over red meat	642	73.5
Traditional sauce 'sofrito' ≥ 2 servings/week	482	55.4
Current smokers	109	11.9
Diabetes	448	48.5
Hypertriacylglycerolaemia	266	28.6
GCKR (rs1260326) (P446L) polymorphism		
PP	284	29.6
PL	488	50.9
LL	187	19.5
GCK (rs1799884) ($-30G > A$) polymorphism		
GG	598	64.6
GA	300	32.4
AA	27	2.9

TC, total cholesterol; HDL-C, HDL-cholesterol; LDL-C, LDL-cholesterol; SBP, systolic blood pressure; DBP, diastolic blood pressure; MD, Mediterranean diet; tbs, tablespoon; GCKR, glucokinase regulatory protein; GCK, glucokinase.

* Score for MD adherence: 14-unit MD score (ranging from 0 (minimum) to 14 points (maximum)).

Table 2. Association between fasting TAG and glucose concentrations (mmol/l) and P446L (rs1260326) glucokinase regulatory protein (*GCKR*) gene, rs1799884 (-30G > A) glucokinase (*GCK*) gene and combined effect (*GCKR* + *GCK*) (Mean values and standard deviations or standard errors)

		P446L (rs1260326) <i>GCKR</i> gene								
		PP (n 288)		PL (n 488)		LL (n 187)		<i>P</i> *	<i>P</i> _{dominant}	
		Mean	SE	Mean	SE	Mean	SE			
TAG (mmol/l)										
Crude										
Mean		1.33		1.51		1.57		0.006	0.002	
SD		0.58		0.88		0.90				
Model 1†		1.33	0.05	1.50	0.04	1.57	0.06	0.012	0.004	
Model 2‡		1.34	0.05	1.50	0.04	1.58	0.07	0.035	0.014	
Glucose (mmol/l)										
Crude										
Mean		6.78		6.89		6.53		0.151	0.068	
SD		2.08		2.25		1.90				
Model 1†		6.77	0.13	6.91	0.10	6.46	0.17	0.071	0.866	
Model 2‡		6.71	0.12	6.84	0.09	6.55	0.14	0.224	0.141	
rs1799884 (-30G > A) <i>GCK</i> gene										
		GG (n 598)		GA (n 300)		AA (n 27)		<i>P</i> *	<i>P</i> _{recessive}	
		Mean	SE	Mean	SE	Mean	SE			
TAG (mmol/l)										
Crude										
Mean		1.47		1.48		1.24		0.258	0.1	
SD		0.82		0.83		0.62				
Model 1†		1.47	0.03	1.48	0.05	1.18	0.16	0.199	0.027	
Model 2‡		1.48	0.04	1.46	0.05	1.17	0.18	0.095	0.033	
Glucose (mmol/l)										
Crude										
Mean		6.74		6.83		7.19		0.517	0.323	
SD		2.16		2.08		1.89				
Model 1†		6.74	0.09	6.80	0.13	7.18	0.43	0.608	0.347	
Model 2‡		6.75	0.08	6.78	0.12	6.09	0.40	0.262	0.105	
Combined effect (<i>GCKR</i> dominant + <i>GCK</i> recessive)										
		PP + AA (n 12)		PP + (GA + AA) (n 253)		(PL + LL) + AA (n 608)		(PP + LL) + (GG + GA) (n 14)		<i>P</i> for joint effect
		Mean	SE	Mean	SE	Mean	SE	Mean	SE	
TAG (mmol/l)										
Crude										
Mean		1.12		1.33		1.35		1.53		0.012
SD		0.49		0.58		0.72		0.90		
Model 1†		1.06	0.24	1.35	0.05	1.27	0.22	1.53	0.03	0.010
Model 2‡		1.11	0.27	1.36	0.06	1.21	0.24	1.52	0.04	0.033
Glucose (mmol/l)										
Crude										
Mean		7.19		6.75		7.20		6.78		0.799
SD		1.69		2.05		2.13		2.16		
Model 1†		7.32	0.64	6.73	0.14	7.05	0.59	6.78	0.09	0.784
Model 2‡		6.49	0.58	6.73	0.12	5.74	0.56	6.77	0.08	0.311

* *P* value for comparisons between genotypes and TAG concentrations. *P* value for log TAG.

† Adjusted for age, sex and BMI.

‡ Model 1+smoking + diabetes + lipid medication.

Further multivariate adjustment did not change the statistical significance ($P=0.001$). The *GCK* SNP was not associated with hypertriglycerolaemia risk. The regression logistic model for combined effects had a low number of participants in both categories of reference for comparisons; therefore, we could not find statistically significant results (see Table S1 of the supplementary material, available online at <http://www>.

[journals.cambridge.org/bjn](http://www.journals.cambridge.org/bjn)). Opposing effects were observed for type 2 diabetes mellitus risk in relation to the L446P and -30G > A polymorphisms. The LL subjects (*GCKR*) had a lower estimated risk for type 2 diabetes mellitus, while the AA subjects had a statistically higher risk (LL (*GCKR*): OR_{adjusted} 0.80, 95% CI 0.66, 0.96; $P=0.017$ and AA (*GCK*): OR_{adjusted} 3.3, 95% CI 1.2, 9.2; $P=0.023$) (see Table S2 of

the supplementary material, available online at <http://www.journals.cambridge.org/bjn>.

Interaction between glucokinase regulatory protein gene (rs1260326) (P446L) polymorphism and adherence to the Mediterranean diet in determining TAG concentrations

We tested whether adherence to the MD modified the associations between the P446L SNP and TAG concentrations, according to groups of adherence. Based on the score obtained in the questionnaire of adherence to the MD, four categories of adherence to the MD were considered. We found statistically significant differences in TAG concentrations by genotype among those with a lower adherence to the MD (group 1–group 2–group 3; $P=0.011$). In contrast, no differences were found between genotypes and TAG concentrations in the group of higher adherence to the MD ($P=0.917$; Fig. 1). However, the test for interaction was not significant.

Joint associations and interactions between glucokinase regulatory protein gene (rs1260326) (P446L) polymorphism and food items of the Mediterranean diet in determining TAG concentrations

We examined the joint associations between the P446L SNP and the major foods that characterise the MD in relation to TAG concentrations (Table 3). Statistically significant joint associations were found for olive oil, vegetables, meat, butter, sweetened carbonated beverages and nuts ($P=0.035$, $P=0.012$, $P=0.017$, $P=0.039$, $P=0.036$ and $P=0.038$, respectively). Subjects with the highest risk (genetic risk (PL + LL)) and who had a high consumption of sweetened carbonated beverages (>1 per d) had 0.19 mmol/l higher TAG




Fig. 1. Gene \times diet interaction for TAG concentrations according to adherence to the Mediterranean diet (AMD). Adjusted means of TAG are shown by the *GCKR* polymorphism according to the strata of AMD. Means were adjusted by sex, age, smoking, BMI, diabetes and lipid medication. P value for interaction terms between TAG concentrations and *GCKR* polymorphism (as a dominant pattern) was obtained in the hierarchical multivariate interaction model (P for *GCKR* (P446L) \times AMD interaction = 0.346). P value for log TAG. G1+G2 + G3, groups 1, 2 and 3 of AMD; G4, group 4 of AMD.




Fig. 2. TAG concentrations according to the joint effects of adherence to the six-point score and the genetic dominant pattern (PP, PL + LL). The analyses were adjusted for sex, age, BMI, smoking, diabetes and lipid medication (P for joint effects = 0.027).

concentrations than those with the lowest allele of risk (PP) and lowest consumption of sweetened carbonated beverages (≤ 1 per d). Similar results were observed for butter, margarine or cream consumption per d, although with lower effects on the reduction of TAG concentrations (2.5%).

For the other main items that characterise the MD, subjects with the highest genetic risk and lower consumption of olive oil had 8.2% (0.25 mmol/l) higher TAG concentrations than did those with no allele risk and higher consumption of olive oil (≥ 4 tablespoons) ($P=0.033$). Similarly, those who consumed less than two servings per d of vegetables, or more than one serving per d of red meat, or less than three servings of nuts per week had 0.30 (23%), 0.06 (4.2%) or 0.11 (8.2%) mmol/l higher TAG concentrations, respectively. All of the means were multivariate adjusted. Further adjustment for energy intake did not change the statistical significance of the results.

We further assessed the potential interactions between the P446L SNP and the major foods characterising the MD in relation to TAG concentrations. No significant interactions were found. We also analysed the specific effect of each food item on TAG concentrations, independently of the genotype. Significant associations were found (inverse) for vegetables and meat consumption ($P=0.022$ and $P=0.035$, respectively), and these effects were enhanced when genotype was considered ($P=0.012$ and $P=0.017$, respectively).

We tried to identify the main items of the MD that specifically could be associated with the genetic risk and TAG concentrations. We created a mini-score with the six items that were significant in the first analysis (olive oil, vegetables, red meat, butter, sweetened carbonated beverages and nuts). In a further analysis, we found a significant joint association. The subjects with the highest genetic risk (PL + LL) and lower food item consumption (1–4 points) had 22.2% higher TAG concentrations. In a stratified analysis, we did not find any differences in TAG concentrations between genotypes in those subjects who consumed five or six items of this mini-score, while significant differences were observed when the consumption was one to four items (Fig. 2).

Table 3. Interaction and joint effects between components of the score of adherence to the Mediterranean diet and the glucokinase regulatory protein gene polymorphism on TAG concentrations (mmol/l)

(Mean values with their standard errors)

	PP		PL + LL		<i>P</i> for joint effects*	<i>P</i> food item*
	Mean	SE	Mean	SE		
Olive oil (tbs/d)						
< 4	1.44	0.08	1.63	0.07	0.035	0.116
≥ 4	1.30	0.06	1.49	0.04		
Vegetables (servings/d)†						
< 2	1.40	0.07	1.59	0.05	0.012	0.022
≥ 2	1.29	0.06	1.47	0.04		
Fruit (including natural fruit juice, servings/d)‡						
< 3	1.33	0.06	1.51	0.05	0.084	0.931
≥ 3	1.34	0.06	1.52	0.05		
Red meat (hamburgers or meat products, servings/d)§						
≥ 1	1.24	0.09	1.41	0.08	0.017	0.035
< 1	1.36	0.06	1.54	0.04		
Butter, margarine or cream (servings/d)						
≥ 1	1.20	0.11	1.38	0.10	0.039	0.145
< 1	1.35	0.06	1.53	0.04		
Sweetened carbonated beverages (servings/d)¶						
≥ 1	1.28	0.10	1.47	0.09	0.036	0.896
< 1	1.34	0.06	1.53	0.04		
Wine (servings/week)**						
< 7	1.32	0.06	1.42	0.01	0.051	0.230
≥ 7	1.42	0.09	1.60	0.08		
Legumes (servings/week)††						
< 3	1.33	0.06	1.52	0.04	0.116	0.703
≥ 3	1.34	0.06	1.53	0.06		
Fish/shellfish (servings/week)‡‡						
< 3	1.32	0.06	1.50	0.05	0.104	0.541
≥ 3	1.35	0.06	1.53	0.05		
Sweets or pastries (servings/week)§§						
< 2	1.34	0.07	1.52	0.06	0.101	0.476
≥ 2	1.33	0.06	1.52	0.04		
Nuts (servings/week)¶¶						
< 3	1.31	0.06	1.50	0.04	0.038	0.314
≥ 3	1.38	0.07	1.57	0.06		
Consumption preferentially of chicken, rabbit or turkey instead of veal, pork, hamburger, sausage¶¶¶						
No	1.27	0.08	1.45	0.06	0.099	0.480
Yes	1.35	0.06	1.54	0.04		
Vegetables, pasta, rice with sofritos (servings/week)***						
< 2	1.32	0.06	1.51	0.05	0.088	0.693
≥ 2	1.34	0.06	1.52	0.04		

tbs, Tablespoon.

* *P* value for log TAG. The analyses were adjusted for sex, age, BMI, smoking, lipid medication and diabetes.

† Serving of vegetables is 200 g.

‡ Serving of fruit is 100–250 g.

§ Serving of red meat is 100–150 g.

|| Serving of butter, margarine or cream is 12 g.

¶ Serving of sweetened carbonated beverages is 200 ml.

** Serving of wine is one cup, 100 ml.

†† Serving of legumes is 150 g.

‡‡ Serving of fish is 100–150 g and 200 g (4–5 units) for shellfish.

§§ Serving of sweets or pastries is 50 g unless other thing is specified in the questionnaire⁽¹⁸⁾.

¶¶ Serving of nuts is 30 g.

¶¶¶ Serving of poultry is 100–150 g.

*** Serving of sofritos is 200 g.

Discussion

The present study was carried out in a high-cardiovascular risk Mediterranean population. We replicated previously reported associations between the minor allele (P variant) of rs1260326 in the *GCKR* gene and higher TAG concentrations^(4–7,9,21–23), and reported for the first time a modulation of this association by adherence to the MD.

The mechanism that may explain the observed results is not yet understood. A recent molecular study⁽²⁴⁾ proposed a mutational mechanism that could explain the reported association of this variant with raised TAG and lower glucose concentrations. This group found that regulation by physiological concentrations of the phosphate ester fructose-6-phosphate is reduced in the *GCKR* T-allele Pro446Leu, which results

indirectly in increased GCK activity⁽²⁴⁾. Altered GCK regulation in the liver is predicted to enhance glycolytic flux, promoting hepatic glucose metabolism and elevating concentrations of malonyl-CoA, a substrate for *de novo* lipogenesis. This provides a mutational mechanism for the reported association of this variant with raised TAG and lower glucose levels⁽²⁴⁾. Taken together, these data support the central position of GCKR in pathways that regulate hepatic TAG as well as glucose metabolism in humans.

With respect to the polymorphism in the *GCK* gene, most previous studies have analysed its effect on glucose concentrations and have reported significant associations^(10,21,25,26). In our population, we found no association with fasting glucose, although the estimated risk of diabetes was lower in the AA subjects, as reported in other studies^(10,23).

Opposing effects of *GCK* and *GCKR* polymorphisms on TAG concentrations have been reported previously. The minor allele of the $-30G > A$ SNP has been associated with lower TAG concentrations. In the present study, we also found a significant association between the A allele and lower TAG concentrations. Furthermore, when we analysed the combined effect of the *GCK* gene (recessive model) and *GCKR* gene (dominant model), we found a potentially increased effect on TAG concentrations in subjects with the AA (*GCK*) and PP (*GCKR*) alleles having lower TAG concentrations when compared with those having the PL + LL and GG + GA alleles.

Several studies have reported additive effects of rs7800094 and rs1799884 on fasting glucose concentrations^(5,7,9), but this combined effect on TAG concentrations has not been widely studied. Tam *et al.*⁽²¹⁾ used additive genetic models in two cohorts of Chinese populations but did not observe an interaction with fasting TAG. Differences between genetic effect size, risk allele frequency, population sample and environmental exposure between European and Asian populations may explain this discrepancy. Additional interaction analyses in further studies are warranted to validate the present findings.

Regarding the gene–diet interaction, we investigated the potential joint effects of genetic variants and the MD on TAG concentrations. We reported a significant difference in TAG between PP *v.* PL + LL in subjects with a lower adherence to the MD, while this effect was not found in the group with a high adherence to the MD. Supporting the presence of gene–diet interactions involving the *GCKR* gene, two recent studies^(15,16) have reported statistically significant interactions between some SNP in the *GCKR* gene and dietary intake in determining glucose-related traits. In one of them⁽¹⁵⁾, carried out in fourteen cohorts comprising about 48 000 participants of European descent aimed at studying interactions of whole grain intake with loci previously associated in genome-wide association studies with fasting glucose (sixteen loci) and/or insulin (two loci), the strongest interaction with whole grain intake was between rs780094 in the *GCKR* gene in determining fasting insulin. In the other work⁽¹⁶⁾ carried out in subjects with the metabolic syndrome participating in the LIPGENE study, the authors found significant interactions between the *GCKR* rs1260326 polymorphism and plasma *n*-3 PUFA levels

modulating insulin resistance and inflammatory markers. These studies^(15,16) did not investigate the potential mechanisms behind the reported gene–diet interactions. In the present study, the first to investigate the combined effects of the MD and the genetic risk of P446L on TAG concentrations, we did not study the biological mechanism to explain the observed results. However, taking into account some previous reports^(27,28) regarding the effect of dietary fat on the postprandial lipid response, we may suggest some hypotheses. Perez-Martinez *et al.*⁽²⁷⁾ studied the postprandial TAG response to a high-fat diet and reported that subjects with the risk allele had higher fasting TAG, hypertriglycerolaemia and a greater postprandial TAG response. Another dietary intervention study (postprandial lipid response after a high-fat challenge)⁽²⁸⁾ also reported results consistent with those published by Perez-Martinez *et al.*⁽²⁷⁾. These findings could support the present results with respect to modulation of the genetic susceptibility of TAG concentrations by the profile of the high fat content of the traditional MD (a high percentage of fat intake (30–35%) is mostly from MUFA, given the consumption of olive oil)^(14,20). A high fat content from SFA could have a negative effect on L carriers with respect to TAG concentrations, while fats from olive oil and nuts (characteristic foods of the MD) could modulate this risk (as shown in our sample for these two individual items). In the present study, we showed a significant joint association (gene–food items of the MD) between TAG concentration and the consumption of olive oil, vegetables and nuts, as well as the low consumption of meat products, butter and sweetened beverages. The MD is well known to have beneficial effects on lipid metabolism as well as on other parameters^(29,30). Currently, the importance of nutrigenomics and personalised diets based on genetics is gaining attention⁽³¹⁾. Therefore, a number of studies have analysed the potential modulation of different phenotypes by nutrient compounds^(32,33). However, the current paradigm is to study the overall dietary pattern, instead of providing a simple assessment of isolated nutrients, because of the synergistic or antagonistic effects of food items and nutrients⁽³⁴⁾. After analysing the overall Mediterranean dietary pattern, we wanted to study the individual items that make up the MD and contribute to the modulation of the genetic effect. As we reported above, six items of the MD were significant, which agrees with the synergistic or antagonistic effects of food items in relation to metabolic diseases.

The benefits of the high consumption of vegetables in cardiovascular intermediate phenotypes have been reported in many studies⁽³⁵⁾. In Mediterranean countries, vegetable consumption is linked to a high consumption of olive oil because it is used to dress salads. Therefore, its role in the prevention of the genetic risk for TAG concentrations could actually be associated with olive oil consumption.

Meat products and butter have a high content of SFA, and SFA have also been associated with cardiovascular risk⁽³⁶⁾. Therefore, LL carriers could find more benefits from a low consumption of these products to decrease their TAG concentrations.

There are some limitations in the present study, mainly derived from the fact that cross-sectional data have been analysed; therefore, causal inference cannot be made.

In conclusion, we found that the P446L SNP in the *GCKR* gene is significantly associated with TAG concentrations in this Mediterranean population, in agreement with previous findings. We also showed combined gene–gene associations with TAG concentrations. We found, for the first time, that the Mediterranean dietary pattern could modulate the effects of the *GCKR* gene variation on TAG concentrations. We also identified the main food items of this pattern that exert their beneficial effects on TAG concentrations (high consumption of olive oil, nuts and vegetables and lower consumption of red meat, butter and sweetened beverages). The present findings suggest that the adoption of the MD may decrease TAG concentrations, especially among the genetically high-risk population. However, more studies analysing dietary patterns are recommended to better establish personalised recommendations.

Acknowledgements

This study was supported by grants from the Ministerio de Ciencia e Innovación and Fondo Europeo de Desarrollo Regional: CIBER CB06/03/0035, PI07-0954, CNIC-06 and AGL2010-22319-C03-03, PFI07-0954, PFI07-0954 and the Generalitat Valenciana (GVACOMP2011-151 and AP111/10). The authors' contributions are as follows: D. C. conceptualised and designed the study; M. S.-P., M. G., J. V. S., O. P., P. G.-S., J. I. G. and D. C. were involved in the acquisition of the data; M. S.-P., L. Q. and D. C. analysed and interpreted the data; M. S.-P. and D. C. drafted the manuscript; M. S.-P., M. G., J. V. S., O. P., P. G.-S., J. I. G., L. Q. and D. C. critically revised the manuscript for important intellectual content. The authors declare that there are no conflicts of interest.

References

1. Grimsby J, Coffey JW, Dvornozniak MT, *et al.* (2000) Characterization of glucokinase regulatory protein-deficient mice. *J Biol Chem* **17**, 7826–7831.
2. Slosberg ED, Desai UJ, Fanelli B, *et al.* (2001) Treatment of type 2 diabetes by adenoviral-mediated overexpression of the glucokinase regulatory protein. *Diabetes* **50**, 1813–1820.
3. Farrelly D, Brown KS, Tieman A, *et al.* (1999) Mice mutant for glucokinase regulatory protein exhibit decreased liver glucokinase: a sequestration mechanism in metabolic regulation. *Proc Natl Acad Sci U S A* **96**, 14511–14516.
4. Saxena R, Voight BF, Lyssenko V, *et al.* (2007) Genome-wide association analysis identifies loci for type 2 diabetes and triglyceride levels. *Science* **316**, 1331–1336.
5. Orho-Melander M, Melander O, Guiducci C, *et al.* (2008) Common missense variant in the glucokinase regulatory protein gene is associated with increased plasma triglyceride and C-reactive protein but lower fasting glucose concentrations. *Diabetes* **57**, 3112–3121.
6. Qi Q, Wu Y, Li H, *et al.* (2009) Association of GCKR rs780094, alone or in combination with GCK rs1799884, with type 2 diabetes and related traits in a Han Chinese population. *Diabetologia* **52**, 834–843.
7. Sparsø T, Andersen G, Nielsen T, *et al.* (2008) The GCKR rs780094 polymorphism is associated with elevated fasting serum triacylglycerol, reduced fasting and OGTT-related insulinaemia, and reduced risk of type 2 diabetes. *Diabetologia* **51**, 70–75.
8. Dupuis J, Langenberg C, Prokopenko I, *et al.* (2010) New genetic loci implicated in fasting glucose homeostasis and their impact on type 2 diabetes risk. *Nat Genet* **42**, 105–116.
9. Vaxillaire M, Cavalcanti-Proença C, Dechaume A, *et al.* (2008) The common P446L polymorphism in GCKR inversely modulates fasting glucose and triglyceride levels and reduces type 2 diabetes risk in the DESIR prospective general French population. *Diabetes* **57**, 2253–2257.
10. Vaxillaire M, Veslot J, Dina C, *et al.* (2008) Impact of common type 2 diabetes risk polymorphisms in the DESIR prospective study. *Diabetes* **57**, 244–254.
11. März W, Nauck M, Hoffmann MM, *et al.* (2004) G(–30)A polymorphism in the pancreatic promoter of the glucokinase gene associated with angiographic coronary artery disease and type 2 diabetes mellitus. *Circulation* **15**, 2844–2849.
12. Rose CS, Ek J, Urhammer SA, *et al.* (2005) –30G > A polymorphism of the beta-cell-specific glucokinase promoter associates with hyperglycemia in the general population of whites. *Diabetes* **54**, 3026–3031.
13. Yamada K, Yuan X, Ishiyama S, *et al.* (1997) Clinical characteristics of Japanese men with glucokinase gene beta-cell promoter variant. *Diabetes Care* **20**, 1159–1161.
14. Estruch R, Martínez-González MA, Corella D, *et al.* (2006) Effects of a Mediterranean-style diet on cardiovascular risk factors: a randomized trial. *Ann Intern Med* **145**, 1–11.
15. Nettleton JA, McKeown NM, Kanoni S, *et al.* (2010) Interactions of dietary whole-grain intake with fasting glucose- and insulin-related genetic loci in individuals of European descent: a meta-analysis of 14 cohort studies. *Diabetes Care* **33**, 2684–2691.
16. Perez-Martinez P, Delgado-Lista J, Garcia-Rios A, *et al.* (2011) Glucokinase regulatory protein genetic variant interacts with omega-3 PUFA to influence insulin resistance and inflammation in metabolic syndrome. *PLoS One*; **6**, e20555.
17. Elosua R, Garcia M, Aguilar A, *et al.* (2000) Validation of the Minnesota Leisure Time Physical Activity Questionnaire in Spanish women. Investigators of the MARATHON Group. *Med Sci Sports Exerc* **32**, 1431–1437.
18. Fernández-Ballart JD, Piñol JL, Zazpe I, *et al.* (2010) Relative validity of a semi-quantitative food-frequency questionnaire in an elderly Mediterranean population of Spain. *Br J Nutr* **103**, 1808–1816.
19. Mataix J (2003) *Tabla de composición de alimentos (Food Composition Tables)*. Granada: University of Granada.
20. Schröder H, Fitó M, Estruch R, *et al.* (2011) A short screener is valid for assessing Mediterranean diet adherence among older Spanish men and women. *J Nutr* **141**, 1140–1145.
21. Tam CH, Ma RC, So WY, *et al.* (2009) Interaction effect of genetic polymorphisms in glucokinase (GCK) and glucokinase regulatory protein (GCKR) on metabolic traits in healthy Chinese adults and adolescents. *Diabetes* **58**, 765–769.
22. Tam CH, Ho JS, Wang Y, *et al.* (2010) Common polymorphisms in MTNR1B, G6PC2 and GCK are associated with increased fasting plasma glucose and impaired beta-cell function in Chinese subjects. *PLoS One* **8**, e11428.
23. Takeuchi F, Katsuya T, Chakrewarthy S, *et al.* (2010) Common variants at the GCK, GCKR, G6PC2-ABCB11 and MTNR1B loci are associated with fasting glucose in two Asian populations. *Diabetologia* **53**, 299–308.

24. Beer NL, Tribble ND, McCulloch LJ, *et al.* (2009) The P446L variant in GCKR associated with fasting plasma glucose and triglyceride levels exerts its effect through increased glucokinase activity in liver. *Hum Mol Genet* **18**, 4081–4088.
25. Webster RJ, Warrington NM, Weedon MN, *et al.* (2009) The association of common genetic variants in the APOA5, LPL and GCK genes with longitudinal changes in metabolic and cardiovascular traits. *Diabetologia* **52**, 106–114.
26. Hu C, Zhang R, Wang C, *et al.* (2010) Effects of GCK, GCKR, G6PC2 and MTNR1B variants on glucose metabolism and insulin secretion. *PLoS One* **5**, e11761.
27. Perez-Martinez P, Corella D, Shen J, *et al.* (2009) Association between glucokinase regulatory protein (GCKR) and apolipoprotein A5 (APOA5) gene polymorphisms and triacylglycerol concentrations in fasting, postprandial, and fenofibrate-treated states. *Am J Clin Nutr* **89**, 391–399.
28. Shen H, Pollin TI, Damcott CM, *et al.* (2009) Glucokinase regulatory protein gene polymorphism affects postprandial lipemic response in a dietary intervention study. *Hum Genet* **126**, 567–574.
29. Estruch R (2010) Anti-inflammatory effects of the Mediterranean diet: the experience of the PREDIMED study. *Proc Nutr Soc* **69**, 333–340.
30. Sotos-Prieto M, Zulet MA & Corella D (2010) Scientific evidence of the Mediterranean diet effects in determining intermediate and final cardiovascular disease phenotypes. *Med Clin (Barc)* **134**, 22–29.
31. Fenech M, El-Sohemy A, Cahill L, *et al.* (2011) Nutrigenetics and nutrigenomics: viewpoints on the current status and applications in nutrition research and practice. *J Nutrigenet Nutrigenomics* **4**, 69–89.
32. Corella D, Tai ES, Sorlí JV, *et al.* (2010) Association between the APOA2 promoter polymorphism and body weight in Mediterranean and Asian populations: replication of a gene-saturated fat interaction. *Int J Obes (Lond)* **35**, 666–675.
33. Sotos-Prieto M, Guillén M, Guillem-Sáiz P, *et al.* (2010) The rs1466113 polymorphism in the somatostatin receptor 2 gene is associated with obesity and food intake in a Mediterranean population. *Ann Nutr Metab* **57**, 124–123.
34. Hu FB (2002) Dietary pattern analysis: a new direction in nutritional epidemiology. *Curr Opin Lipidol* **13**, 3–9.
35. Bendinelli B, Masala G, Saieva C, *et al.* (2011) Fruit, vegetables, and olive oil and risk of coronary heart disease in Italian women: the EPICOR Study. *Am J Clin Nutr* **93**, 275–283.
36. Polychronopoulos E, Pounis G, Bountziouka V, *et al.* (2010) Dietary meat fats and burden of cardiovascular disease risk factors, in the elderly: a report from the MEDIS study. *Lipids Health Dis* **9**, 30.