



Weeds

VOLUME 7
JULY 1959
NUMBER 3

Journal of the Weed Society of America

Weeds

Issued Quarterly by the Weed Society of America

C. J. WILLARD, *Editor*, Department of Agronomy, H. & F. Bldg., The Ohio State University, Columbus 10, Ohio.

W. C. JACOB, *Business Manager*, Department of Agronomy, University of Illinois, Urbana, Illinois.

EDITORIAL COMMITTEE

O. C. LEE, Department of Botany and Plant Pathology, Purdue University, Lafayette, Indiana.

R. A. PETERS, Department of Plant Science, University of Connecticut, Storrs, Connecticut.

W. C. ROBOCKER, Crops Research Division, ARS, USDA, State College of Washington, Pullman, Washington.

E. G. RODGERS, Department of Agronomy, University of Florida, Gainesville, Florida.

WEEDS is a quarterly journal published by the Weed Society of America. Editorial office is located at Department of Agronomy, H. and F. Bldg., The Ohio State University, Columbus 10, Ohio. Printing is by the W. F. Humphrey Press Inc., Geneva, New York. Subscription price is \$6.00 yearly for four issues; single copies \$1.50. Address all communications regarding subscriptions, advertising, and reprints to W. C. Jacob, Department of Agronomy, University of Illinois, Urbana, Illinois. Inquiries concerning information on manuscripts and other material for publication should be addressed to the Editorial Office. All checks, money orders, and other remittances should be made payable to the Weed Society of America.

Entered as second-class matter at the post office at Urbana, Illinois with additional entry at Geneva, New York.

Table of Contents

	<i>Page</i>
Oxalate and Mineral Contents of <i>Halogeton glomeratus</i> . Howard L. Morton, Robert H. Haas, and Lambert C. Erickson	255
Weeds in Alaska and Some Aspects of their Control. C. H. Dearborn	265
Amino Triazole for Control of Redroot in Cranberries. W. F. Meggitt and R. J. Aldrich	271
Effect of Amitrol Sprays on the Growth and Development of the Cranberry. Malcolm N. Dana	277
Residual Herbicidal Activity of Some Chloro-Substituted Benzoic Acids in Soil. W. M. Phillips	284
The Translocation of 2,4-D-1-C ¹⁴ in <i>Potamogeton pectinatus</i> , a Submersed Aquatic Plant. F. D. Aldrich and N. E. Otto	295
The Absorption and Translocation of C ¹⁴ -Labeled Simazin by Corn, Cotton, and Cucumber. D. E. Davis, H. H. Funderburk, Jr., and N. G. Sansing	300
Control of Certain Aquatic Weeds in Missouri Farm Ponds. Charles R. Walker	310
Chemical Weed Control in Seedling Alfalfa. II. Control of Broad-leaved Weeds. Paul F. Sand and M. K. McCarty	317
Logarithmic Sprayer Evaluations of Herbicides in Horticultural Crops. L. L. Danielson and Robert E. Wester	324
Effects of DCB, DCB-xylene Mixtures, Amitrol, and Sodium Salt of Dalapon in Irrigation Water on Corn and Rutabagas. V. F. Bruns and J. H. Dawson	333
A Three-year Study of the Effect of Some Phenoxy Herbicides on Cotton. W. K. Porter, Jr., C. H. Thomas, and J. B. Baker	341
Substituted Urea Herbicides: Their Electrophoretic Behavior and the Influence of Clay Colloid in Nutrient Solution on Their Phytotoxicity. C. W. Coggins, Jr. and A. S. Crafts	349
Brief Papers	
The Effect of Temperature and Light on Germination of Seed of Goosegrass, <i>Eleusine indica</i> . J. R. Fulwider and R. E. Engel	359
The Persistence of 4-(2-methyl-4-chlorophenoxy)butyric acid in Peas. H. A. Glastonbury, Margaret D. Stevenson, and R. W. E. Ball	362
Spectrofluorometric Determination of 1,1'-Ethylene 2,2'-bipyridylum Dibromide (FB/2). V. H. Freed and Robert E. Hughes, Jr.	364
Sustaining Members	365
News and Notes	366
Bibliography of Weed Investigations, October to December, 1958	367

Advertisers Index

duPont de Nemours & Co.	ii
U. S. Borax & Chemical Co.	iii
Stauffer Chemical Co.	iv
Chipman Chemical Co.	v
Spraying Systems Co.	vi
Union Carbide Chemicals Co.	vii

For Agriculture and Industry . . .

Du Pont WEED KILLERS

*offer new economies
and efficiency in killing
weeds, grass and brush*

KARMEX® diuron for weed control in many crops; irrigation and drainage ditches; and industrial areas.

KARMEX® DL for pre-emergence weed control in cotton.

TELVAR® monuron for weed control in many crops; irrigation and drainage ditches; and industrial areas.

KLOBEN neburon for weed and grass control in nursery plantings and tomatoes.

DYBAR fenuron—a pelleted product for dry application to the soil for control of brush in fence rows, drainage ditches, utility and railroad right-of-ways, and other non-cultivated areas.

TRYBEN 200, a new weed killer based on trichlorobenzoic acid, for control of bindweed and other noxious weeds; mixed broad-leaf weeds; and certain woody vines and brush.

AMMATE® X for long-term, low-cost brush control.



Better Things for Better Living *Through Chemistry*

**Four
easy
ways to
Destroy
Weeds**

1. UREABOR®

A nonselective, granular complex of sodium borate and substituted urea. Low application rates are a feature. Apply with the special new PCB Spreader for best results.

2. DB® Granular

A combination of 2,4-D and sodium borates. Kills deep-rooted, noxious weeds. Low application rates for maximum control with the utmost economy; use the PCB Spreader.
(Not intended for control of grass.)

3. POLYBOR-CHLORATE®

Highly soluble; for spray or dry application. It gives a quick knock-down; destroys top growth and roots. A general nonselective herbicide.

4. Concentrated BORASCU®

A nonselective, granular material. Apply by hand or with a mechanical spreader. Long residual action.

When you want Nonselective Herbicides

for Dependable Action

*look to United States Borax
& Chemical Corporation*

PACIFIC COAST BORAX COMPANY DIVISION



630 Shatto Place, Los Angeles 5, Calif.

New label granted **Eptam**[®] herbicide

EPTAM (EPTC) is now recommended for use in the following crops:

Snap beans	Forage legumes
Dry beans (except limas)	Irish potatoes
Table and sugar beets	Established Strawberries
Carrots	Established Tomatoes
Field and sweet corn	Established Nursery Stock
Flax	Established Ornamentals

For control of the following:

Annual Grasses: Annual blue grass, crab grass, crow-foot grass, foxtails (barley, bristly, green, giant and yellow), goose grass, oats, old witch grass, rye grass, sandbur, stinkgrass, water grass (barnyard grass), wild cane (black amber cane) and wild oats.

Perennials: Johnson grass (seedlings), nutgrass (yellow and purple), quackgrass (seedlings and established).

Broadleaves: Careless weed, chickweed (common), corn spurry, cud weed, dead nettle (henbit), fiddleneck, Florida purslane, kochia, lamb's quarters, pigweed (prostrate and red-root), purslane, ragweed, shepherd's purse and smartweed.

EPTAM should be thoroughly incorporated into the soil immediately by rotary tilling, disking or spike-tooth harrowing.

Broadcast applications of either spray or granular EPTAM give effective weed control of the entire treated area. For row crops such as corn, EPTAM can be applied in bands at the time of planting for control in the rows. Incorporation into the soil can then be accomplished by shallow disking in the same direction as the rows.

Testing of EPTAM to control grasses and weeds in other crops is continuing. For full information, write to the address below.

®EPTAM is Stauffer Chemical Company's trade-mark for ethyl di-n propylthiolcarbamate (EPTC), a herbicide.



STAUFFER CHEMICAL COMPANY
Sales Development Department
Agricultural Chemicals Division
380 Madison Avenue, New York 17, N. Y.



ATLACIDE: Safer chlorate weed killer...widely used for non-selective eradication of bindweed, Canada thistle, quack grass, Johnson grass and other tough perennials. Kills roots...discourages regrowth. Applied dry or dissolved in water.

ATLACIDE-2,4-D: Offers dual killing action of sodium chlorate and 2,4-D.

CHLOREA: A combination of sodium chlorate, borate and monuron in powder form. Kills weeds and grasses. Combines the proven effectiveness of chlorate on deep-rooted weeds with the soil-surface action of monuron on shallow-rooted grasses and annual seedling growth. Lasting residual effect inhibits new growth. Does not create a fire hazard when used as directed. Applied dry or as a water-mixed spray.

CHLOREA GRANULAR Similar to Chlorea, but a "pelletized" material. No mixing...easy to apply by hand or with mechanical spreader.

CHLORAX "40": A composition of sodium chlorate and borate...for weed and grass control. Has lasting residual effect. Does not create a fire hazard. Applied dry or as a spray.

ATLAS "A": A 40% sodium arsenite solution (4 lbs. arsenic trioxide per gal.). Destroys vegetation and algae in ponds and lakes. Controls certain turf weeds. Used as general weed killer and to kill trees and stumps. Also used to kill potato vines prior to harvesting.

SODIUM ARSENITE: A powder containing 75% arsenic trioxide. Used for the same purposes as Atlas "A". Applied dry or as a spray.

CHIP-CAL: New low-lime calcium arsenate formulation in granular and spray powder forms. Used as crabgrass preventatives for established turf. Applied any time before crabgrass seed starts to germinate.

2,4-D & 2,4,5-T: 2,4-D Amine and 2,4-D Ester liquids; 2,4-D Ester dusts; Low Volatile 2,4,5-T and Brush Killer.

METHOXONE: Contains MCP sodium salt. Used for control in small grains, flax, rice and grass. Controls same weeds as 2,4-D; considered safer for selective spraying.

BUTOXONE: A liquid containing 4-(2,4-DB). For selective control of many broadleaf weeds in established alfalfa and certain other legumes.

Write for Weed Control Booklets

CHIPMAN CHEMICAL COMPANY, INC.

Chicago, Ill.

Bound Brook, N. J.

Portland, Ore.

Palo Alto, Calif.

Pasadena, Tex.


Bessemer, Ala.

Manufacturers of Weed Killers Since 1912




the precision nozzle for effective spraying

Supplied in a full range of interchangeable orifice tip and strainer sizes to meet every capacity requirement. TeeJet Spray Nozzles for Weed Control by spraying make it possible to take maximum advantage of the chemical and sprayer unit. TeeJet nozzles are precision built and provide a flat spray with uniform distribution. Atomization is properly controlled to give coverage with an absolute minimum of driftage. Patented tip design, with set-back orifice opening protects precision orifice from accidental damage. TeeJet spray nozzles are built for use on spray booms and portable sprayers.




TEEJET SPRAY NOZZLE male pipe connection



TEEJET SPRAY NOZZLE female pipe connection

OFF-CENTER SPRAY NOZZLES

Spraying Systems Spray Nozzles with TeeJet tips are supplied in a variety of special body types to meet any unusual spraying requirement. For example, one type of off-center spray nozzle with swivel body provides a flat spray up to 35 feet wide for spraying areas with a single nozzle, that are not accessible with a boom.



INTERCHANGEABLE ORIFICE TIPS flat and cone spray types

SUPPLEMENTARY EQUIPMENT

Complete accessories relating to nozzle use are supplied. These include strainers, special nozzle fittings, and hand valve equipment.

TeeJet Spray Nozzles are supplied for Weed Control... as well as all other types of agricultural spraying. For complete information and reference data write for Catalog 30.

SPRAYING SYSTEMS CO.
Engineers and Manufacturers

3296 RANDOLPH STREET BELLWOOD, ILLINOIS

CRAG Mylone
BRAND SOIL FUMIGANT

HELPS

FOREST TREE SEEDLINGS

**GET OFF TO A
BETTER START**


In numerous tests throughout the country, formulations with CRAG Mylone soil fumigant have given excellent control of weeds, nematodes, and soil fungi in forest tree seed beds. Seedlings have a better chance to grow without interference from these pests.

When formulated, Mylone is an easy-to-handle powder that can be applied to the soil with a fertilizer spreader or as a drench. Application should be made at least three weeks before seeds are planted, except tobacco seed. Soil mixing is not necessary; no plastic cover is needed over the bed.

Formulations of CRAG Mylone are now commercially available for use in certain ornamental propagating beds, tobacco seed beds, and in tomato, pepper, egg plant, lettuce, and cabbage seed beds. It is also sold for weed and dry rot control in gladiolus in Florida. Experimental work is continuing with other crops.

Write to the address below for formulations in test quantities or more information. Names and addresses of commercial formulators are also available.

"Crag", "Mylone", and "Union Carbide" are trade marks of Union Carbide Corporation.



**FORMULATIONS CONTAINING
MYLONE NOW AVAILABLE
FOR VEGETABLE SEED BEDS**

A commercial label has been accepted for Mylone pre-planting use in seed beds of certain vegetables. Growers can now use it for the control of weeds, nematodes, and soil fungi in tomato, pepper, cabbage, egg plant, and lettuce seed beds.



CRAG Agricultural Chemicals
Union Carbide Chemicals Company

Division of Union Carbide Corporation
180 South Broadway, White Plains, New York