

Disadvantaged pre-schoolers attending day care in Salvador, Northeast Brazil have a low prevalence of anaemia and micronutrient deficiencies

Rebecca L Lander¹, Karl B Bailey¹, Alastair G Lander¹, Abdulmonem A Alsaleh², Hugo C Costa-Ribeiro³, Angela P Mattos³, Danile L Barreto³, Lisa A Houghton¹, Ian M Morison², Sheila M Williams⁴ and Rosalind S Gibson^{1,*}

¹Department of Human Nutrition, University of Otago, PO Box 56, Dunedin, New Zealand: ²Department of Pathology, University of Otago, Dunedin, New Zealand: ³Hospital Universitário Professor Edgard Santos, Fima Lifshitz Research Unit, Salvador, Bahia, Brazil: ⁴Department of Preventive and Social Medicine, University of Otago, Dunedin, New Zealand

Submitted 27 November 2012: Final revision received 23 June 2013: Accepted 25 July 2013: First published online 5 September 2013

Abstract

Objective: To examine the micronutrient status of disadvantaged pre-schoolers from Northeast Brazil, following the introduction of pro-poor policies, by assessing the prevalence of anaemia and micronutrient deficiencies and the role of sociodemographic factors, genetic Hb disorders and parasitic infections.

Design: In a cross-sectional study, data on sociodemographic status, health, growth, genetic Hb disorders, parasites and nutrient supply from day-care meals were obtained. Fasting blood samples were collected and analysed for Hb, serum ferritin, transferrin receptor, folate, vitamin B₁₂, retinol, Zn and Se.

Setting: Seven philanthropic day-care centres serving urban slums in Salvador, Northeast Brazil.

Subjects: Pre-schoolers aged 3–6 years from disadvantaged households.

Results: Of the 376 sampled children, 94% were of black or mixed race; 33% and 29% had at least one genetic Hb disorder and intestinal parasite, respectively. Stunting and underweight were ≤5%; 14% were overweight. Day-care centres supplied micronutrient-dense meals and snacks each weekday. Less than 10% of pre-schoolers had anaemia and micronutrient deficiencies. Predictors ($P < 0.05$) of Hb were α^{3-7} thalassaemia, Se and retinol (but not ferritin). Micronutrient predictors ($P < 0.05$) were: elevated α_1 -glycoprotein for ferritin, Hb AS and BMI Z-score >1 for transferrin receptor, Zn and elevated α_1 -glycoprotein for retinol, sex and helminths for Se, helminths for vitamin B₁₂, and *Giardia intestinalis* infection for serum folate.

Conclusions: Impaired growth, anaemia and micronutrient deficiencies were uncommon among these disadvantaged pre-schoolers attending day care. A range of interventions including provision of micronutrient-dense, fortified day-care meals, deworming and vitamin A supplementation likely contributed to improved micronutrient status, suggesting expanded coverage of these programmes.

Keywords

Anaemia
Micronutrient deficiencies
Predictors
Pre-schoolers
Disadvantaged
Northeast Brazil

Many young children living in urban poverty today are vulnerable to malnutrition⁽¹⁾. In Brazil, the Northeast (NE) region is the poorest in the country and where young children living in low-income urban slums are at increased risk of anaemia, micronutrient deficiencies and morbidity^(2–4). Together, these deleterious effects can have a long-lasting adverse impact on the cognitive and developmental potential of young children⁽⁵⁾.

Numerous diet and host-related factors may contribute to micronutrient malnutrition among urban slum pre-schoolers in NE Brazil. Their staple diets are based on

cereals and legumes, and are devoid of expensive micronutrient-dense animal-source foods. Thus, the adequacy of key essential micronutrients such as bioavailable Fe and Zn, preformed vitamin A and vitamin B₁₂ may be compromised by low intakes and/or poor bioavailability from these plant-based diets⁽⁶⁾. In urban slum settings, the water supply and sanitation are poor⁽²⁾, so intestinal parasitic infections are often widespread⁽⁷⁾. Such infections can exacerbate poor micronutrient bioavailability by increasing permeability and reducing transit time in the intestine⁽⁸⁾. Furthermore, in Salvador,

*Corresponding author: Email rosalingibson@otago.ac.nz

NE Brazil, 80% of the population is of West African descent⁽⁹⁾. Hence, some disadvantaged pre-schoolers in Salvador are likely to be at risk for genetic Hb disorders, particularly thalassaemias and sickle cell disease⁽¹⁰⁾, which can be associated with anaemia and Zn deficiency, respectively^(11,12).

Over recent decades, Brazil has implemented several key pro-poor policies to improve the micronutrient status, health, physical and cognitive development of urban pre-schoolers living in low-income environments. Policies include supplementation of young children with vitamin A and Fe⁽¹³⁾, mandatory fortification of wheat and corn flour with Fe and folic acid⁽¹³⁾, implementation of sewage and sanitation programmes⁽⁷⁾, and promotion of day-care facilities for the care and education of pre-schoolers⁽¹⁴⁾.

Nevertheless, there have been few comprehensive studies that have examined the micronutrient status of disadvantaged urban pre-schoolers attending day care in NE Brazil since the implementation of these policies. Consequently, little is known about the prevalence of micronutrient deficiencies among children in these settings today. In earlier reports, we have examined the growth, intestinal parasitic infections and adequacy of the nutrient supply from day-care meals in a group of pre-schoolers living in urban slums in Salvador, NE Brazil^(15,16). Here, we extend the research by investigating the prevalence of anaemia and micronutrient deficiencies in these same children and exploring the major predictors of Hb and micronutrient biomarkers.

Methods

Study sites and participants

The present cross-sectional study was conducted between August and November 2010 in Salvador, in the state of Bahia, NE Brazil. Seven philanthropic pre-school day-care centres within the city centre and peri-urban areas participated, as described earlier⁽¹⁵⁾. These day-care centres were selected based on the criteria that all the children in attendance were from poor communities and participated full-time (i.e. 07.30–17.00 hours). The service provided was without charge and provided five meals per day, with an on-site dietitian.

The children (n 376) enrolled in the day-care centres (maximum class size of twenty-five children) were from low-income families and attended day care five days per week, except holidays, until school age. The day-care centres supplied filtered drinking water and the majority of the weekday food intake for the children, and provided flush toilets, hand basins and shower facilities. Inclusion criteria for the study were apparently healthy children enrolled in the day-care classes for 3- and 4-year-olds in the 2010 school year (February to December). The study protocol was approved by the Human Ethics Committees of the Federal University of Bahia, Salvador and the University of Otago, New Zealand. Informed

written permission to participate in the study was given by the parents or primary guardians of the children.

Questionnaire and assessment of children's growth, parasite status and day-care meals

Data on sociodemographic status and health of the participants were collected via a culturally appropriate structured questionnaire, administered by trained research assistants. Methodological details are provided elsewhere⁽¹⁵⁾. Sociodemographic data were used to calculate an overall socio-economic status (SES) score for each participant based on a model designed to assess the poverty level of Brazilian families in low socio-economic urban populations⁽¹⁷⁾. Briefly, points were assigned for family and housing size and structure, parental education and occupation, marital status, ownership of house and household assets, toilet and sewage facilities, type of drinking water, availability of electricity and susceptibility of the house to flooding during heavy rain, and used to calculate two SES categories: extremely low and low SES, based on scores ≤ 34 and ≥ 35 , respectively.

Health status variables of the child supplied by maternal report included coffee intake within the past 24 h, smoking in the home and history of asthma in the mother or siblings. The mother also reported on the use of Fe supplements and deworming treatment within the past 6 months, whereas details of vitamin A supplementation was obtained from the child's health card. Ethnicity of the child was determined by skin colour, hair and facial characteristics, which were recorded by the research assistants, as performed in the national census⁽⁹⁾.

Weight and height measurements were taken with children wearing light clothes and no shoes using standardized techniques and calibrated equipment⁽¹⁸⁾. Standardized Z-scores for height-for-age (HAZ), weight-for-age (WAZ), weight-for-height (WHZ) and BMI (BMIZ) were calculated using the WHO 2006/2007 growth reference data^(18,19). Overweight and obesity in the children was defined as BMIZ >1 to ≤ 2 and BMIZ >2 , respectively.

Of the pre-schoolers, 86% (325/376) provided a stool sample which was examined by microscopy for helminths and protozoan intestinal parasites. Positive samples for *Giardia intestinalis* were identified by an enzyme immunoassay. Major food sources and adequacy of the micronutrient supply from twenty daily day-care menus was evaluated by procedures described earlier⁽¹⁶⁾.

Biochemical assessment

Morning fasting peripheral venepuncture blood samples were obtained with participants in a sitting position, after applying a topical local vasodilator anaesthetic (amethocaine; AmetopTM) to minimize any discomfort. Blood was drawn into a trace-element-free evacuated tube (Becton Dickinson, Franklin Lakes, NJ, USA) for the micronutrient and infection biomarkers and into a paediatric evacuated tube containing EDTA (Becton Dickinson) for a complete

blood count and for testing for genetic Hb disorders. All blood samples were refrigerated immediately following collection⁽²⁰⁾, protected from UV light, and the serum separated within 2 h using trace-element-free techniques. Aliquots of serum and washed red blood cells were frozen within 10 min, initially at -30°C and later at -70°C . An aliquot of EDTA-anticoagulated whole blood was haemolysed by a 1:10 dilution in 1% ascorbic acid solution (w/w) and frozen for erythrocyte folate analysis⁽²¹⁾. Frozen samples were shipped on dry ice to the University of Otago, New Zealand for analysis.

Serum ferritin was determined on an Elecsys 2010 auto-analyser (Roche, New Zealand; CV = 3%) using a Ferritin Elecsys reagent kit (Roche Diagnostics GmbH, Mannheim Germany) and soluble transferrin receptor (sTfR) via an enzyme immunoassay (Ramco Laboratories Inc., Houston, TX, USA; CV = 7%). Serum Zn was analysed by flame atomic absorption spectrophotometry (ContraAA 700; Analytik Jena AG, Jena, Germany; CV = 5%)⁽²²⁾ and serum Se by electrothermal atomic absorption spectrophotometry (AA-800, Perkin Elmer 2690; Ebos Group Ltd, Auckland, New Zealand; CV = 7%)⁽²³⁾. Serum retinol was analysed by HPLC⁽²⁴⁾ (CV = 2%) and serum vitamin B₁₂ by an electrochemiluminescence immunoassay (CV = 4%) using a Vitamin B₁₂ Elecsys reagent kit (Roche Diagnostics GmbH) on an Elecsys 2010 auto-analyser (Roche, New Zealand). Serum and whole-blood folate concentrations were measured by microbiological assay⁽²⁵⁾ in ninety-six-well microtitre plates using chloramphenicol-resistant cryopreserved *Lactobacillus rhamnosus* (ATCC 27773; American Type Culture Collection, Manassa, VA, USA) and 5-methyltetrahydrofolate as the calibrator (Merck & Cie, Schaffhausen, Switzerland; CV = 14%). Erythrocyte folate concentrations were calculated from whole blood values by using individual packed cell volumes and correction for serum folate concentration. Serum C-reactive protein (CRP) and α_1 -glycoprotein (AGP) concentrations were assayed by immunoturbidimetry (Roche Diagnostics GmbH) on a Cobas Mira II auto-analyser (CV = 10% and 3%, respectively). The precision of the biochemical assays was checked using a pooled serum sample and their accuracy established using certified reference materials or appropriate manufacturers' controls; values fell within the certified ranges.

The complete blood count was determined using an automatic electronic analyser (Coulter LH 750 Hematology Analyzer; Beckman Coulter Inc., São Paulo, Brazil) in the Professor Edgar Santos Hospital haematology laboratory, Salvador. Haemoglobinopathies were detected by alkaline and acid Hb electrophoresis analysis (Sebia Hydrasys Electrophoresis Analyzer; Sebia Inc., Norcross, GA, USA), which separated normal haemoglobins (A, A₂ and F) and detected major Hb variants, including the heterozygous variant of Hb S (Hb AS) and Hb C (Hb AC). The presence of α^{3-7} thalassaemia was determined by using a multiplex PCR reaction on extracted DNA⁽²⁶⁾. Positive Hb

and DNA controls were used for the electrophoresis analysis and PCR reactions, respectively. A negative control confirmed the absence of contamination for the PCR results, as described elsewhere⁽²⁷⁾.

Anaemia and other haematological disturbances were defined using the following interpretive criteria: Hb <110 g/l and <115 g/l for children aged <5 years and ≥ 5 years, respectively⁽²⁸⁾; mean cell volume (MCV) <73 fl and <74 fl for children aged <5 years and ≥ 5 years, respectively⁽²⁸⁾; and red cell distribution width (RDW) >14%⁽²⁹⁾. For storage Fe depletion, serum ferritin was defined as <12 $\mu\text{g/l}$ and <15 $\mu\text{g/l}$ for children aged <5 years and ≥ 5 years, respectively⁽²⁸⁾. Tissue Fe deficiency was defined as sTfR >8.5 mg/l⁽³⁰⁾. Hb AS and Hb AC, and α^{3-7} thalassaemia, homozygous or heterozygous, were identified as present or absent.

Interpretive criteria to define micronutrient deficiencies were: serum Zn <9.9 $\mu\text{mol/l}$ ⁽³¹⁾, Se ≤ 0.82 $\mu\text{mol/l}$ ⁽³²⁾, retinol <0.70 $\mu\text{mol/l}$ ⁽³³⁾ and vitamin B₁₂ <150 pmol/l⁽³⁴⁾. Marginal vitamin A status was defined as serum retinol ≥ 0.7 but <1.05 $\mu\text{mol/l}$ ⁽³⁵⁾. Low serum folate concentration was defined as <6.8 nmol/l and low erythrocyte folate as <317 nmol/l⁽³⁶⁾. Acute and chronic inflammation were assessed by serum CRP >5 mg/l⁽³⁷⁾ and AGP >1.0 g/l⁽³⁸⁾, respectively.

Statistical analysis

Selected characteristics of the children and households, and prevalence of intestinal parasites and genetic Hb disorders are presented as percentages. Means and standard deviations were calculated for haematology and micronutrient biomarkers, and adjusted where necessary for inflammation (i.e. ferritin and retinol)^(39,40). Associations between elevated AGP concentrations (>1 g/l) and BMIZ >1 and between the presence of genetic Hb disorders and ethnicity were investigated using Fisher's exact test. AGP was used as a dichotomized variable because the precision of the assay precluded using the data as a continuous variable. Data were log-transformed if the variable had a strongly skewed distribution (e.g. ferritin, sTfR, retinol and serum folate). Multiple regression models included variables if associations using univariate regression analysis were $P < 0.2$. The sandwich estimator was used to obtain robust standard errors, to account for the sampling procedure. Statistical analyses were carried out using the statistical software package STATA version 11.

Results

Sociodemographic, health, growth and parasite status

Of the 438 eligible children, 378 (86%) were recruited. Reasons for non-participation included children on the roll who had moved or were moving during the study, children who were chronically ill and not in regular

day-care attendance, and parental refusal. Mean age of the children was 4.2 (SD 0.61) years and 52% were boys (Table 1). Of the households, 48.4% (182/376) were classified as extremely low SES and 51.6% as low SES. Most of the pre-schoolers were black (42.2%) or mixed

race (i.e. brown; 51.8%); only 6.0% were white. More than 50% had received vitamin A supplements and deworming treatment, but less than 20% had reportedly been supplemented with Fe.

Prevalence of stunting and underweight was low ($\leq 5\%$), whereas that of overweight and obesity was 11.0% and 3.3%, respectively (Table 1). Stool samples were provided by 86% (325/376) of the pre-schoolers, of whom nearly 30% were infected with at least one intestinal parasite (Table 1). No significant differences in age, sex, sociodemographic status and deworming treatment were found between participants who provided a stool sample and those participants who did not ($n = 51$), except for the use of vitamin A supplements. Daily supply of almost all micronutrients from the day-care meals appeared adequate except for vitamin A, where risk of adequacy was $\leq 50\%$, also reported earlier⁽¹⁶⁾. Intake of animal-source foods from the day-care meals was 80 g/d.

Table 1 Sociodemographic, growth, health, parasite status, genetic Hb disorders and elevated biomarkers of inflammation among pre-schoolers aged 3–6 years from disadvantaged households, Salvador, Northeast Brazil, August–November 2010

	<i>n</i>	%
Age (years)	376	
Mean	4.2	
SD	0.61	
Sex (male)	196/376	52.1
SES (extremely low)	182/376	48.4
Ethnicity		
Black	154/365	42.2
Mixed (i.e. brown)	189/365	51.8
White	22/365	6.0
Growth measurements		
Stunted (HAZ < -2)	10/364	2.8
Underweight (WAZ < -2)	2/364	0.6
Overweight (BMIZ > 1 to ≤ 2)	40/364	11.0
Obese (BMIZ > 2)	12/364	3.3
Smoking		
Mother smoking in the house	47/376	12.5
Adult smoking in the house	89/376	23.7
History of asthma in mother or sibling	68/376	18.1
Coffee intake within past 24 h	149/376	39.6
Fe syrup within 6 months	70/376	18.6
Vitamin A supplementation ever received	200/376	53.2
Deworming treatment within 6 months	192/376	51.1
Parasite present	95/325	29.2
Helminths	58/325	17.8
<i>Trichuris trichiura</i>	39/325	12.0
<i>Ascaris lumbricoides</i>	34/325	10.5
Hookworm	3/325	0.9
<i>Giardia intestinalis</i>	42/325	12.9
Overall prevalence of selected Hb disorders	101/311	32.5
Haemoglobinopathies		
Hb AS	20/358	5.6
Hb AC	9/358	2.5
α^{3-7} Thalassaemia		
Heterozygous	70/306	22.9
Homozygous	6/306	2.0
AGP > 1.0 g/l	112/358	31.3
CRP > 5 mg/l	46/358	12.9

SES, socio-economic status; HAZ, height-for age Z-score; WAZ, weight-for-age Z-score; BMIZ, BMI Z-score; AGP, α_1 -glycoprotein; CRP, C-reactive protein.

Genetic Hb disorders, biomarkers of infection and haematology

Almost a third of the pre-schoolers had at least one genetic Hb disorder, of which the most prevalent was heterozygous α^{3-7} thalassaemia (23%). Prevalence of Hb AS and Hb AC was low (Table 1). Four children had both a haemoglobinopathy and α^{3-7} thalassaemia. Prevalence of genetic Hb disorders was independent of ethnicity. Hb AS tended to be higher in the black children (60%, 12/20) compared with those of mixed race (i.e. 30%, 6/20) or white (10%, 2/20), although the difference was not significant ($P = 0.084$). Only 13% had elevated CRP, although nearly a third had elevated AGP (Table 1). Overweight (i.e. BMIZ > 1) was more prevalent among those with AGP > 1 g/l compared to those with AGP ≤ 1 g/l (i.e. 49%, 24/49 *v.* 29%, 87/303; $P = 0.007$).

Significant differences existed across the Hb variants for Hb, MCV and RDW ($P < 0.001$), with Hb AA children having greater mean Hb and MCV, and lower RDW, compared with children with Hb AS and Hb AC and/or α^{3-7} thalassaemia. There were no significant differences across the groups for serum ferritin and sTfR (Table 2).

Table 2 Mean values (and standard deviation) for haematological variables and iron biomarkers for normal Hb type (Hb AA) and the four major abnormal Hb variants among pre-schoolers aged 3–6 years from disadvantaged households, Salvador, Northeast Brazil, August–November 2010

	Hb AA (<i>n</i> 210)		Hb AS (<i>n</i> 20)		Hb AC (<i>n</i> 9)		α^{3-7} Thalassaemia heterozygous (<i>n</i> 66)		α^{3-7} Thalassaemia homozygous (<i>n</i> 6)		<i>P</i>
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	
Hb (g/l)	124.1	7.1	122.5	6.6	121.9	13.5	119.4	6.5	111.8	4.4	<0.001
MCV (fl)	80.7	3.2	79.4	4.1	72.9	5.1	75.4	3.4	63.8	2.4	<0.001
RDW (%)	13.9	1.0	13.9	2.6	15.2	0.9	14.4	1.5	16.4	1.1	<0.001
Ferritin† (μ g/l)	41.3	19.9	43.6	18.5	34.1	10.2	43.6	17.2	38.3	14.4	0.084
sTfR (mg/l)	7.1	1.5	8.1	2.6	7.8	1.5	7.4	1.7	7.2	1.4	0.259

MCV, mean cell volume; RDW, red cell distribution width; sTfR, soluble transferrin receptor.

†Participants with concomitant Hb AS or Hb AC were excluded ($n = 4$).

‡Adjusted for infection⁽³⁹⁾.

Table 3 Prevalence (% and 95% confidence interval) of anaemia and micronutrient deficiencies, and mean values (and standard deviation) of specific micronutrient biomarkers, including adjusted mean values for those impacted by infection, among pre-schoolers aged 3–6 years from disadvantaged households, Salvador, Northeast Brazil, August–November 2010

	<i>n</i>	% or Mean	95% CI or SD
Hb			
<5 years of age (<110 g/l)	11/319	3.4	1.7, 6.1
≥5 years of age (<115 g/l)	3/40	7.5	1.5, 21.9
Serum ferritin			
<5 years of age (<12 µg/l)	6/318	1.9	0.7, 4.1
≥5 years of age (<15 µg/l)	1/40	2.5	0.1, 13.9
Serum sTfR >8.5 mg/l	54/358	15.1	11.5, 19.2
Serum Zn (µmol/l)	358	12.8	1.8
Low serum Zn (<9.9 µmol/l)	13/358	3.6	1.9, 6.1
Serum retinol (µmol/l)	358	1.09†	0.24
Low serum retinol‡ (<0.7 µmol/l)	10/358	2.8	1.3, 5.1
Marginal retinol deficiency (0.7–1.05 µmol/l)	158/358	44.1	38.9, 49.4
Serum Se (µmol/l)	358	0.99	0.14
Low serum Se (≤0.82 µmol/l)	34/358	9.5	6.7, 13.0
Serum vitamin B ₁₂ (pmol/l)	357	649	224
Low serum vitamin B ₁₂ (<148 pmol/l)	1/357	0.3	0.01, 1.7
Marginal vitamin B ₁₂ deficiency (148–221 pmol/l)	5/357	1.4	0.5, 3.3
Serum folate (nmol/l)	355	51	21
Low serum folate (<6.8 nmol/l)	0/355	0	0, 1.0
Erythrocyte folate (nmol/l)	355	1253	390
Low erythrocyte folate (<317 nmol/l)	0/355	0	0, 1.0

sTfR, soluble transferrin receptor.

†Adjusted for infection⁽³⁹⁾.

‡Adjusted for infection⁽³⁸⁾.

Anaemia and micronutrient deficiencies

Mean concentrations of Hb and the micronutrient biomarkers and the prevalence of anaemia and micronutrient deficiencies are shown in Table 3. Prevalence of anaemia and micronutrient deficiencies was <10%, and ranged from 0% for folate to 9.5% for Se (Table 3); 13% (46/358) had at least one micronutrient deficiency. The prevalence of anaemia was higher among children with at least one Hb disorder than among those without Hb disorders (i.e. 69%, 9/13 *v.* 31%, 4/13; $P=0.006$). Of the few children with depleted Fe stores (2%, 7/358), only one child had an Hb disorder. However, 15% (54/358) had elevated sTfR indicative of tissue Fe deficiency, of whom 30% (16/54) had an abnormal Hb variant.

Predictors of Hb and micronutrient biomarkers

Table 4 shows the significant predictors of Hb and biomarkers of Fe status. Both homozygous and heterozygous α^{3-7} thalassaemia variants were negative determinants of Hb ($P\leq 0.001$), whereas Se and retinol were positive determinants of Hb; serum ferritin was not a significant determinant of Hb. BMIZ > 1 and Hb AS were positive determinants of sTfR but had no effect on ferritin. Elevated AGP was a positive determinant of ferritin.

In contrast, AGP was a negative determinant of retinol, whereas Zn was a positive predictor of retinol (Table 5). Being male was positively associated with Se, whereas helminths were a negative predictor of Se and vitamin B₁₂. *G. intestinalis* was a positive predictor of serum but not erythrocyte folate. The associations between either

Hb or micronutrient biomarkers and age, sex (with the exception of Se), ethnicity, supplementation with Fe or vitamin A, deworming treatment, smoking in the home, family history of asthma and recent coffee intake were not statistically significant.

Discussion

Our findings highlight the low prevalence of anaemia and micronutrient deficiencies in these disadvantaged pre-schoolers attending philanthropic day-care centres. To our knowledge, these data are the first in Brazil to examine six micronutrient biomarkers in this age group concurrently and to investigate the complex interrelationships between Hb, micronutrient status, parasitic infections and genetic Hb disorders.

The low prevalence of anaemia reported here (i.e. <4%) was unexpected. Earlier studies of urban pre-schoolers in NE Brazil⁽⁴¹⁾, including Salvador⁽³⁾, have reported much greater anaemia rates, sometimes as high as 40%⁽⁴²⁾. Several factors may account for our unexpected finding. The most frequent genetic Hb disorder among these pre-schoolers was heterozygous α^{3-7} thalassaemia (23%), which, unlike the homozygous variant, is relatively benign and is not always associated with anaemia.

Moreover, review of the day-care meals indicated that the supply of all the micronutrients with a major role in the maintenance of normal haematopoietic function (i.e. Fe, folate and vitamin B₁₂), with the exception of

Table 4 Predictors of Hb and two biomarkers of iron status based on both univariate and multiple regression models, as shown by β coefficient (and 95% confidence interval), among pre-schoolers aged 3–6 years from disadvantaged households, Salvador, Northeast Brazil, August–November 2010

	Univariate β	Multivariate β	95% CI	P
Hb†				
α^{3-7} Thalassaemia homozygous	-12.29*	-12.19	-14.78, -9.60	<0.001
α^{3-7} Thalassaemia heterozygous	-4.84*	-5.28	-7.44, -3.11	0.001
Hb AS	-0.44			
Hb AC	-1.00			
Se	11.71*	11.71	6.26, 17.17	0.001
Retinol (log-transformed)	7.62*	5.08	0.65, 9.51	0.030
Ferritin (log-transformed)	2.11(*)	2.35	-0.76, 5.46	0.117
Zn	0.71*	0.42	-0.09, 0.92	0.092
AGP > 1 g/l	0.01(*)	0.29	-2.09, 2.68	0.779
Serum ferritin (log-transformed)‡				
Age	0.03			
BMIZ > 1	0.10*	0.04	-0.02, 0.10	0.138
AGP > 1 g/l	0.31*	0.30	0.16, 0.44	0.001
Hb AS	0.07			
Hb AC	-0.12			
α^{3-7} Thalassaemia heterozygous	0.04			
α^{3-7} Thalassaemia homozygous	-0.05			
Serum sTfR (log-transformed)§				
Age	-0.04*	-0.03	-0.06, 0.01	0.086
Sex	0.04			
BMIZ > 1	0.10*	0.09	0.00, 0.17	0.050
Helminths	-0.06(*)	-0.06	-0.14, 0.03	0.163
Hb AS	0.16*	0.17	0.04, 0.31	0.020
Hb AC	0.09			
α^{3-7} Thalassaemia heterozygous	0.03			
α^{3-7} Thalassaemia homozygous	0.01			

AGP, α_1 -glycoprotein; BMIZ, BMI Z-score.

All variables with $P < 0.2$ in the univariate analysis were included in the multivariate regression.

For all binary categorical variables, the reference was the absence of the condition specified.

(*) $P < 0.2$ based on univariate regression; * $P < 0.05$ based on univariate regression.

† R^2 for multivariate regression model = 0.242.

‡ R^2 for multivariate regression model = 0.098.

§ R^2 for multivariate regression model = 0.083.

vitamin A, met the requirements of the pre-schoolers. For example, animal-source foods provided in the day-care meals (80 g/d), provision of legumes and use of fortified cereal flours in meal preparation⁽¹⁶⁾ and some supplementation sources of Fe and vitamin A (albeit minimal coverage)⁽¹⁵⁾ contributed to the daily nutrient requirements in these children. Despite the seeming lack of impact of mandatory Fe fortification alone on anaemia in Brazilian pre-schoolers⁽⁴³⁾, these multiple strategies were most likely responsible, at least in part, for the lower prevalences of anaemia and vitamin A deficiency reported here compared with earlier studies of pre-schoolers attending day-care centres in NE Brazil^(41,44).

Notwithstanding the apparently low prevalence of micronutrient deficiencies, there was some evidence of marginal vitamin A (45%) and Se (9.5%) status. The existence of suboptimal vitamin A status was attributed to an inadequate supply of vitamin A from day-care meals together with rather poor coverage of vitamin A supplementation (i.e. 50%). Zn status may have also played a role in vitamin A status in view of the positive association between Zn and retinol observed here. This relationship is not unexpected; Zn has a role in the hepatic synthesis of retinol-binding protein, and thus in the transport and

utilization of retinol, although in other studies where a similar relationship has been observed, the prevalence of Zn deficiency has been higher⁽⁴⁵⁾.

Reasons for the suboptimal Se status are uncertain. It could be associated with low soil Se levels and thus low levels of Se in the major locally grown plant-based staples. However, we were unable to locate values for the concentrations of Se in locally grown foods, so the supply of Se from the day-care meals is unknown.

It is of interest that both serum retinol and Se each had independent and significant positive associations with Hb (Table 4). The positive relationship between retinol and Hb is attributed to the role of vitamin A in the mobilization of Fe from the spleen or liver into the circulation⁽⁴⁶⁾, a mechanism that may account for the low prevalence of storage Fe depletion (i.e. <3%) reported here. There are several plausible mechanisms whereby Se status might impact on Hb, including its affect on the activity of thioredoxin reductase, a selenoenzyme postulated to be implicated in the up-regulation of hepatic haem oxygenase-1 involved in haem catabolism⁽⁴⁷⁾. In addition, reduced activity of glutathione peroxidase, a selenoenzyme that may protect Hb against oxidation in red blood cells⁽⁴⁸⁾, can result in increased inflammation and

Table 5 Predictors of micronutrients based on both univariate and multiple regression models, as shown by β coefficient (and 95% confidence interval), among pre-schoolers aged 3–6 years from disadvantaged households, Salvador, Northeast Brazil, August–November 2010

	Univariate β	Multivariate β	95% CI	<i>P</i>
Serum retinol (log-transformed) [†]				
Age	-0.01			
Sex (male)	-0.04 ^(*)	-0.04	-0.09, 0.01	0.076
Zn	0.02*	0.02	0.003, 0.04	0.031
AGP > 1 g/l	-0.12*	-0.12	-0.17, -0.07	0.001
Helminths	-0.08			
<i>Giardia intestinalis</i>	-0.06 ^(*)	-0.06	-0.15, 0.03	0.167
Vitamin A supplementation	0.00			
Serum Zn [‡]				
Age	0.03			
Sex (male)	0.01			
Se	1.88 ^(*)	1.99	-0.38, 4.37	0.088
AGP > 1 g/l	-0.19			
Helminths	-0.61 ^(*)	-0.50	-1.33, 0.32	0.190
<i>Giardia intestinalis</i>	-0.27			
Serum Se [§]				
Age	0.03			
Sex (male)	0.04 ^(*)	0.05	0.01, 0.09	0.018
AGP > 1 g/l	-0.03 ^(*)	-0.04	-0.08, 0.01	0.080
Helminths	-0.05*	-0.06	-0.10, -0.03	0.006
<i>Giardia intestinalis</i>	-0.01			
Serum vitamin B ₁₂				
Age	-33.70 ^(*)	-33.06	-67.02, 0.90	0.055
Sex (male)	-62.25 ^(*)	-49.44	-144.02, 45.14	0.256
AGP > 1 g/l	38.33			
Helminths	-116.01*	-97.63	-188.33, -6.93	0.038
<i>Giardia intestinalis</i>	-70.56*	-49.12	-102.88, 4.64	0.068
Serum folate (log-transformed) [¶]				
Age	0.02			
Sex (male)	-0.12 ^(*)	-0.12	-0.26, 0.02	0.088
Helminths	0.04			
<i>Giardia intestinalis</i>	0.14 ^(*)	0.14	0.01, 0.28	0.041

AGP, α_1 -glycoprotein.

All variables with $P < 0.2$ in the univariate regression were included in the multivariate regression.

For all binary categorical variables, the reference was the absence of the condition specified.

^(*) $P < 0.2$ based on univariate regression; ^(*) $P < 0.05$ based on univariate regression.

[†] R^2 for multivariate regression model = 0.111.

[‡] R^2 for multivariate regression model = 0.041.

[§] R^2 for multivariate regression model = 0.073.

^{||} R^2 for multivariate regression model = 0.063.

[¶] R^2 for multivariate regression model = 0.036.

oxidative stress. This inflammatory response may also be induced by intestinal parasites and may account for the independent and negative impact of helminths on Se concentrations.

During inflammation, pro-inflammatory cytokines such as IL-6 and leptin stimulate an increase in circulating hepcidin produced by both the liver and adipose tissue, which in turn down-regulates Fe absorption independent of Fe status, leading to functional Fe deficiency⁽⁴⁹⁾. Chronic inflammation also accompanies overweight and obesity and thus may account for the observed positive relationship between overweight and sTfR (Table 4).

Parasitic infections are also known to play a role in the aetiology of anaemia. In the present study parasitic infections were not predictors of Hb, but instead were associated with vitamin B₁₂ and folate status of the pre-schoolers. Their effect, however, although negative for vitamin B₁₂, was positive for folate status. The negative relationship between vitamin B₁₂ status and infection with

helminths, and to a lesser extent *G. intestinalis*, is well recognized^(50,51). Such infections are not only associated with changes in small intestine morphology that lead to reductions in the absorptive surface⁽⁸⁾, but also with increased bacterial overgrowth that causes malabsorption, specifically of vitamin B₁₂⁽⁵¹⁾. Risk of bacterial overgrowth is especially high for Brazilian pre-school children living in urban slum environments⁽⁵²⁾, emphasizing the importance of the adequate supply of animal-source foods in the day-care meals.

The positive association between serum folate status and *Giardia* infection observed here was unexpected but is not implausible. Mandatory folate fortification of cereal flours may have created a folate-rich environment which provided optimal conditions for the adherence of *Giardia* trophozoites to the intestinal epithelium^(53,54). Alternatively, high levels of circulating unmetabolized folic acid from folate fortification of flour⁽⁵⁵⁾ have been associated with decreased innate immune function⁽⁵⁶⁾. This finding was

attributed to reduced natural killer cell cytotoxicity, a mechanism responsible for targeting invading pathogens. However, to date these findings have been observed only in postmenopausal women and need to be replicated in other studies. Clearly, further research is warranted to confirm the positive association noted here between serum folate and *Giardia* infection, and establish the underlying biological mechanism.

Our findings are based on a cross-sectional study and hence preclude causal inferences from being made. Sampling was restricted to seven philanthropically funded day-care centres that provide a free service, so results may not apply to other day-care settings. Unfortunately, very limited data exist from the Salvador city registration on the number of philanthropic, public and private day-care centres for children of this age group. To our knowledge, the philanthropic day-care centres included here are unique to Salvador. The public government day care is also usually free of charge, but like the private day care, provides fewer meals (i.e. four and three, respectively) than the philanthropically funded day-care centres studied here. Furthermore, although we investigated several genetic Hb disorders, we were unable to measure all genetic factors, including glucose-6-phosphate dehydrogenase deficiency, which may have affected Hb and micronutrient levels, especially among the pre-schoolers of West African descent. Nevertheless, we did investigate Hb, six micronutrients, micronutrient supply from day-care meals, acute and chronic inflammation, parasite and health status variables, which together provide a greater understanding of factors influencing Hb and micronutrient status in these disadvantaged pre-schoolers.

Conclusions

In conclusion, even though the pre-schoolers were living in urban slum settings, impaired growth and anaemia and micronutrient deficiencies were uncommon. These results are most likely a reflection of the provision of micronutrient-rich day-care meals fortified with Fe and folic acid, parasite control and vitamin A supplementation, and coverage of these programmes should be expanded. Nevertheless, the existence of functional Fe deficiency associated with overweight in these pre-schoolers is of concern, and highlights the importance of strengthening efforts to prevent the emerging problem of overweight in disadvantaged pre-schoolers in NE Brazil.

Acknowledgements

Sources of funding: The work was funded in part by the University of Otago Research Fund, but the Board administering the Fund had no role in the design of the study, the data analysis or the writing of this article. *Conflicts of interest:* None. *Authors' contributions:* R.L.L.

contributed to the research project design, conduct of the study, statistical analysis and manuscript draft; K.B.B., A.G.L. and A.A.A. contributed to the analysis of the specimens; H.C.C.-R. and A.P.M. contributed to the research study oversight and recruitment of the day-care centres; D.B.L. assisted with the conduct of the study; L.A.H. and I.M.M. provided oversight of analytical procedures and contributed to the manuscript draft; S.M.W. contributed to the statistical analysis and manuscript draft; R.S.G. contributed to the research project design, study oversight and manuscript draft. All authors have read and approved the final draft. *Acknowledgements:* The authors thank the day-care organizations Santa Casa de Misericórdia and Mansão do Caminho for their support during planning and implementation of this study, the parents of the participating children, and the coordinators of the seven day-care centres. They also thank all nutritionists from the Fima Lifshitz Research Unit of the Hospital Universitário Professor Edgard Santos who assisted with the data collection.

References

1. UNICEF (2012) *Children in an Urban World. State of the World's Children 2012*. New York: UNICEF.
2. Prado M, Strina A, Barreto M *et al.* (2003) Risk factors for infection with *Giardia duodenalis* in pre-school children in the city of Salvador, Brazil. *Epidemiol Infect* **131**, 899–906.
3. Assis A, Barreto M, Gomes G *et al.* (2004) Childhood anemia prevalence and associated factors in Salvador, Bahia, Brazil. *Cad Saude Publica* **20**, 1633–1641.
4. Martins M, Santos L & Assis A (2004) Prevalence of hypovitaminosis A among preschool children from north-eastern Brazil, 1998. *Rev Saude Publica* **38**, 537–542.
5. Grantham-McGregor S, Cheung Y, Cueto S *et al.* (2007) Developmental potential in the first 5 years for children in developing countries. *Lancet* **369**, 60–70.
6. Antunes M, Sichieri R & Salles-Costa R (2010) Food intake among children under three years of age in an area with high food insecurity. *Cad Saude Publica* **26**, 1642–1650.
7. Barreto M, Genser B, Strina A *et al.* (2010) Impact of a citywide sanitation program in northeast Brazil on intestinal parasites infection in young children. *Environ Health Perspect* **118**, 1637–1642.
8. Müller N & von Allmen N (2005) Recent insights into the mucosal reactions associated with *Giardia lamblia* infections. *Int J Parasitol* **35**, 1339–1347.
9. Instituto Brasileiro de Geografia e Estatística (2010) *Census 2010*. Rio de Janeiro: IBGE.
10. Bain B (2006) *Haemoglobinopathy Diagnosis*, 2nd ed. Oxford: Blackwell Publishing.
11. Zemel B, Kawchak D, Fung E *et al.* (2002) Effect of zinc supplementation on growth and body composition in children with sickle cell disease. *Am J Clin Nutr* **75**, 300–307.
12. Fung E (2010) Nutritional deficiencies in patients with thalassemia. *Ann N Y Acad Sci* **1202**, 188–196.
13. Brazil Ministry of Health (2007) *Deficiencies of Micronutrients. Cadernos de Atencao Basica*. Brasilia: Ministry of Health.
14. Brazil Ministry of Health (1996) *Public Law on the Rights and Basis of Education. Law 9.394, December 20, 1996*. Brasilia: Ministry of Health.
15. Lander R, Lander A, Houghton L *et al.* (2012) Factors influencing growth and intestinal parasitic infections in preschoolers attending philanthropic daycare centers

- in Salvador, Northeast region of Brazil. *Cad Saude Publica* **28**, 2177–2188.
16. Lander R (2012) The nutritional status of disadvantaged preschool children attending daycares in Salvador, Northeast Brazil: a cross-sectional study. PhD Thesis, University of Otago.
 17. Issler R, Guigliani E, Kreutz G *et al.* (1996) Poverty levels and children's health status: study of risk factors in an urban population of low socioeconomic level. *Rev Saude Publica* **30**, 506–511.
 18. WHO Multicentre Growth Reference Study Group (2006) WHO child growth standards based on length/height, weight and age. *Acta Paediatr* **95**, 76–85.
 19. de Onis M, Onyango AW, Borghi E *et al.* (2007) Development of a WHO growth reference for school-aged children and adolescents. *Bull World Health Organ* **85**, 660–667.
 20. Tamura T, Johnston K, Freeberg L *et al.* (1994) Refrigeration of blood samples prior to separation is essential for the accurate determination of plasma or serum zinc concentrations. *Biol Trace Elem Res* **41**, 165–173.
 21. Thurlow R, Winichagoon P, Green T *et al.* (2005) Only a small proportion of anemia in northeast Thai children is associated with iron deficiency. *Am J Clin Nutr* **82**, 380–387.
 22. Smith J, Butrimovitz G & Purdy W (1979) Direct measurement of zinc in plasma by atomic absorption spectroscopy. *Clin Chem* **25**, 1487–1491.
 23. Jacobson B & Lockitch G (1988) Direct determination of selenium in serum by graphite-furnace atomic absorption spectrometry with deuterium background detection and a reduced palladium modifier: age specific reference ranges. *Clin Chem* **34**, 709–714.
 24. Thurnham D, Smith E & Flora P (1988) Concurrent liquid-chromatographic assay of retinol, α -tocopherol, β -carotene, α -carotene, lycopene, and β -cryptoxanthin in plasma with tocopherol acetate as an internal standard. *Clin Chem* **34**, 377–381.
 25. Molloy A & Scott J (1997) Microbiological assay for serum, plasma, and red cell folate using cryopreserved, microtitre plate method. *Meth Enzymol* **281**, 43–53.
 26. Tan A, Quah T, Low P *et al.* (2001) A rapid and reliable 7-deletion multiplex polymerase chain reaction assay for alpha thalassemia. *Blood* **98**, 250–251.
 27. Alsaleh A (2011) Haemoglobinopathies and thalassemia among Brazilian children: a cross-sectional study. Masters Dissertation, University of Otago.
 28. World Health Organization (2001) *Iron Deficiency Anaemia: Assessment, Prevention and Control – A Guide for Programme Managers*. Geneva: WHO.
 29. Gibson R (2005) *Principles of Nutritional Assessment*, 2nd ed. Oxford: Oxford University Press.
 30. Cook J, Skikne B & Baynes R (1993) Serum transferrin receptor. *Annu Rev Med* **44**, 63–74.
 31. Hotz C, Peerson J & Brown K (2003) Suggested lower cut-offs of serum zinc concentrations for assessing zinc status: reanalysis of the second National Health and Nutrition Examination Survey data (1976–1980). *Am J Clin Nutr* **78**, 756–764.
 32. Thomson C (2004) Assessment of requirements for selenium and adequacy of selenium status: a review. *Eur J Clin Nutr* **58**, 391–402.
 33. de Pee S & Dary O (2002) Biochemical indicators of vitamin A deficiency: serum retinol and serum retinol binding protein. *J Nutr* **132**, 9 Suppl., 2895S–2901S.
 34. de Benoist B (2008) Conclusions of a WHO technical consultation on folate and vitamin B₁₂ deficiencies. *Food Nutr Bull* **29**, 2 Suppl., S238–S244.
 35. Ballew C, Bowman B, Sowell A *et al.* (2001) Serum retinol distributions in residents of the United States: third National Health and Nutrition Examination Survey, 1988–1994. *Am J Clin Nutr* **73**, 586–593.
 36. Pfeiffer C, Johnson C, Jain R *et al.* (2007) Trends in blood folate and vitamin B-12 concentrations in the United States, 1988–2004. *Am J Clin Nutr* **86**, 718–727.
 37. Thurnham D, Mburu A, Mwaniki D *et al.* (2005) Micronutrients in childhood and the influence of subclinical inflammation. *Proc Nutr Soc* **64**, 502–509.
 38. World Health Organization/Centers for Disease Control and Prevention (2007) *Assessing the Iron Status of Populations*, 2nd ed. Geneva: WHO.
 39. Thurnham D, McCabe G, Northrop-Clewes C *et al.* (2003) Effects of subclinical infection on plasma retinol concentrations and assessment of prevalence of vitamin A deficiency: meta-analysis. *Lancet* **362**, 2052–2058.
 40. Thurnham D, McCabe L, Haldar S *et al.* (2010) Adjusting plasma ferritin concentrations to remove the effects of subclinical inflammation in the assessment of iron deficiency: a meta-analysis. *Am J Clin Nutr* **92**, 546–555.
 41. Vieira A, Diniz A, Cabral P *et al.* (2007) Nutritional assessment of iron status and anemia in children under 5 years old at public daycare centers. *J Pediatr (Rio J)* **83**, 370–376.
 42. Jordão R, Bernardi J & Filho A (2009) Prevalence of iron-deficiency anaemia in Brazil: a systematic review. *Rev Paul Pediatr* **27**, 90–98.
 43. Assunção M, Santos I, Barros A *et al.* (2012) Flour fortification with iron has no impact on anaemia in urban Brazilian children. *Public Health Nutr* **15**, 1796–1801.
 44. Paiva A, Rondo P, Gonçalves-Carvalho C *et al.* (2006) Prevalence and factors associated with vitamin A deficiency in preschool children from Teresina, Piauí, Brazil. *Cad Saude Publica* **22**, 1979–1987.
 45. Muñoz E, Rosado J, Lopez P *et al.* (2000) Iron and zinc supplementation improves indicators of vitamin A status of Mexican preschoolers. *Am J Clin Nutr* **71**, 789–794.
 46. Fishman S, Christian P & West KJ (2000) The role of vitamins in the prevention and control of anaemia. *Public Health Nutr* **3**, 125–150.
 47. Mostert V, Hill K & Burk R (2003) Loss of activity of the selenoenzyme thioredoxin reductase causes induction of hepatic heme oxygenase-1. *FEBS Lett* **541**, 85–88.
 48. Nagababu E, Chrest F & Rifkind J (2003) Hydrogen-peroxide-induced heme degradation in red blood cells: the protective roles of catalase and glutathione peroxidase. *Biochim Biophys Acta* **1620**, 211–217.
 49. Yanoff L, Menzie C, Denkinger B *et al.* (2007) Inflammation and iron deficiency in the hypoferrremia of obesity. *Int J Obes (Lond)* **31**, 1412–1419.
 50. Solomons N (1993) Pathways to the impairment of human nutritional status by gastrointestinal pathogens. *Parasitology* **107**, Suppl., S19–S35.
 51. Allen L, Rosado J, Casterline J *et al.* (1995) Vitamin B-12 deficiency and malabsorption are highly prevalent in rural Mexican communities. *Am J Clin Nutr* **62**, 1013–1019.
 52. dos Reis J, de Moraes M, Oliva C *et al.* (2007) Breath hydrogen tests in the diagnosis of environmental enteropathy in children living in an urban slum. *Dig Dis Sci* **52**, 1253–1258.
 53. Sousa M, Gonçalves C, Bairos V *et al.* (2001) Adherence of *Giardia lamblia* trophozoites in Int-407 human intestinal cells. *Clin Diagn Lab Immunol* **8**, 258–265.
 54. Khademi R, Ghaffarifar F & Asl H (2006) *In vitro* effect of folic acid and cobalamin (vitamin B₁₂) on adhesion and growth of *Giardia lamblia*. *Iran J Parasitol* **1**, 47–52.
 55. Smith A (2010) Folic acid nutrition: what about the little children? *Am J Clin Nutr* **91**, 1408–1410.
 56. Troen A, Mitchell B, Sorensen B *et al.* (2006) Unmetabolized folic acid in plasma is associated with reduced natural killer cell cytotoxicity among postmenopausal women. *J Nutr* **136**, 189–194.