

baseline levels ($P < .05$), whereas counts returned to baseline levels by day 4 in the quaternary ammonium-alcohol disinfectant group. Recovery of MRSA and enterococci was significantly reduced on days 1–7 in the continuously active disinfectant group versus untreated controls (3 of 93, 3% vs 20 of 97, 21% respectively; $P = .002$), but not in the quaternary ammonium-alcohol disinfectant (11 of 97, 11%; $P = .12$). **Conclusions:** A single spray application of a continuously active disinfectant resulted in sustained reductions in total aerobic colony counts over 7 days and reduced recovery of MRSA and enterococci. The continuously active disinfectant could potentially reduce the risk for transmission of pathogens by portable devices.

Funding: None

Disclosures: None

Doi:10.1017/ice.2020.770

Presentation Type:

Poster Presentation

Evaluation of a Disease State Stewardship Intervention for Urinary Tract Infections at an Academic Medical Center

Xhilda Xhemali, University of Kentucky HealthCare; Derek Forster, University of Kentucky School of Medicine; Bryant Clemons, University of Kentucky HealthCare; Sarah Cotner, University of Kentucky HealthCare; Jeremy VanHoose, University of Kentucky HealthCare; Donna Burgess, University of Kentucky HealthCare; David Burgess, University of Kentucky College of Pharmacy; Thein Myint, University of Kentucky HealthCare; Mitu Karki Maskey, University of Kentucky HealthCare; Katie Wallace, University of Kentucky HealthCare

Background: Urinary tract infections (UTIs) are often misdiagnosed and mismanaged. Disease state stewardship initiatives targeting UTIs through the development of institutional guidelines and real-time prospective audit and feedback (PAAF) on provider management may have a significant impact on the overuse of antimicrobials. **Objective:** Our study evaluated the effectiveness of a UTI focused disease state stewardship intervention by assessing institutional guideline adherence before and after implementation. **Methods:** This retrospective quasi-experimental study was conducted at a tertiary-care academic medical center. Patients ≥ 18 years of age receiving antimicrobials for a UTI were included. A previously performed retrospective review of UTI management from September–November 2017 was used as the baseline. The UTI management guideline was implemented in July 2018, and service lines were educated. A PAAF initiative began in June 2019, whereby the antimicrobial stewardship team performed daily reviews of patients receiving antimicrobials for UTIs. Data was collected on their management, and providers were contacted in real time with recommendations based on the institutional guideline. Patients reviewed June–October 2019 were included in the postimplementation analysis. Patients were excluded if they were pregnant, underwent a urological procedure with risk of mucosal bleeding, or were an outside hospital transfer already on UTI therapy. The primary outcome of this study was to evaluate guideline adherence before and after the implementation of PAAF for the management of UTIs. **Results:** In total, 198 patients in the pre-intervention group and 246 in the PAAF group were included. The emergency department was the primary ordering service of urinalyses (60.1% vs 66.1%; $P = .2287$) in both periods and altered mental status as the main indication for testing (35.2% vs 31.3%; $P = .5465$). Treatment of asymptomatic bacteriuria and pyuria decreased significantly between the 2 periods: 74.8% versus

36.2% ($P = .0001$). Appropriate ordering of urinalyses (33.8% vs 68.3%; $P = .0001$) and urine cultures (29.3% vs 61.0%; $P = .0001$) also improved in the PAAF group. Recommendations made during PAAF included therapy discontinuation (66.7%), antimicrobial therapy change (15.5%), or duration modification (15.5%), and 59.5% of first interventions were accepted. Overall guideline compliance significantly improved from 13.1% in the pre-intervention period to 26.1% in the PAAF period ($P = .0011$).

Conclusions: A UTI disease state intervention was associated with significant reductions in the treatment of asymptomatic presentations as well as an improvement in overall guideline adherence. We believe that this approach represents a powerful stewardship strategy for decreasing unnecessary antimicrobial usage.

Funding: None

Disclosures: None

Doi:10.1017/ice.2020.771

Presentation Type:

Poster Presentation

Evaluation of a Sick Employee Online Log System for Tracking Sick Hospital Employees During Two Influenza Seasons

William Cleve, Vidant Medical Center; Kathy Cochran, Vidant Medical Center; Keith M. Ramsey, Vidant Medical Center

Background: Since 2009, Vidant Health has used a Sick Employee Online Log (SEOL) system to track illnesses among employees and to capture this information in real time. The CDC assessed the 2017–2018 influenza season as a high-severity influenza season, whereas the 2018–2019 influenza season was of moderate severity. **Objective:** In this research project, we sought to determine whether the influenza season severity would affect either the hospital-based employee illness surveillance system results or would correlate with state influenza-like illness (ILI) visits. **Methods:** The SEOL system is an internet-based system initiated in December 2008. When a hospital employee calls in sick, the department manager records whether the employee reports the following symptoms: nausea, vomiting, diarrhea, upper respiratory infection, fever, sore throat, headache, conjunctivitis, rash, and/or cough. The information is confidential, with raw data access restricted to review by occupational health and infection control leadership. The correlation value was determined for each symptom using the North Carolina Division of Human Services (NC DHHS) percentage of ILI visits in statewide emergency departments.¹ The data collection dates covered January 1–May 31 for each year. In this study, only symptoms related to influenza were included: upper respiratory infection, fever, influenza-like illness, cough and self-reported influenza. Correlation values were calculated using MS Excel software. **Results:** There were no breaks in confidentiality. All of the correlation values had a correlation value of 0.5 or better (Fig. 1), showing good correlation with the NC DHHS ILI data for both years; however, the more severe 2017–2018 influenza season had correlation values of ≥ 0.7 for all symptoms, versus 0.52–0.59 for URI and ILI, respectively, only during 2018–2019. **Conclusions:** The higher-severity influenza season did correlate with a higher r values when compared to North Carolina's DHHS ILI emergency department data than did the influenza season of moderate severity. A possible explanation is that a higher-severity influenza season would correlate better than a moderate influenza season because it shows fewer ILI peaks and troughs. In conclusion, the SEOL system served as an early warning that influenza is present among our staff, and it correlates well with the state system for ILI surveillance. Potential




Fig. 1.

limitations of SEOL are that respiratory symptoms are not specific to influenza; thus, they are subject to variation due to other respiratory viruses circulating among our employees.

1. The North Carolina Disease Event Tracking and Epidemiologic Collection Tool. NC DETECT website. <http://www.ncdetect.org>. Accessed Nov 8, 2019.

Funding: None

Disclosures: None

Doi:10.1017/ice.2020.772

Presentation Type:

Poster Presentation

Evaluation of Antibiotic Prophylaxis and Postsurgery Antibiotics for Urological Surgeries at an Academic Medical Center

Emily Drwiega, Grady Health System; Saira Rab, Grady Health System; Sheetal Kandiah, Emory University; Jane Kriengkauykiat, Grady Health System; Jordan Wong, Grady Health System

Objective: The purpose of this study was to evaluate antibiotic use in patients undergoing urological procedures. **Methodology:** This single-center, IRB-approved, retrospective, observational study was conducted at Grady Health System. Patients were included if they underwent their first inpatient urologic procedure between April 1, 2016, and April 1, 2018. Patients were excluded if they were <18 years old, pregnant, or a prisoner. The primary outcome was percentage of overall adherence to our institutional guidelines for surgical prophylaxis as a composite of antibiotic selection, dose, preoperative timing, and postoperative duration. Secondary outcomes include individual components of the composite outcome, nephrotoxicity, *Clostridium difficile* infection, and discharge antibiotic prescriptions. Descriptive statistics were used. **Results:** Of

the 100 patients evaluated, 11% achieved adherence with the primary outcome. Of the 89 patients who did not achieve composite outcome, only 8 selected the appropriate perioperative antibiotic. Overall, 30% were dosed appropriately, 47% were administered at the appropriate time with respect to time of incision, and 46% received perioperative antibiotics for no more than 24 hours. Also, 19 patients did not receive perioperative antibiotics. Overall, 14 different perioperative antibiotic regimens were utilized, despite institutional guidelines recommending 1 of 3 options. All 9 patients who developed nephrotoxicity received noncompliant perioperative prophylaxis. No patient developed *Clostridium difficile* infection within 30 days of surgery. Moreover, 58 patients were discharged with a prescription for at least 1 antibiotic. **Conclusions:** Most perioperative antibiotic prophylaxes for genitourinary surgeries are not compliant with institution guideline recommendations. Despite having institutional guidelines, there was a large variety in the antibiotic regimens that patients received. All of the patients identified as having an evaluated antibiotic-related adverse effect did not receive appropriate perioperative antibiotic prophylaxis. More than half of the patients received a prescription at discharge for at least 1 antibiotic.

Funding: None

Disclosures: None

Doi:10.1017/ice.2020.773

Presentation Type:

Poster Presentation

Evaluation of Environmental Decontamination of Surfaces Using Continuous Application of Low-Level Hydrogen Peroxide

Robert Garcia, Stony Brook University Hospital