

Discovery of a strong magnetic field in the rapidly rotating B2Vn star HR 7355

Mary E. Oksala^{1,2}, Gregg A. Wade², Wagner L. F. Marcolino^{3,4}, Jason H. Grunhut^{2,5}, David Bohlender⁶, Nadine Manset⁷, Richard H. D. Townsend⁸, and the MiMeS Collaboration

¹Department of Physics and Astronomy, University of Delaware, Newark, DE, USA

²Department of Physics, Royal Military College of Canada, Kingston, Ontario, Canada

³LAM-UMR 6110, CNRS & Univ. de Provence, Marseille, France

⁴Observatório Nacional, Rio de Janeiro, Brazil

⁵Department of Physics, Queen's University, Kingston, Ontario, Canada

⁶National Research Council of Canada, Herzberg Institute of Astrophysics, Victoria, Canada

⁷Canada-France-Hawaii Telescope Corporation, Kamuela, HI, USA

⁸Department of Astronomy, University of Wisconsin-Madison, Madison, WI, USA

Abstract. We report on the detection of a strong, organized magnetic field in the helium-variable early B-type star HR 7355 using spectropolarimetric data obtained with ESPaDOnS on CFHT by the MiMeS large program. We also present results from new V-band differential photometry obtained with the CTIO 0.9m telescope. We investigate the longitudinal field, using a technique called Least-Squares Deconvolution (LSD), and the rotational period of HR 7355. These new observations strongly support the proposal that HR 7355 harbors a structured magnetosphere similar to that in the prototypical helium-strong star, σ Ori E.

Keywords. stars: magnetic fields, stars: rotation, stars: early-type, stars: circumstellar matter, stars: individual (HR 7355), techniques: polarimetric

1. Introduction

HR 7355 (HD 182180) is a bright B2Vn helium-strong star originally classified as a Be star due to H α emission present in its spectrum (Abt & Cardona 1984). Previous studies of this star show a $v \sin i \sim 300$ km s⁻¹ (Abt *et al.* 2002) with a $P_{\text{rot}} \sim 0.52$ d (Koen & Eyer 2002), as well as variation in helium, H α , and brightness, suggesting the presence of a magnetosphere (Rivinius *et al.* 2008). HR 7355 is the most rapidly rotating helium-strong star, rotating near its critical velocity, providing an excellent testbed for magnetospheres under the effects of rapid rotation.

2. Method

Least-Squares Deconvolution (LSD) describes the stellar spectrum as the convolution of a mean Stokes I or V profile, representative of the average shape of the line profile, and a line mask, describing the position, strength and magnetic sensitivity of all lines in the spectrum. From the LSD mean Stokes I and V profiles, we calculate the longitudinal magnetic field, B_{ℓ} :

$$B_{\ell} = -2.14 \times 10^{11} \frac{\int vV(v)dv}{\lambda gc \int [1 - I(v)]dv} \quad (2.1)$$




Figure 1. Top: Longitudinal magnetic field measurements for HR 7355 and the best-fit first order sine curve. Oksala *et al.* (2010) (asterisks) and Rivinius *et al.* (2010) (diamonds) with 1σ error bars. Bottom: The V-band photometric light curve for HR 7355 including both HIPPARCOS photometry (asterisks) and new CTIO data (plus signs).

(Wade *et al.* 2000), where λ is the average wavelength and g is the average Landé factor in the mask. I_c is the continuum value of the intensity profile. The integral is evaluated over the full velocity range of the mean profile.

3. Results

We detect a strong magnetic field on HR 7355, the most rapidly rotating helium-strong star discovered thus far. A simultaneous independent confirmation of the field detection has been obtained with FORS at the VLT by Rivinius *et al.* 2010. The longitudinal magnetic field varies sinusoidally with the rotation period, with extrema -2 to 2.5 kG. Assuming a dipole magnetic field, the polar value of the magnetic field is ~ 13 -17 kG. The photometric (brightness) light curve constructed from HIPPARCOS archival data and new CTIO measurements shows two minima separated by 0.5 in rotational phase and occurring 0.25 cycles before/after the magnetic extrema. Using the Scargle periodogram, eclipse-like photometric variations give a highly precise $P_{rot} = 0.5214404(6)$ days. We confirm spectral variability of helium and metal lines, as well as variability of $H\alpha$ emission. $H\alpha$ emission indicates circumstellar material extending out to $5 R_*$ from the star, rotating rigidly with the stellar surface. We conclude that HR 7355 is a magnetic oblique rotator with a magnetosphere, mirroring the physical picture for σ Ori E (Townsend *et al.* 2005).

References

- Abt, H. A. & Cardona, O. 1984, *ApJ*, 285, 190
 Abt, H. A., Levato, H., & Grosso, M. 2002, *ApJ*, 573, 359
 Koen, C. & Eyler, L. 2002, *MNRAS*, 331, 45
 Oksala, M. E., Wade, G. A., Marcolino, W. L. F., Grunhut, J. *et al.* 2010, *MNRAS*, 405, 51
 Rivinius, T., Štefl, S., Townsend, R. H. D., & Baade, D. 2008, *A&A*, 482, 255
 Rivinius, T., Szeifert, T., Barrera, L., Townsend, R. H. D. *et al.* 2010, *MNRAS*, 405, 46
 Townsend, R. H. D., Owocki, S. P., & Groote, D. 2005, *ApJ* (Letters), 630, 81
 Wade, G. A., Donati, J.-F., Landstreet, J. D., & Shorlin, S. L. S. 2000, *MNRAS*, 313, 851