

Neon abundances in three Wolf-Rayet stars observed with the ISO Short Wavelength Spectrometer

Patrick W. Morris¹ and Karel A. van der Hucht

SRON, Sorbonnelaan 2, NL-3584 CA Utrecht, the Netherlands

Allan J. Willis, Luc Dessart, and Paul A. Crowther


*Department of Physics & Astronomy, University College London,
Gower St, London WC1E 6BT, UK*

Peredur M. Williams

Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, UK

1. Introduction

We are motivated to study the Ne/He abundance ratio at the surfaces of Wolf-Rayet stars by evolutionary models, which have predicted enhancements by more than an order of magnitude over cosmic levels during the WC stage (*e.g.*, Maeder 1991). This should occur early in the WC phase, as core-hydrogen is exhausted and the CNO bi-cycle gives way to the following chain of reactions for the Galactic stars:



The latest evolutionary models, which include the effects of interior mixing due to stellar rotation, indicate that the enhancement may be somewhat more gradual, but that the Ne/He ratio is essentially unchanged in its peak value, coinciding with complete H-depletion (*e.g.*, Meynet, these Proceedings).

2. Latest ISO results

The first WC star to be studied in detail, γ^2 Vel (WR 11, WC8+O8.5III), showed only a minor enhancement of Ne/He (Barlow *et al.* 1988). This was the only WC star for which an *IRAS-LRS* spectrum was available to measure the [NeIII] 15.55 μm line emission. With *ISO*, γ^2 Vel and several other WR stars have been observed using the short wavelength spectrometer (SWS). The reader is referred to Figure 4 in the review by A. Willis (these Proceedings) for a montage of several WR star spectra over the 12–17 μm range that would include the lines of [NeII] 12.81 μm , [NeV] 14.32 μm , and [NeIII] 15.55 μm .

The SWS and ground-based spectrum of WR 146 (WC6+O) has been studied by Willis *et al.* (1997), who found Ne/He to be much closer to the evolutionary

¹Present address: Astronomical Institute, University of Amsterdam, the Netherlands

predictions. Subsequently, Morris *et al.* (1998) presented preliminary results for WR 135 (WC8) and WR 147 (WN8(h)+B0.5V), and re-assessed the results for γ^2 Vel in light of the new distance established by *Hipparcos* (van der Hucht *et al.* 1997) and the mass-loss rate estimates by St-Louis *et al.* (1988) and Stevens *et al.* (1996).

2.1. WR 135 (WC8)

The Ne/He ratio of WR 135 is roughly a factor four higher than expected for an environment where H/He = 0.0 and C/He = 0.13, but this is not as much as the evolutionary models predict. There is some hint of [MgV] 13.54 μm emission, which could indicate that α capture by ^{22}Ne has occurred. We do not notice magnetic dipole transitions of [MgV] 5.608 μm or [MgIV] 4.488 μm .

2.2. WR 11 (WC8+O8.5III)

The Ne-line fluxes at 12.81 μm and 15.55 μm are in good agreement with the values quoted by Barlow *et al.* (1988), but the stellar distance is now known to be nearly a factor of two closer based on *Hipparcos* observations. This by itself does not influence the results much, since the ion fraction γ_i scales as $d^2/\dot{M}^{3/2}$, and \dot{M} scales as $d^{3/2}$. However, much lower mass-loss rates are given by polarimetry studies by St-Louis *et al.* (1988) and X-ray observations by Stevens *et al.* (1996), roughly three times lower than estimated by Barlow *et al.* (1988) from the radio free-free continuum. Using an average $\dot{M} = 1.3 \times 10^{-4} M_{\odot}\text{yr}^{-1}$ and C/He = 0.14 from recombination analysis of the SWS lines, a lower value of Ne/He $\geq 6.3 \times 10^{-3}$ is obtained, a factor of ~ 10 higher than a cosmic environment where H/He = 0.0 and C/He = 0.14.

2.3. WR 147 (WN8(h)+B0.5V)

The SWS spectrum of WR 147 shows very strong [NeIII] 15.55 μm emission, and no [NeII] 12.81 μm line (*e.g.*, Morris *et al.* 1998). Nonetheless, Ne in the ratio with He should be normal, since no net Ne production occurs during the CNO bi-cycle. To check Ne/He, we relied on the mass-loss rate estimated by Hamann *et al.* (1995) from optical spectroscopy. Their value was re-scaled to the distance and interstellar reddening given by Churchwell *et al.* (1992), using $M_v = v - \text{DM} - A_v$, $A_v = 4.1 E_{b-v}$, and equations (4) and (5) in Hamann *et al.* (1993). This leads to a lower limit of Ne/He $\geq 1.0 \times 10^{-3}$, which is a factor of 2 or 3 higher than expected for an environment where H/He is somewhere between 0.0 and 0.4. With respect to levels predicted for WC stars, the enhancement is hardly that, but it is significant for a WN star, and may indicate some exposure of core-processed material via rotational mixing, as predicted to occur in WR stars (Meynet, these Proceedings).

3. The \dot{M} problem

As mentioned, the Ne-ion fraction scales as $\dot{M}^{-3/2}$ in the two-level atom, mass-loss normalizing method by Barlow *et al.* (1988). Among published results using spherical, homogeneous wind models, differences by a factor of three in the value of \dot{M} are not unusual due to distance and (to a lesser extent) reddening

uncertainties, and the inclusion of line-blanketing by metals. More important may be the assumptions on geometry and homogeneity of the wind. It is known that clumping may lower the mass-loss rate by factors of two to three, which would serve to increase the ion fractions *in the clumps*, according to the inverse scaling with \dot{M} . This is emphasized by the case of WR 11, and might rectify the low Ne/He abundance ratio in WR 135. It would normally, however, exacerbate an already unexpected (albeit minor at the moment) over-abundance in the WN8 star of WR 147. But the system is highly reddened at $A_V \simeq 11.5$ mag (Churchwell *et al.* 1992), and the foreground extinction properties are probably anomalous, contributing to the difficulty in arriving at a clean spectroscopic solution for the physical parameters of the system. These issues are addressed in papers in preparation, where we will present the ISO-SWS spectra of WR 11, WR 135, and WR 147 in detail.

References

- Barlow, M.J., Roche, P.F., Aitken, D.K. 1988, MNRAS 232, 821
 Churchwell, E., Bieging, J.H., van der Hucht, K.A., Williams, P.M., Spoelstra, T.A.Th., Abbott, D.C. 1992, ApJ 393, 329
 Hamann, W.-R., Koesterke, L., Wessolowski, U. 1993, A&A 274, 397
 Hamann, W.-R., Koesterke, L., Wessolowski, U. 1995, A&A 299, 151
 van der Hucht, K. A., Schrijver, H., Stenholm, B., *et al.* 1997, New Astron. 2, 245
 Maeder, A. 1991, A&A 242, 93
 Morris, P.W., van der Hucht, K.A., Willis, A.J., Williams, P.M. 1998, in: L.B.F.M. Waters, C. Waelkens, K.A. van der Hucht & P.A. Zaal (eds.), *ISO's View on Stellar Evolution*, ApSS 255, 157
 St-Louis, N., Moffat, A.F.J., Drissen, L., *et al.* 1988, ApJ 330, 286
 Stevens, I.R., Cocoran, M.F., Willis, A.J., *et al.* 1996, MNRAS 283, 589
 Willis, A.J., Dessart, L., Crowther, P.A., Morris, P.W., Maeder, A., Conti, P.S., van der Hucht, K.A. 1997, MNRAS 290, 371

