

Disorders, Psychiatric Hospital ⁴ *Department for PTSD Treatment, Psychiatric Hospital*

Within its sociotherapeutical programme St. John Psychiatric Hospital has been continuously since 1963 conducting the treatment and rehabilitation of psychiatric patients accommodated in heterogeneous families according to the Belgian model. Research conducted in the West as well as the experience gained at our hospital suggests that this sociotherapeutical method is very effective in the process of prolonged treatment, rehabilitation and resocialisation as well as the improvement of the quality of life of psychiatric patients. The ultimate objective of this investigation was to examine the quality of life of the groups of schizophrenic and depressive patients under observation, and the impact of a heterogeneous family, primary or secondary family and hospital ambient on the process of treatment and rehabilitation. The assumption is that in the process of prolonged treatment patients accommodated in heterogeneous families psychosocially function better and attain a higher quality of life than those who are under long-term outpatient treatment and live with the primary or secondary family or, alternatively, have been hospitalised. Applied was a comparative method of investigating groups of patients affected by schizophrenia and depression that were in the process of prolonged treatment accommodated in heterogeneous families, primary or secondary family, or those who were hospitalised. Individually assessed by way of appropriate questionnaires, completed by the very subjects, were the health-related quality of life and the subjective quality of life.

P417

Socio-psychological problems in the families of patients with mental illnesses in republic of belarus

K. Mironova. *Department of Psychiatry, Belarusian Medical Academy of Post-Graduate Education, Minsk, Belarus*

The study was carried out at the Department of Psychiatry of the Belarusian Medical Academy of Post-Graduate Education.

The purpose of the study was investigating the problems connected with negative consequences of mental illness stigma and the family burden.

Method and objective: Study participants were: the caregivers of the patients admitted to the Republican Clinical Psychiatric Hospital, suffering from schizophrenia, affective disorders and the control group (participants from mentally healthy families).

The inclusion criteria were: 1) living with the patient in the same house; 2) age > 18 years; 3) absence of somatic and psychiatric pathology; 4) informed consent.

We used the following assessment instruments: the Medical Outcomes Study Short-Form36 (SF-36) survey; Family Experiences with the Stigma of Mental Illness Questionnaire by H.Stuart; the Coping Strategies Questionnaire by E.Heim; the Questionnaire of Family Burden according to the WHO «Quality Assurance in Mental Health Care».

Result: The obtained results show that relatives of mentally ill patients suffer from negative consequences of stigma and discrimination. They have the burden in economic, psychological, social environment, family interaction areas. Relatives feel social isolation, shame, guilt, worry and distress.

Conclusion: One of the causes of such tragic consequences of stigma is the lack of specific information about mental illnesses and choice of inadequate coping strategies. We apply

psychoeducational project in the families of patients with schizophrenia to reduce negative influence on their lives.

P418

Psychiatrist and social dialogue

E. Koic ¹, V. Djordjevic ², S. Nadj ¹, E.N. Gruber ³, D. Poredos ³.
¹ *Psychiatric Department, General Hospital Virovitica, Virovitica, Croatia* ² *Psychiatric Clinic, Clinical Hospital Center Zagreb, Zagreb, Croatia* ³ *Neuropsychiatric Hospital "Dr.Ivan Barbot", Popovaca, Croatia*

Background and aims: The authors presented their activities in prevention of mobbing, suicidality, pathological gambling, and antistigma programs aimed for the persons treated for combat PTSD.

Methods: Social dialogue is achieved through numerous outpatient activities that included systematic information and education, public discussions, conducted studies, public program promotions, performance presentations, koordination, and active engagement of various professionals (psychiatrists, psychologists, pedagogues, social workers, jurists, politicians, employees in civil service, associations and syndicate, employers, and volunteers).

Results: Numerous public discussions and professional gatherings have been held, professional and scientific studies have been conducted, programs have been presented in public through various media, web-pages have been created and brochures have been written. The created positive atmosphere leveled up the awareness, what resulted in propositions for drawing up broader, national strategies, and some pending legal solutions.

Conclusion: Social dialogue is represented by all types of information exchange, consultations, and discussions between social partners and government representatives about issues of common interest related to social or economic policy. In situations when some primarily social events partially participate in the genesis of psychological disorders, a psychiatrist may take part in the dialogue, pointing out the existence of the problem, defining the problem and offering professional solutions for prevention. In order to have a social dialogue it is necessary to have most extensive public support and understanding that only with joint endeavor of all citizens the problem may be, and must be, solved.

P419

An investigation on spouses' mutual abuse behavior

M. Nasiri. *Psychiatric Nursing Department, Isfahan, Iran*

Introduction: The family plays an essential role in individual and social life. That is the first center to grow and blossom the talents appropriately and a starting point for all social changes each modification should start from the family.

Method: This is a descriptive analytic study. On 394 men and 404 women. The spouses' abuse behavior was defined by a questionnaire. Results:

The findings showed that there was a significant association between abuse behaviors such as aggression, screaming, punishment and physical fight, mutual scolding, contempt and criticizing, silence and not speaking, insulting not talking to each other, criticizing in public, throwing and breaking things, sleeping separately between men and women ($p < 0.001$)

The findings showed 88% of the men and 51% of the women, stubbornness in 73% of men and 70% of women, aggression and screaming in 63% of men and 51% women, physical punishment