

## MATTER-BOUNDED PHOTOIONIZED CLOUDS


L. BINETTE<sup>1</sup>, A.S. WILSON<sup>2</sup>, T. STORCHI-BERGMANN<sup>3</sup>

<sup>1</sup> *Observatoire de Lyon - 9 av. Charles André  
F-69561 Saint-Genis-Laval Cedex, France*


<sup>2</sup> *Astronomy Department - University of Maryland  
College Park, MD 20742, USA*

<sup>3</sup> *Instituto de Fisica - UFRGS, Campus do Vale  
91500 Porto Alegre, RS, Brasil*

The *extended* ionized gas in Seyfert and Radio-Galaxies is characterized by large values of the ratio  $\text{He II}/\text{H}\beta$ , which exceeds the value predicted by the standard photoionization model in which the ionizing continuum consists of a power-law. This has led to the suggestion of considering a matter-bounded (MB) component [3],[5],[2] for explaining such extreme values. We now find[1] that it is also possible to resolve the temperature problem[3] if the thickness and the ionization parameter of the MB is appropriately selected. Adopting a canonical power law ( $\alpha=-1.3$ ) and solar abundances ( $Z=1$ ), we can account for the observed trends in excitation (represented for example by the ratio  $[\text{O II}]/[\text{O III}]$  in Fig. 1)) by varying the relative number of MB clouds (which emit the high excitation lines C IV, [Ne V], He II... and most of [O III]) versus the number of ionization-bounded (IB) clouds (which emit [N II],[S II] [O II], [O I]...). We obtain a one-parameter sequence (solid line) which is function of the weight  $A_{\text{M/I}}$  of the MB component relative to the IB component. This  $A_{\text{M/I}}$ -sequence successfully reproduces the observed range in  $\text{He II}/\text{H}\beta$ . Note the failure of the traditional U-sequence (long dashed line). Fig. 2 indicates that we can also reproduce the ratio  $R_{\text{OIII}} = [\text{O III}]\lambda 4363/[\text{O III}]\lambda 5007$  and therefore resolve the temperature problem. Interestingly, our model indicates a temperature difference of 5 000 K between the IB component ([N II] temperature  $\simeq 10\,000$  K) and the MB component ([O III] temperature  $\simeq 15\,000$  K) while the traditional U-sequence predicts a difference of only 1 000 K. Such difference of 5 000 K has been reported[4] in the extended gas of Cygnus A.



**Figure 1.** Diagram of the line ratios  $[O II]/[O III]$  against  $He II/H\beta$ . Filled and open symbols denote Seyfert and radio galaxies, respectively. Larger symbols denote the nuclear values. A dotted line joins measurements at different locations in the same galaxy. The parameter  $A_{M/I}$  of our model represents the relative weight of the MB component and increases from left to right along the solid line ( $0.04 \leq A_{M/I} \leq 16$ ). The long dashed line represents the traditional U-sequence.



**Figure 2.** Diagram of the line ratios  $R_{OIII}$  ( $4363\text{\AA}/5007\text{\AA}$ ) against  $He II/H\beta$ . The symbols have the same meaning as in Fig. 1.  $A_{M/I}$  is increasing from left to right. The two short-dash lines correspond to metallicity sequences ( $0.2 \leq Z \leq 2.0$ ) at  $A_{M/I}=0.4$  and  $4.0$ .

## References

- [1] Binette, L., Wilson, A. S. and Storchi-Bergmann, T., 1996. *A&A*, submitted.
- [2] Morganti, R., Robinson, A., Fosbury, R. A. E., di Serego Alighieri, S., Tadhunter, C. N., and Malin D. F., 1991. *MNRAS*, **249**, 91.
- [3] Tadhunter, C.N., Robinson, A. and Morganti, R., 1989. ESO Conf. and Workshop Proc. no.32, Garching, p 293.
- [4] Tadhunter, C. N., Metz, S. and Robinson, A., 1994. *MNRAS*, **268**, 989.
- [5] Viegas, S. M. and Prieto, A., 1992. *MNRAS*, **258**, 483.